B.M.
E.G.
27.c.
0.
B.

Meyrick Types of Microlepidoptera in the British Museum (Natural History) London

Catalogue of the Type Specimens of Microlepidoptera in the British Museum (Natural History) described by Edward Meyrick

BY

J. F. Gates Clarke

Bureau of Entomology and Plant Quarantine
U.S. Department of Agriculture

VOLUME II

STENOMIDAE
XYLORYCTIDAE
COPROMORPHIDAE

LONDON
PRINTED BY ORDER OF THE TRUSTEES
OF THE BRITISH MUSEUM

Issued 1955

Price: Six pounds

595.782

Sold at The British Museum (Natural History), Cromwell Road, S.W.7
and by
B. Quaritch, Ltd.
and H.M. Stationery Office

MADE AND PRINTED IN GREAT BRITAIN
BY JARROLD AND SONS LTD, NORWICH

FAMILY STENOMIDAE

FAMILY XYLORYCTIDAE . . . see p. 397
FAMILY COPROMORPHIDAE . . see p. 511















ANADASMUS

ANADASMUS Walsingham

Anadasmus accurata (Meyrick), new combination

Plate 1, Figures 1–1a, 2–2a

Stenoma accurata Meyrick, 1916, Exotic Microlepidoptera, 1: 526.
Stenoma perjura Meyrick, 1925, Exotic Microlepidoptera, 3: 207 (new synonymy).

[accurata]

"♂. 19–20 mm. . . . French Guiana, St. Jean, R. Maroni, in July; two specimens."

Lectotype of *accurata:* ♂, 19 mm., with "LeM. .7.15" at the bottom of the pin-label. There is a median-ventral hair-pencil from the first and second abdominal sternites. Slide No. 5315.

Figure 1, left wings; 1a, ventral aspect of male genitalia with aedeagus *in situ.*

[perjura]

"♂. 21 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex."

Type of *perjura:* The above indicated specimen dated "1.20". Slide No. 5316.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ.*

Anadasmus arenosa (Meyrick), new combination

Plate 1, Figures 3–3a

Stenoma arenosa Meyrick, 1916, Exotic Microlepidoptera, 1: 519.

"♂. 30 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. This species is very closely related to *Anadasmus leontodes* (Meyrick, q.v.) and *Anadasmus quadratella* (Walker) (new combination). Slide No. 5205.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ.*

Anadasmus byrsinitis (Meyrick), new combination

Plate 1, Figures 4–4b

Stenoma byrsinitis Meyrick, 1912, Trans. Ent. Soc. London, 1911: 713.

"♂♀. 25–30 mm. . . . Colombia, San Antonio, 5,800 feet, in November and December; nine specimens."

Lectotype: ♂, dated ".12.07". Slide No. 5284.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Anadasmus capnocrossa (Meyrick), new combination

Plate 2, Figures 1–1b

Stenoma capnocrossa Meyrick, 1925, Exotic Microlepidoptera, 3: 197.

"♂. 25 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex."

Type: The above indicated specimen dated "1.20". Slide No. 5206.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Anadasmus chionodora (Meyrick), new combination

Plate 2, Figures 2–2a

Stenoma chionodora Meyrick, 1915, Exotic Microlepidoptera, 1: 464.

"♂♀. 25–27 mm. . . . British Guiana, Mallali, in March (*Parish*); three specimens."

Lectotype: ♂, dated ".3.13". The three original specimens have identical data. Slide No. 5273.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Anadasmus dictyogramma (Meyrick), new combination

See Anadasmus venosella (Walker), p. 11.

Anadasmus endochra (Meyrick), new combination

Plate 2, Figures 3–3a

Stenoma endochra Meyrick, 1925, Exotic Microlepidoptera, 3: 198.

"♂. 23 mm. . . . Brazil, Teffé, in January (*Parish*); 2 ex."

Lectotype: One of the above males, both of which bear identical data and are dated "1.20". Slide No. 5211.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Anadasmus entephras (Meyrick), new combination

Plate 2, Figures 4–4b

Stenoma entephras Meyrick, 1915, Exotic Microlepidoptera, 1: 453.

"♂. 27 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated ".3.13". Slide No. 5323.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANADASMUS



ANADASMUS

Anadasmus incitatrix (Meyrick), new combination

Plate 3, Figures 1–1a

Stenoma incitatrix Meyrick, 1925, Exotic Microlepidoptera, 3: 207.

"♂. 28 mm. . . . Argentina, R. Parana; 1 ex. Allied to *pleurotricha*."

Type: The above indicated specimen. At the bottom of the pin-label appears "LeM. .15". Slide No. 5291.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Anadasmus ischioptila (Meyrick), new combination

Plate 3, Figures 2–2b

Stenoma ischioptila Meyrick, 1925, Exotic Microlepidoptera, 3: 211.

"♂. 30 mm. . . . Colombia, Sosomoko, 2,650 feet, February; 1 ex."

Type: A single male, measuring exactly 30 mm., and agreeing in all details with Meyrick's description and date, is in his collection. There appears little doubt that this is the actual type, but the data on the pin-label read, "S. Antonio, Colombia, 6,600′ .2.20." and I suspect a *lapsus* was responsible for the recorded data.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Anadasmus leontodes (Meyrick), new combination

Plate 3, Figures 3–3a

Gonioterma leontodes Meyrick, 1915, Exotic Microlepidoptera, 1: 384.

Stenoma quadratella (Walker) Meyrick (*not* Walker), 1925, Exotic Microlepidoptera, 3: 192.

"♂. 28 mm. . . . Dutch Guiana, Paramaribo, in December; one specimen."

Type: The above indicated specimen. Slide No. 5632.

I have compared Meyrick's *leontodes* with Walker's type. The latter is a female, and is clay-coloured, whereas Meyrick's species is reddish-brown. A male, from which I have made a genitalic preparation, which matches Walker's *quadratella* is easily distinguished from Meyrick's *leontodes* by the cluster of small stout transverse cornuti and the long cylindrical spine-covered distal portion of the vesica. The vesica of *leontodes* is armed with one large cornutus and its distal portion is covered with concentric bands of fine spiculate cornuti.

For the above reasons I remove *leontodes* from synonymy.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Anadasmus lithogypsa (Meyrick), new combination

Plate 3, Figures 4–4b

Stenoma lithogypsa Meyrick, 1932, Exotic Microlepidoptera, 4: 299.

"♂. 18 mm. . . . Brazil, Pará, July (*Parish*); 1 ex."

Type: The above indicated specimen dated "7.–19". Slide No. 5308.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Anadasmus obmutescens (Meyrick), new combination

Plate 4, Figures 1–1b, 2–2b

Stenoma obmutescens Meyrick, 1916, Exotic Microlepidoptera, 1: 525.
Stenoma pleurotricha Meyrick, 1925, Exotic Microlepidoptera, 3: 206 (new synonymy).

[obmutescens]

"♂. 31 mm. . . . French Guiana, St. Jean, R. Maroni; one specimen."

Type: The male referred to above. At the bottom of the pin-label is the inscription, "LeM. .15". Slide No. 5292.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

[pleurotricha]

"♂. 29–30 mm. . . . Brazil, Teffé, January (*Parish*); 3 ex."

Lectotype: One of the above indicated males dated "1.20". One specimen is missing. Slide No. 5289.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Anadasmus paurocentra (Meyrick), new combination

Plate 4, Figures 3–3b

Stenoma paurocentra Meyrick, 1912, Trans. Ent. Soc. London, 1911: 713.

"♂. 28–34 mm. . . . Colombia, San Antonio, 4,400–5,800 feet, in November; two specimens."

Lectotype: The male labelled, "San Antonio, W. Colombia, R. 5,800′ .11.07". Slide No. 5296.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Anadasmus pelinitis (Meyrick), new combination

Plate 4, Figures 4–4b

Stenoma pelinitis Meyrick, 1912, Trans. Ent. Soc. London, 1911: 714.

"♂. 34–38 mm. . . . Colombia, San Antonio, 5,800 feet, in November and December; eleven specimens."

Lectotype: A male dated ".12.07". Meyrick records only males, but there are three females in the type series. Slide No. 5295.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANADASMUS


1

1a

1b

2

2a

2b


3

3a

3b


4

4a


4b

ANADASMUS

Anadasmus plebicola (Meyrick), new combination

Plate 5, Figures 1–1b

Stenoma plebicola Meyrick, 1918, Exotic Microlepidoptera, 2: 205.

"♂♀. 26–28 mm. . . . French Guiana, R. Maroni; British Guiana, Bartica, January (*Parish*); five specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana, LeM. .15". Two specimens are missing. There is a median-ventral hair-pencil from the first and second sternites. Slide No. 5322.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Anadasmus pleurotricha (Meyrick), new combination

See Anadasmus obmutescens (Meyrick), p. 8.

Anadasmus pleximorpha (Meyrick), new combination

See Anadasmus venosella (Walker), below.

Anadasmus porinodes (Meyrick), new combination

Plate 5, Figures 2–2b

Stenoma porinodes Meyrick, 1915, Exotic Microlepidoptera, 1: 454.

"♂ 22 mm., ♀ 24 mm. . . . British Guiana, Mallali, in March (*Parish*); Peru, Chanchamayo, in June; two specimens."

Lectotype: ♂, "Chanchamayo, Peru, M. .1.13". The June date as recorded by Meyrick is a misprint. Slide No. 5326.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Anadasmus venosella (Walker), new combination

Plate 5, Figures 3–3b

Cryptolechia venosella Walker, 1864, List of the specimens of lepidopterous insects in the collection of the British Museum, 29: 718.

Stenoma pleximorpha Meyrick, 1930, Exotic Microlepidoptera, 4: 32 (new synonymy).

Stenoma dictyogramma Meyrick, 1932, Exotic Microlepidoptera, 4: 301 (new synonymy).

[pleximorpha]

"♂. 25 mm. . . . Bolivia, 1 ex. (Coll. Paravicini)."

Type: The above indicated male, "Prov. Cochabamba. Y. d. Espiritu Santo. P. Germain." A small white label bears the inscription, "M829". The specimen is also marked "Holotype". Slide No. 5335. [*continued on p. 140.*

ANTAEOTRICHA Zeller

Antaeotricha acrobapta Meyrick

Plate 6, Figures 1–1c

Antaeotricha acrobapta Meyrick, 1933, Exotic Microlepidoptera, 4: 434.

"♀. 22 mm. . . . Argentina, Alta Gracia (*C. Bruch*); 1 ex. Nearest *oxycentra* Meyr."

Type: The above indicated specimen with "CB, .32." at the bottom of the pin-label. Slide No. 4862.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, ventral view to show ostium and genital plate; 1c, signum.

Antaeotricha acrograpta (Meyrick)

Plate 6, Figures 2–2b

Aphanoxena acrograpta Meyrick, 1915, Exotic Microlepidoptera, 1: 387.

"♂. 16 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen."

Type: The above indicated specimen dated "1.13". Slide No. 4815.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha acronephela Meyrick

Plate 6, Figures 3–3b

Antaeotricha acronephela Meyrick, 1915, Exotic Microlepidoptera, 1: 392.

"♂. 18 mm. . . . British Guiana, Bartica and Mallali, in December and March (*Parish*); two specimens."

Lectotype: ♂, "Mallali, Brit. Guiana, Parish. 3.13". Slide No. 4806.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha actista (Meyrick), new combination

Plate 6, Figures 4–4b

Stenoma actista Meyrick, 1913, Trans. Ent. Soc. London, 1913: 186.

"♀. 33–34 mm. . . . Venezuela, Palma Sola; British Guiana, R. Demerara; two specimens."

Lectotype: ♀, "Palma Sola, Venezuela, D. .06". Slide No. 5113.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, enlarged view of ostium and genital plate.


1
1a
1b
1c
2
2a
2b
3
3a
3b
4
4a
4b



ANTAEOTRICHA

Antaeotricha adornata (Meyrick), new combination

Plate 7, Figures 1–1b

Stenoma adornata Meyrick, 1915, Exotic Microlepidoptera, 1: 443.

"♂. 18 mm. . . . Peru, Pacaya, in September (*Mounsey*); one specimen."

Type: The above indicated specimen dated ".9.12". Slide No. 5385.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha aequabilis (Meyrick), new combination

Plate 7, Figures 2–2a

Stenoma aequabilis Meyrick, 1916, Exotic Microlepidoptera, 1: 513.

"♂♀. 23–24 mm. . . . French Guiana, St. Jean and Godebert, R. Maroni; five specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana, LeM. .15". Slide No. 4984.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha aggravata (Meyrick), new combination

Plate 7, Figures 3–3b

Stenoma aggravata Meyrick, 1916, Exotic Microlepidoptera, 1: 514.

"♂. 16 mm. . . . French Guiana, R. Maroni; one specimen. Notable by the peculiar antennae."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5147.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha aglypta Meyrick

Plate 7, Figures 4–4b

Antaeotricha aglypta Meyrick, 1925, Exotic Microlepidoptera, 3: 174.

"♂. 14–16 mm. . . . Brazil, Teffé, December, January (*Parish*); 4 ex."

Lectotype: One of the above specimens dated "1.20". Slide No. 4941.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha agrioschista (Meyrick), new combination

Plate 8, Figures 1–1b

Stenoma agrioschista Meyrick, 1927, Exotic Microlepidoptera, 3: 365.

"♂♀. 20–21 mm. . . . Texas, Alpine, 5,000–8,000 feet, April to July; 8 ex. Nearest *humilis* but quite distinct."

Lectotype: ♂, dated ".4.26". The original eight specimens are in the Meyrick collection, all but one being labelled "Brenster Co. Texas" (=Alpine of Meyrick). Slide No. 5403.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha ampherista Meyrick

Plate 8, Figures 2–2b

Antaeotricha ampherista Meyrick, 1925, Exotic Microlepidoptera, 3: 163.

"♂. 24–27 mm. . . . Bolivia, Prov. del Sara, 1,500 feet; Brazil, Obidos, September; 3 ex."

Lectotype: ♂, "Prov. del Sara, Dep. Sta. Cruz, Bolivia; 450 m.". Slide No. 4833. The male genitalia of this and *dynastis* appear to be identical.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha amphilyta Meyrick

Plate 8, Figures 3–3b

Antaeotricha amphilyta Meyrick, 1916, Exotic Microlepidoptera, 1: 503.

"♂. 17–18 mm. . . . French Guiana, St. Jean, R. Maroni; five specimens."

Lectotype: One of the above indicated specimens with "LeM. .7.15" at the bottom of the pin-label. Slide No. 4927.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha anaclintris Meyrick

Plate 8, Figures 4–4b

Antaeotricha anaclintris Meyrick, 1916, Exotic Microlepidoptera, 1: 499.

"♂. 22–23 mm. . . . French Guiana, St. Jean, R. Maroni; nine specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana, LeM. 15". Slide No. 4825. Only four of the original specimens remain in the Meyrick collection.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha annixa Meyrick

See Antaeotricha dissimilis (Kearfott), p. 48.

Antaeotricha anticharis (Meyrick), new combination

See Antaeotricha tanysta (Meyrick), p. 127.



ANTAEOTRICHA


1


1a

2a

2


1b


2b

3

3a

3b


4

4a


4b

ANTAEOTRICHA

Antaeotricha aporodes Meyrick

Plate 9, Figures 1–1b

Antaeotricha aporodes Meyrick, 1915, Exotic Microlepidoptera, 1: 400.

"♂♀. 24–25 mm. . . . Dutch Guiana, Paramaribo, in December; four specimens. Allied to *binubila* and *addon*."

Lectotype: ♀, "Paramaribo, Dutch Guiana. E. .12.92". Three specimens are missing. Slide No. 4902.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum.

Antaeotricha arachnia (Meyrick), new combination

Plate 9, Figures 2–2b

Stenoma arachnia Meyrick, 1915, Exotic Microlepidoptera, 1: 429.

"♀. 24 mm. . . . British Guiana, Bartica, in December (*Parish*); one specimen."

Type: The specimen indicated above and dated ".12.12". Slide No. 5486.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum.

Antaeotricha arystis Meyrick

Plate 9, Figures 3–3b

Antaeotricha arystis Meyrick, 1915, Exotic Microlepidoptera, 1: 402.

"♂. 17–18 mm. . . . British Guiana, Bartica, in January (*Parish*); two specimens."

Lectotype: ♂, 18 mm., dated "1.13". Slide No. 4931.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha asphalopis (Meyrick), new combination

See Antaeotricha vannifera (Meyrick), p. 136.

Antaeotricha astynoma (Meyrick)

Plate 9, Figures 4–4b

Aphanoxena astynoma Meyrick, 1915, Exotic Microlepidoptera, 1: 388.

"♂. 18 mm. . . . British Guiana, Mallali, March (*Parish*); one specimen."

Type: The above indicated male dated ".3.13". Slide No. 4922.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha atmospora (Meyrick), new combination

Plate 10, Figures 1–1b

Stenoma atmospora Meyrick, 1925, Exotic Microlepidoptera, 3: 209.

"♀. 38–39 mm. . . . Colombia, Minero and Sosomoko, 2,650 feet, April, October; 2 ex."

Lectotype: ♀, "Minero, Colombia, .10.20". Slide No. 5114.

Figure 1, left wings; 1a, ventral aspect of female genitalia; 1b, ventral view to show detail of genital plate and ostium.

Antaeotricha axena Meyrick

Plate 10, Figures 2–2b

Antaeotricha axena Meyrick, 1916, Exotic Microlepidoptera, 1: 501.

"♂. 21–22 mm. . . . French Guiana, St. Jean, R. Maroni; four specimens."

Lectotype: One of the above indicated specimens with "LeM. .7.15" at the bottom of the pin-label. Slide No. 4918.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha balanocentra (Meyrick)

Plate 10, Figures 3–3b

Aphanoxena balanocentra Meyrick, 1915, Exotic Microlepidoptera, 1: 387.

"♂♀. 16–19 mm. . . . British Guiana, Bartica and Mallali, from December to March (*Parish*); eight specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. .2.13". Slide No. 4814.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha ballista (Meyrick)

Plate 10, Figures 4–4b

Stenoma ballista Meyrick, 1916, Exotic Microlepidoptera, 1: 516.

"♂ 19–20 mm., ♀ 24 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana, LeM. .16". Slide No. 5105. On the second and third abdominal segments there are lateral pockets for the reception of hair-pencils.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA


1


1a

1b

2

2a

2b

3

3a

4

4a

4b

ANTAEOTRICHA

Antaeotricha bathrotoma (Meyrick), new combination

Plate 11, Figures 1–1b

Stenoma bathrotoma Meyrick, 1925, Exotic Microlepidoptera, 3: 189.

"♂. 16 mm. . . . Brazil, Obidos, September (*Parish*); 1 ex. Very close to *insidiata*. . ."

Type: The above indicated specimen dated ".9–19". Slide No. 5176.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha biarcuata Meyrick

Plate 11, Figures 2–2b

Antaeotricha biarcuata Meyrick, 1926, Exotic Microlepidoptera, 3: 238.

"♂. 20 mm. . . . Colombia, Minero, October; 1 ex."

Type: The above indicated specimen dated "10.20". Slide No. 4857.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha bilinguis (Meyrick), new combination

Plate 11, Figures 3–3a

Stenoma bilinguis Meyrick, 1918, Exotic Microlepidoptera, 2: 202.

"♂. 17 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen indicated above with "LeM. .15" at the bottom of the pin-label. Slide No. 5169.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha brachysaris Meyrick

Plate 11, Figures 4–4b

Antaeotricha brachysaris Meyrick, 1916, Exotic Microlepidoptera, 1: 504.

"♂. 18 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: One of the above males with "LeM. .15" at the bottom of the pin-label. Slide No. 4917.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha brochota Meyrick

Plate 12, Figures 1–1b

Antaeotricha brochota Meyrick, 1915, Exotic Microlepidoptera, 1: 396.

"♂. 22 mm. . . . Peru, Yquitos, in May (*Mounsey*); one specimen."

Type: The above indicated specimen dated "29.5.12". Slide No. 4851.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha caenochytis (Meyrick), new combination

Plate 12, Figures 2–2a

Stenoma caenochytis Meyrick, 1915, Exotic Microlepidoptera, 1: 415.

"♂ 16–17 mm., ♀ 18–20 mm. . . . British Guiana, Bartica and Mallali, from January to March (*Parish*); six specimens."

Lectotype: The male labelled "Mallali, Brit. Guiana, Parish, 3.13". Slide No. 5159.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha camarina Meyrick

Plate 12, Figures 3–3b

Antaeotricha camarina Meyrick, 1915, Exotic Microlepidoptera, 1: 401.

"♂. 17 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated "3.13". Slide No. 4919.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha campylodes Meyrick

Plate 12, Figures 4–4a

Antaeotricha campylodes Meyrick, 1916, Exotic Microlepidoptera, 1: 494.

"♂. 18 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The male denoted above with "LeM. .16". Slide No. 4894.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.



ANTAEOTRICHA


1

1a

2

2a

2b


3

3d


3c

3b

4


4a

ANTAEOTRICHA

Antaeotricha cantharitis (Meyrick)

Plate 13, Figures 1–1a

Aphanoxena cantharitis Meyrick, 1916, Exotic Microlepidoptera, 1: 490.

"♂. 17–18 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: One of the above indicated specimens all of which bear identical data with "LeM. .15" at the bottom of the pin-label. Slide No. 4854.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha capsiformis (Meyrick), new combination

Plate 13, Figures 4–4a

Stenoma capsiformis Meyrick, 1930, Exotic Microlepidoptera, 4: 25.

"♂. 14 mm. . . . Brazil, San Antonio da Barra, Bahia; 1 ex."

Type: The specimen referred to above and dated "11-12-88". A small white label bears the inscription "M855". The type lacks both forewings. Slide No. 5578.

Figure 4, lateral aspect of aedeagus; 4a, ventral view of male genitalia with aedeagus removed.

Antaeotricha capsulata Meyrick

Plate 13, Figures 2–2b

Antaeotricha capsulata Meyrick, 1918, Exotic Microlepidoptera, 2: 199.

"♂. 18 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, "R. Maroni, French Guiana. LeM. .15". Slide No. 4914.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha carabodes (Meyrick), new combination

Plate 13, Figures 3–3c

Stenoma carabodes Meyrick, 1915, Exotic Microlepidoptera, 1: 435.

"♀. 23 mm. . . . British Guiana, Bartica, in February (*Parish*); one specimen. Allied to *actista*."

Type: The above indicated specimen dated ".2.13". Slide No. 5501.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium; 3c, signum.

Antaeotricha carbasea (Meyrick), new combination

Plate 14, Figures 1–1c

Stenoma carbasea Meyrick, 1915, Exotic Microlepidoptera, 1: 426.

"♀. 20 mm. . . . Brazil, Novo Friburgo; one specimen."

Type: The female indicated above with "R. .06" at the bottom of the pin-label. Slide No. 5138.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Antaeotricha carphitis Meyrick

Plate 14, Figures 2–2b

Antaeotricha carphitis Meyrick, 1912, Trans. Ent. Soc. London, 1911: 710.

"♂. 24 mm. . . . Brazil, Santa Caterina; one specimen."

Type: The above indicated specimen with "D. .92" at the bottom of the pin-label. Slide No. 4846.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha cathagnista Meyrick

Plate 14, Figures 3–3b

Antaeotricha cathagnista Meyrick, 1925, Exotic Microlepidoptera, 3: 165.

"♂. 15–17 mm. . . . Brazil, R. Trombetas, Teffé, September, January (*Parish*); 2 ex. Very like *pactota*, but all veins present. . . ."

Lectotype: ♂, "R. Trombetas, Brazil, Parish. 9–19". Slide No. 4809.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha cedroxyla Meyrick

Plate 14, Figures 4–4b

Antaeotricha cedroxyla Meyrick, 1930, Exotic Microlepidoptera, 4: 23.

"♂. 18 mm. . . . Brazil, Queluz (*Germain*); 1 ex. (Coll. Paravicini). Allied to group of *gymnolopha*."

Type: The above indicated specimen. A small white label bears the inscription "M857". Slide No. 4948.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1
1a
1b
1c
2
2a
2b
3
3a
3b
4
4a
4b

1

2c

1a

2

2a

2b



3

3a

3b



4

4a

4b

ANTAEOTRICHA

Antaeotricha celidotis Meyrick

Plate 15, Figures 1–1a

Antaeotricha celidotis Meyrick, 1925, Exotic Microlepidoptera, 3: 169.

"♂. 18–19 mm. . . . Peru, R. Napo, May (*Parish*); 2 ex. Next *purulenta*."

Lectotype: ♂, 19 mm., dated "5.20". Slide No. 4873.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha chalastis (Meyrick)

Plate 15, Figures 2–2c

Stenoma chalastis Meyrick, 1915, Exotic Microlepidoptera, 1: 413.

"♂. 14 mm. . . . British Guiana, Bartica, in February (*Parish*); one specimen."

Type: The above indicated specimen dated ".2.13". Slide No. 5345. A hair-pencil from the anal area of hindwing is inserted into a tubular abdominal fold which extends to the fifth segment and continues as a blind sac to the seventh segment.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed; 2c, denuded abdomen to show abdominal folds and internal sacs.

Antaeotricha chalinophanes (Meyrick), new combination

Plate 15, Figures 3–3b

Stenoma chalinophanes Meyrick, 1931, Exotic Microlepidoptera, 4: 42.

"♂. 28 mm. . . . Bolivia, Cochabamba (*Germain*); 1 ex. (Coll. Paravicini).

Type: The above indicated male labelled "Bolivia, Cochabamba (Yunga del Esperitu Santo). P. Germain. 1888–89." A small white label bears the inscription "M922". Slide No. 5579.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha chelobathra Meyrick

Plate 15, Figures 4–4b

Antaeotricha chelobathra Meyrick, 1916, Exotic Microlepidoptera, 1: 498.

"♂. 25–26 mm. . . . French Guiana, St. Jean, R. Maroni; eight specimens."

Lectotype: One of the above specimens with "LeM. .15" at the bottom of the pin-label. Slide No. 4880.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha chilosema (Meyrick), new combination

Plate 16, Figures 1–1b

Stenoma chilosema Meyrick, 1918, Exotic Microlepidoptera, 2: 208.

"♂. 19–20 mm. . . . French Guiana, Godebert, R. Maroni; two specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .16." Slide No. 5453.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha chloromis (Meyrick), new combination

Plate 16, Figures 2–2c

Stenoma chloromis Meyrick, 1915, Exotic Microlepidoptera, 1: 433.

"♀. 25 mm. . . . British Guiana, Bartica, in December (*Perish*); one specimen."

Type: The specimen denoted above and dated "12.12". Slide No. 5100.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Antaeotricha christocoma Meyrick

Plate 16, Figures 3–3b

Antaeotricha christocoma Meyrick, 1915, Exotic Microlepidoptera, 1: 398.

"♂. 17–18 mm. . . . Peru, Pacaya and Contamano, R. Ucuyali, in June, August, September and December (*Mounsey*); five specimens."

Lectotype: ♂, "Pacaya, Peru. M. .8.12." Slide No. 4861.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha cirrhoxantha (Meyrick), new combination

Plate 16, Figures 4–4b

Stenoma cirrhoxantha Meyrick, 1915, Exotic Microlepidoptera, 1: 477.

"♂. 20–22 mm. . . . French Guiana, Godebert, R. Maroni; two specimens. Allied to *virens*; . . ."

Lectotype: One of the above specimens with "LeM. .15" at the bottom of the pin-label. Slide No. 5479.

Figure 4, right wings with image reversed; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA



1

1a

1b



2

2a



2b



3

3a

3b

4

4a

4b

ANTAEOTRICHA

Antaeotricha cleopatra Meyrick

Plate 17, Figures 1–1b

Antaeotricha cleopatra Meyrick, 1925, Exotic Microlepidoptera, 3: 166.

"♂. 15 mm. . . . Brazil, Teffé, December (*Parish*); 1 ex. near *pseudochyta*."

Type: The above indicated specimen dated "12.19". Slide No. 4812.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha clivosa Meyrick

Plate 17, Figures 2–2b

Antaeotricha clivosa Meyrick, 1918, Exotic Microlepidoptera, 2: 199.

"♀. 22 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .16" at the bottom of the pin-label. Slide No. 4869.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium.

Antaeotricha cnemosaris (Meyrick), new combination

Plate 17, Figures 3–3b

Stenoma cnemosaris Meyrick, 1925, Exotic Microlepidoptera, 3: 181.

"♂♀. 18–19 mm. . . . Brazil, Pará, July (*Parish*); 7 ex. Near *oxydecta*."

Lectotype: One of the above specimens all of which have identical data. Slide No. 5133.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha colposaris (Meyrick), new combination

Plate 17, Figures 4–4b

Stenoma colposaris Meyrick, 1925, Exotic Microlepidoptera, 3: 185.

"♂♀. 12–13 mm. . . . Brazil, Teffé, December (*Parish*); 2 ex."

Lectotype: ♂, dated "12.19". Slide No. 5152.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha compsographa Meyrick

Plate 18, Figures 1–1b

Antaeotricha compsographa Meyrick, 1916, Exotic Microlepidoptera, 1: 491.

"♂. 19–20 mm. . . . French Guiana, St. Jean, R. Maroni, in July; four specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 4804.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha compsoneura (Meyrick), new combination

Plate 18, Figures 2–2b

Stenoma compsoneura Meyrick, 1925, Exotic Microlepidoptera, 3: 217.

"♂. 16–17 mm. . . . Brazil, Pará, July (*Parish*); French Guiana, R. Maroni; 2 ex."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5442.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha congelata Meyrick

Plate 18, Figures 3–3b

Antaeotricha congelata Meyrick, 1926, Exotic Microlepidoptera, 3: 236.

"♂. 24 mm. . . . Peru, Cocapata, 12,000 feet; 1 ex."

Type: The above indicated specimen with "20" in the lower right of the pin-label. Slide No. 4834.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha coniopa (Meyrick), new combination

Plate 18, Figures 4–4a

Stenoma coniopa Meyrick, 1925, Exotic Microlepidoptera, 3: 184.

"♂. 11–12 mm. . . . Brazil, Obidos, August, September (*Parish*); 2 ex."

Lectotype: ♂, the male dated "9.19". Slide No. 5156.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.


1

1a

1b


2

2a

2b

3

3a

3b

4


4a

ANTAEOTRICHA


1

1a


1b


2

2a

2b

3

3a

3b

4

4a

4b

ANTAEOTRICHA

Antaeotricha constituta (Meyrick), new combination

Plate 19, Figures 1–1b

Stenoma constituta Meyrick, 1925, Exotic Microlepidoptera, 3: 191.

"♂. 18 mm. . . . French Guiana, R. Maroni; 1 ex."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5177.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha constricta (Meyrick), new combination

Plate 19, Figures 2–2b

Stenoma constricta Meyrick, 1926, Exotic Microlepidoptera, 3: 226.

"♂. 24 mm. . . . Colombia, Mt. Socorro, 12,500 feet, July, 1 ex. Near *fluminalis*."

Type: The above indicated specimen dated ".7.20". Slide No. 5432.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha copromima Meyrick

Plate 19, Figures 3–3b

Antaeotricha copromima Meyrick, 1930, Exotic Microlepidoptera, 4: 22.

"♀. 30 mm. . . . French Guiana, R. Maroni, 1 ex. (Coll. Paravicini). Perhaps nearest *cryeropis*; . . ."

Type: The above indicated female, without abdomen. A small white label bears the inscription "M870". The genitalia are figured from a male from "New Bremen, Brazil, 20 March 1936, Fritz Hoffman" in the U.S. National Museum. I have compared the male carefully with the type female and have no doubt about the identity. Slide No. 4949.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha coriodes Meyrick

Plate 19, Figures 4–4b

Antaeotricha coriodes Meyrick, 1915, Exotic Microlepidoptera, 1: 397.

"♂. 16–17 mm. . . . British Guiana, Mallali and Bartica, in February and March (*Parish*); two specimens. Allied to *confixella*."

Lectotype: ♂, "Mallali, Br. Guiana, Parish. .3.13." Slide No. 4852.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha corvigera Meyrick

Plate 20, Figures 1–1b

Antaeotricha corvigera Meyrick, 1915, Exotic Microlepidoptera, 1: 390.

"♂. 19 mm. . . . British Guiana, Mallali, in March (*Parish*); Peru, Contamano, R. Ucuyali, in December (*Mounsey*); two specimens."

Lectotype: ♂, "Mallali, Brit. Guiana, Parish. .3.13." Slide No. 4878.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha corvula Meyrick

Plate 20, Figures 2–2b

Antaeotricha corvula Meyrick, 1912, Trans. Ent. Soc. London, 1911: 710.

"♂. 21 mm. . . . Colombia, San Antonio, 5,800 feet, in November; one specimen."

Type: The above indicated specimen dated "11.07". Slide No. 4796.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia (damaged) with aedeagus removed.

Antaeotricha cremastis (Meyrick), new combination

Plate 20, Figures 3–3c

Stenoma cremastis Meyrick, 1925, Exotic Microlepidoptera, 3: 194.

"♀. 24 mm. . . . Peru, Jurimaguas, March (type); Brazil, Manaos, November (*Parish*); 2 ex."

Type: ♀, "Peru, Jurimaguas, Parish. 3.20", designated by Meyrick. Slide No. 4991.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Antaeotricha cryeropis Meyrick

Plate 20, Figures 4–4b

Antaeotricha cryeropis Meyrick, 1925, Exotic Microlepidoptera, 3: 167.

"♂♀. 23–25 mm. . . . Mexico, Guerrero, June to September; 4 ex."

Lectotype: One of the above specimens dated "9.20". Slide No. 4889.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA

1

1a

1b

2

2a

2c

3

2b

3a

4

4a

4b

ANTAEOTRICHA

Antaeotricha crypsiphaea (Meyrick), new combination

Plate 21, Figures 1–1b

Stenoma crypsiphaea Meyrick, 1925, Exotic Microlepidoptera, 3: 190.

"♂. 19 mm. . . . Brazil, Pará, July (*Parish*); 1 ex."

Type: The above indicated specimen dated ".7.19". Slide No. 5182.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha crypsithias (Meyrick), new combination

Plate 21, Figures 2–2c

Stenoma crypsithias Meyrick, 1930, Exotic Microlepidoptera, 3: 558.

"♀. 19 mm. . . . Brazil, Caraça; 1 ex. (Coll. Paravicini). Next *paracta*."

Type: The above indicated specimen labelled "Brazil, Caraça, P. Germain, 2e Semestre 1884" on a green label. A small white label bears the inscription "M82". Slide No. 5581.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Antaeotricha cycnolopha (Meyrick), new combination

Plate 21, Figures 3–3a

Stenoma cycnolopha Meyrick, 1925, Exotic Microlepidoptera, 3: 186.

"♂. 17 mm. . . . Peru, R. Napo, May (*Parish*); 1 ex."

Type: The above indicated male dated "5.20". Slide No. 5153.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha cycnomorpha Meyrick

Plate 21, Figures 4–4b

Antaeotricha cycnomorpha Meyrick, 1925, Exotic Microlepidoptera, 3: 169.

"♂. 20 mm. . . . Brazil, R. Trombetas, September (*Parish*); 1 ex. Next *deltopis*."

Type: The above indicated specimen dated ".9.19". Slide No. 4874.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha cymogramma (Meyrick), new combination

Plate 22, Figures 1–1b

Stenoma cymogramma Meyrick, 1925, Exotic Microlepidoptera, 3: 193.

"♂. 18–19 mm. . . . Peru, R. Napo, Iquitos, March to May (*Parish*); 2 ex."

Lectotype: The male labelled "Rio Napo, Peru, Parish. 5.20". Slide No. 5183.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha cynopis (Meyrick)

Plate 22, Figures 2–2b

Antaeotricha cynopis Meyrick, 1915, Exotic Microlepidoptera, 1: 434.

"♂. 28–29 mm. . . . British Guiana, Bartica, in January and February (*Parish*); five specimens."

Lectotype: A male dated ".2.13". Slide No. 5108.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha deltopis Meyrick

Plate 22, Figures 3–3b

Antaeotricha deltopis Meyrick, 1915, Exotic Microlepidoptera, 1: 390.

"♂. 17–20 mm. . . . British Guiana, Bartica, from December to February (*Parish*); five specimens. The absence of vein 4 of hindwings . . . is a notable character, . . ."

Lectotype: A male dated "1.13". Slide No. 4879.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha deridens Meyrick

Plate 22, Figures 4–4b

Antaeotricha deridens Meyrick, 1925, Exotic Microlepidoptera, 3: 162.

"♂. 20 mm. . . . Bolivia, Prov. del Sara, 1,500 feet; 1 ex. Near *cantharitis*."

Type: The specimen indicated above and labelled "Prov. del Sara, Dep. Sta. Cruz, Bolivia, 450 m.". Slide No. 4859.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1
1a
1b
2
2a
2b
3
3a
3b
4
4a
4b

1



1a

2



2a



3

3a



3b

4

4a



4b

ANTAEOTRICHA

Antaeotricha desecta (Meyrick), new combination

Plate 23, Figures 1–1a

Stenoma desecta Meyrick, 1918, Exotic Microlepidoptera, 2: 203.

"♂♀. 17–18 mm. . . . French Guiana, R. Maroni; five specimens."

Lectotype: The male with "LeM. .15" at the bottom of the pin-label. Slide No. 4983.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha diacta (Meyrick), new combination

Plate 23, Figures 2–2a

Stenoma diacta Meyrick, 1916, Exotic Microlepidoptera, 1: 513.

"♂. 17 mm. . . . French Guiana, R. Maroni; one specimen. Allied to *tephrodesma*."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5180. A hair-pencil from base of hindwing is inserted in a fold in first segment of abdomen; edges of fold studded with short, slender spines.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha diffracta Meyrick

Plate 23, Figures 3–3b

Antaeotricha diffracta Meyrick, 1916, Exotic Microlepidoptera, 1: 500.

"♂. 17–18 mm. . . . French Guiana, St. Jean, R. Maroni; three specimens."

Lectotype: The only remaining male in the Meyrick collection bearing the data "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15". Slide No. 4905.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha diplarcha Meyrick

Plate 23, Figures 4–4b

Antaeotricha diplarcha Meyrick, 1915, Exotic Microlepidoptera, 1: 403.

"♂. 23 mm. . . . British Guiana, Bartica, in December (*Parish*); one specimen."

Type: The above indicated specimen dated "12.12". Slide No. 4876.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha diplophaea Meyrick

Plate 24, Figures 1–1b

Antaeotricha diplophaea Meyrick, 1916, Exotic Microlepidoptera, 1: 494.

"♂ 17–18 mm., ♀ 22 mm. . . . French Guiana, Godebert and St. Jean, R. Maroni; three specimens."

Lectotype: ♂, "Godebert, R. Maroni, Fr. Guiana. LeM. .9.14." Slide No. 4891.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha diplosaris (Meyrick), new combination

Plate 24, Figures 2–2a

Stenoma diplosaris Meyrick, 1915, Exotic Microlepidoptera, 1: 418.

"♂♀. 15 mm. . . . British Guiana, Bartica, in January (*Parish*); three specimens."

Lectotype: ♂, dated "1.13". Slide No. 5127.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha disjecta (Zeller), new combination

Plate 24, Figures 3–3b

Cryptolechia disjecta Zeller, 1854, Linnaea Entomologica, 9: 368.
Stenoma intersecta Meyrick, 1916, Exotic Microlepidoptera, 1: 515; l.c., 3: 192; 1925.

[intersecta]

"♂. 24–25 mm. . . . French Guiana, R. Maroni; three specimens. Allied to *disjecta* and *lebetias*."

Lectotype: One of the above specimens all of which bear identical data with "LeM. .15" at the bottom of the pin-label.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha dissimilis (Kearfott)

Plate 24, Figures 4–4b

Stenoma dissimilis Kearfott, 1911, Ent. News, 22: 126.
Antaeotricha annixa Meyrick, 1918, Exotic Microlepidoptera, 2: 198.

[annixa]

"♂ 21 mm., ♀ 26–27 mm. . . . Brazil, São Paulo, Novo Friburgo; three specimens. Apparently allied to *tibialis*."

Lectotype: ♂, "São Paulo, Brazil. WDK. .11." Slide No. 7249.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA

1

1a

1b


2

2a


2b


2c


3

3a


3b


4

4a


4b

ANTAEOTRICHA

Antaeotricha dissona (Meyrick), new combination

Plate 25, Figures 1–1b

Stenoma dissona Meyrick, 1925, Exotic Microlepidoptera, 3: 191.

"♂. 15 mm. . . . Brazil, Manaos, November (*Parish*); 1 ex."

Type: The above indicated specimen dated "11.19". Slide No. 5181.

Figure 1, right wings with image reversed; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha doleropis (Meyrick), new combination

Plate 25, Figures 2–2c

Stenoma doleropis Meyrick, 1915, Exotic Microlepidoptera, 1: 421.

"♀. 23 mm. . . . British Guiana, Bartica, in April (*Parish*); one specimen."

Type: The above indicated specimen dated ".4.13". Slide No. 5502.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Antaeotricha dromica (Meyrick), new combination

Plate 25, Figures 3–3b

Stenoma dromica Meyrick, 1925, Exotic Microlepidoptera, 3: 185.

"♂ 14–15 mm., ♀ 16 mm. . . . Brazil, Parintins, October (*Parish*); 5 ex."

Lectotype: ♂, dated "10–19". All specimens bear identical data. Slide No. 5149.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha dynastis Meyrick

Plate 25, Figures 4–4b

Antaeotricha dynastis Meyrick, 1915, Exotic Microlepidoptera, 1: 398.

"♂. 25 mm. . . . Peru, Chanchamayo, in January (*Mounsey*); two specimens. Allied to *griseana*, . . ."

Lectotype: The male with the right forewing missing, dated ".1.13". Slide No. 4831. This appears to be only a colour form of *ampherista*.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha emollita Meyrick

Plate 26, Figures 1–1a

Antaeotricha emollita Meyrick, 1926, Exotic Microlepidoptera, 3: 234.

"♂. 25 mm. . . . Colombia; 1 ex. Next *melanopis*."

Type: The above indicated specimen with ".20" in the lower right of the pin-label. Slide No. 4800. The two hooked processes of the anellus are dorsal to the aedeagus.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha encyclia Meyrick

Plate 26, Figures 2–2b

Antaeotricha encyclia Meyrick, 1915, Exotic Microlepidoptera, 1: 403.

"♂. 28 mm. . . . Colombia, San Antonio, 5,800 feet, in December; one specimen."

Type: The above indicated specimen dated "12.07". Slide No. 4875.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha enodata Meyrick

Plate 26, Figures 3–3b

Antaeotricha enodata Meyrick, 1916, Exotic Microlepidoptera, 1: 493.

"♂. 24–27 mm. . . . French Guiana, Godebert and Nouveau Chantier, R. Maroni, in May; five specimens."

Lectotype: ♂, "Godebert, R. Maroni, Fr. Guiana, LeM. .5.15." Slide No. 4936.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha epignampta Meyrick

Plate 26, Figures 4–4b

Antaeotricha epignampta Meyrick, 1915, Exotic Microlepidoptera, 1: 395.

"♂. 19 mm. . . . Peru, Pacaya, in June (*Mounsey*); one specimen."

Type: The male denoted above, dated ".6.12". Slide No. 4890.

Figure 4, left wings; 4a, ventral view of aedeagus; 4b, lateral aspect of male genitalia (harpes broken off) with aedeagus removed.



ANTAEOTRICHA



ANTAEOTRICHA

Antaeotricha episimbla (Meyrick)

Plate 27, Figures 1–1c

Aphanoxena episimbla Meyrick, 1915, Exotic Microlepidoptera, 1: 389.

"♀. 15–17 mm. . . . British Guiana, Bartica and Mallali, from January to March (*Parish*); three specimens."

Lectotype: ♀, "Bartica, Brit. Guiana, Parish. 1.13." Slide No. 4921.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Antaeotricha erotica (Meyrick), new combination

Plate 27, Figures 2–2a

Stenoma erotica Meyrick, 1916, Exotic Microlepidoptera, 1: 534.

"♂. 18 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: One of the above indicated males with "LeM. .15" at the bottom of the pin-label. Slide No. 5392.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha eucoma Meyrick

Plate 27, Figures 3–3b

Antaeotricha eucoma Meyrick, 1925, Exotic Microlepidoptera, 3: 168.

"♂. 13 mm. . . . Brazil, Manaos, November (*Parish*); 1 ex. Allied to *christocoma* and *coriodes*."

Type: The above indicated specimen dated "11.19". Slide No. 4855.

Figure 3, right wings with image reversed; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha euthrinca Meyrick

Plate 27, Figures 4–4b

Antaeotricha euthrinca Meyrick, 1915, Exotic Microlepidoptera, 1: 399.

"♂. 28–29 mm. . . . Colombia, San Antonio, 5,800 feet, in November; two specimens."

Lectotype: ♂, 29 mm., with "R. 5800′ .11.07" at the bottom of the pin-label. Slide No. 4850.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha exasperata (Meyrick), new combination

Plate 28, Figures 1–1b

Stenoma exasperata Meyrick, 1916, Exotic Microlepidoptera, 1: 533.

"♂. 22–23 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .15." The second specimen is a female. Slide No. 5389.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha excisa Meyrick

Plate 28, Figures 2–2b

Antaeotricha excisa Meyrick, 1916, Exotic Microlepidoptera, 1: 496.

"♂. 18–20 mm. . . . French Guiana, St. Jean, R. Maroni, in July; three specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 4858.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha exusta Meyrick

Plate 28, Figures 3–3a

Antaeotricha exusta Meyrick, 1916, Exotic Microlepidoptera, 1: 492.

"♂. 27 mm. . . . French Guiana, R. Maroni; one specimen. Allied to *melanopis*."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 4797.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha falsidica (Meyrick), new combination

Plate 28, Figures 4–4b

Stenoma falsidica Meyrick, 1915, Exotic Microlepidoptera, 1: 426.

"♂. 19 mm. . . . Dutch Guiana, Berg-en-Daal, in April; one specimen."

Type: The above indicated specimen with "S. 4.92" at the bottom of the pin-label. Slide No. 5135.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a


1b


2

2a

2b

3

3a

4

4a

4b

ANTAEOTRICHA



ANTAEOTRICHA

Antaeotricha fascicularis Zeller

Plate 29, Figures 1–1c

Antaeotricha fascicularis Zeller, 1854, Linnaea Entomologica, 9: 393, pl. 3, figs. 32, 33.
Antaeotricha leucogramma Meyrick, 1915, Exotic Microlepidoptera, 1: 404.

[leucogramma]

"♀. 30 mm. . . . British Guiana, R. Demerara; one specimen."

Type: The female indicated above with "R. .06" at the bottom of the pin-label. Slide No. 4871.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of ostium; 1c, signum.

Antaeotricha fulta Meyrick

Plate 29, Figures 2–2b

Antaeotricha fulta Meyrick, 1926, Exotic Microlepidoptera, 3: 234.

"♂. 27 mm. . . . Colombia, Monte del Eden, 9,550 feet, October; 1 ex. Between *semiovata* and *Ribbei*."

Type: The specimen denoted above with ".20" in the lower right of the pin-label. Slide No. 4867.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha generatrix Meyrick

Plate 29, Figures 3–3b

Antaeotricha generatrix Meyrick, 1926, Exotic Microlepidoptera, 3: 239.

"♂. 17 mm. . . . Brazil, Santa Cruz, Rio Grande do Sul; 1 ex."

Type: The specimen denoted above. Slide No. 4828.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha glaciata Meyrick

Plate 29, Figures 4–4b

Antaeotricha glaciata Meyrick, 1909, Trans. Ent. Soc. London, 1909: 30.

"♂. 27–29 mm. . . . Bolivia, Songo; three specimens."

Lectotype: The male with "B. .07" at the bottom of the pin-label. Slide No. 4830.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha glaphyrodes (Meyrick), new combination

Plate 30, Figures 1–1c

Stenoma glaphyrodes Meyrick, 1913, Trans. Ent. Soc. London, 1913: 186.

"♀. 26–27 mm. . . . French Guiana, St. Laurient; Brazil, Iquitos [=? Peru]; two specimens."

Lectotype: ♀, labelled "Iquitos, Brazil. S. .93". Slide No. 5488.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Antaeotricha glaucescens (Meyrick), new combination

Plate 30, Figures 2–2b

Stenoma glaucescens Meyrick, 1916, Exotic Microlepidoptera, 1: 537.

"♂ 16–19 mm., ♀ 23 mm. . . . French Guiana, R. Maroni; four specimens."

Lectotype: A male with "LeM. .16" at the bottom of the pin-label. Slide No. 5478.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha glycerostoma Meyrick

Plate 30, Figures 3–3b

Antaeotricha glycerostoma Meyrick, 1915, Exotic Microlepidoptera, 1: 399.

"♂. 26 mm. . . . Colombia, San Antonio, 5,800 feet, in November; two specimens."

Lectotype: ♂, "San Antonio, Colombia. R. 5800′ .11.07." Slide No. 4849.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha gravescens Meyrick

Plate 30, Figures 4–4b

Antaeotricha gravescens Meyrick, 1926, Exotic Microlepidoptera, 3: 237.

"♂. 20 mm. . . . Colombia, Minero, October; 1 ex. Near *christocoma*."

Type: The above indicated specimen. Slide No. 4860.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1
1a
1b
1c
2
2a
2b
3
3a
3b
4
4a
4b

1

1a


2

2a


2b


3

3a

3b


4

4a

4b

ANTAEOTRICHA

Antaeotricha gubernatrix Meyrick

Plate 31, Figures 1–1a

Antaeotricha gubernatrix Meyrick, 1925, Exotic Microlepidoptera, 3: 173.

"♂. 18 mm. . . . Peru, R. Napo, May (*Parish*); 1 ex."

Type: The male denoted above, dated "5.20". Slide No. 4912.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha gymnolopha Meyrick

Plate 31, Figures 2–2b

Antaeotricha gymnolopha Meyrick, 1925, Exotic Microlepidoptera, 3: 174.

"♂. 15–16 mm. . . . Brazil, Parintins, Manaos, October, November (*Parish*); 2 ex."

Lectotype: ♂, "Parintins, Brazil, Parish. 10–19." Slide No. 4939.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha gypsoterma (Meyrick), new combination

Plate 31, Figures 3–3b

Stenoma gypsoterma Meyrick, 1915, Exotic Microlepidoptera, 1: 424.

"♂. 16 mm. . . . British Guiana, Mallali, in March (*Parish*); two specimens."

Lectotype: One of the above indicated males dated "3.13". Slide No. 5146. There is a paired, ventral hair-pencil from second segment.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha habilis (Meyrick), new combination

Plate 31, Figures 4–4b

Stenoma habilis Meyrick, 1915, Exotic Microlepidoptera, 1: 427.

"♂. 17–20 mm. . . . British Guiana, Bartica, from December to February (*Parish*); nine specimens."

Lectotype: ♂, dated "2.13". Slide No. 5137.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha haplocentra Meyrick

Plate 32, Figures 1–1b

Antaeotricha haplocentra Meyrick, 1925, Exotic Microlepidoptera, 3: 170.

"♂♀. 16–17 mm. . . . Brazil, Obidos, Parintins, August to October (*Parish*); 3 ex."

Lectotype: ♂, "Obidos, Brazil, Parish. .8.19." Slide No. 4909. The vesica contains much extraneous matter which could not be removed.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha hapsicora Meyrick

Plate 32, Figures 2–2c

Antaeotricha hapsicora Meyrick, 1915, Exotic Microlepidoptera, 1: 399.

"♀. 26–28 mm. . . . Brazil, São Paulo and Petropolis; two specimens."

Lectotype: ♀, "Petropolis, Brazil". Slide No. 4848.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of ostium and genital plate; 2c, signum.

Antaeotricha harpobathra Meyrick

Plate 32, Figures 3–3b

Antaeotricha harpobathra Meyrick, 1916, Exotic Microlepidoptera, 1: 499.

"♂. 27 mm. . . . Argentina, Parana, in August; one specimen."

Type: The above indicated specimen with "LeM. .8.15" at the bottom of the pin-label. Slide No. 4886.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha helicias Meyrick

Plate 32, Figures 4–4a

Antaeotricha helicias Meyrick, 1916, Exotic Microlepidoptera, 1: 502.

"♂. 25 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen, without date, with "LeM" at the bottom of the pin-label. Slide No. 4904. Sacculus armed with a curved, flat process.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

1

1a


1b

2

2a


2b

3

3a

3b

4

2c

4a

ANTAEOTRICHA



ANTAEOTRICHA

Antaeotricha herilis Felder & Rogenhofer

Plate 33, Figures 1–1b

Antaeotricha herilis Felder & Rogenhofer, 1875, Reise der Osterreichischen Fregatte Novara, 2: pl. 138, fig. 66.

Antaeotricha xanthoptila Meyrick, 1912, Trans. Ent. Soc. London, 1911: 709 (new synonymy).

[xanthoptila]

"♂. 30 mm. . . . Dutch Guiana, interior, in August; one specimen. Near *basalis* Zell."

Type: The specimen referred to above with "E. 8.92" at the bottom of the pin-label. Slide No. 4882.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha heteropa (Meyrick), new combination

See Antaeotricha nictitans (Zeller), p. 91.

Antaeotricha heterosaris (Meyrick), new combination

Plate 33, Figures 2–2b

Stenoma heterosaris Meyrick, 1915, Exotic Microlepidoptera, 1: 418.

"♂. 15 mm. . . . British Guiana, Bartica, in February (*Parish*); Dutch Guiana, Onoribo, in March; two specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. 2.13." Slide No. 5126.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha himaea Meyrick

Plate 33, Figures 3–3a, 3c

Antaeotricha himaea Meyrick, 1916, Exotic Microlepidoptera, 1: 505.

"♂♀. 17 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .15". Slide No. 4942.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3c, ventral view of male genitalia with aedeagus removed.

Antaeotricha homologa (Meyrick)

Plate 33, Figures 4–4c

Aphanoxena homologa Meyrick, 1915, Exotic Microlepidoptera, 1: 388.

"♀. 19 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen."

Type: The above indicated specimen dated "1.13". Slide No. 4810.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum; 4c, detail of genital plate and ostium.

Antaeotricha horizontias (Meyrick), new combination

Plate 34, Figures 1–1b

Stenoma horizontias Meyrick, 1925, Exotic Microlepidoptera, 3: 188.

"♂. 13 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex."

Type: The above indicated specimen dated "1.20". Slide No. 5167.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha hydrophora Meyrick

Plate 34, Figures 2–2a

Antaeotricha hydrophora Meyrick, 1925, Exotic Microlepidoptera, 3: 171.

"♂. 13 mm. . . . Peru, Iquitos, March (*Parish*); 1 ex."

Type: The male denoted above dated "3.20". Slide No. 4935.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha illepida Meyrick

Plate 34, Figures 3–3b

Antaeotricha illepida Meyrick, 1916, Exotic Microlepidoptera, 1: 495.

"♂. 20–21 mm. . . . French Guiana, Godebert, R. Maroni; three specimens."

Lectotype: One of the above specimens with "LeM. .10.15" at the bottom of the pin-label. Slide No. 4896.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha imminens (Meyrick), new combination

Plate 34, Figures 4–4c

Stenoma imminens Meyrick, 1915, Exotic Microlepidoptera, 1: 431.

"♀. 28 mm. . . . Dutch Guiana, Onoribo, in January; one specimen."

Type: The specimen indicated above with "E. .1.93" at the bottom of the pin-label. Slide No. 5505.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, detail of genital plate and ostium; 4c, signum.



ANTAEOTRICHA



ANTAEOTRICHA

Antaeotricha immota Meyrick

Plate 35, Figures 1–1a

Antaeotricha immota Meyrick, 1916, Exotic Microlepidoptera, 1: 502.

"♂♀. 14–15 mm. . . . French Guiana, R. Maroni; British Guiana, Mallali, in March (*Parish*); five specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .15." The Mallali specimens (2 ♀♀) were collected by Parish. Slide No. 4929.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha impedita (Meyrick), new combination

Plate 35, Figures 2–2c

Stenoma impedita Meyrick, 1915, Exotic Microlepidoptera, 1: 422.

"♀. 19 mm. . . . Peru, Pacaya, in June (*Mounsey*); one specimen."

Type: The above indicated specimen dated ".6.12". Slide No. 5184. The genitalia are badly damaged by mould.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signum.

Antaeotricha incrassata Meyrick

Plate 35, Figures 3–3a

Antaeotricha incrassata Meyrick, 1916, Exotic Microlepidoptera, 1: 504.

"♂. 21 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: One of the above indicated males. Slide No. 4916.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha infrenata (Meyrick), new combination

Plate 35, Figures 4–4b

Stenoma infrenata Meyrick, 1918, Exotic Microlepidoptera, 2: 202.

"♂ 17 mm., ♀ 20 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 5189.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha innexa (Meyrick), new combination

Plate 36, Figures 1–1b

Stenoma innexa Meyrick, 1925, Exotic Microlepidoptera, 3: 190.

"♂♀. 16–19 mm. . . . Peru, Jurimaguas, Iquitos, March (*Parish*); 4 ex."

Lectotype: ♂, "Iquitos, Peru, Parish. 3.20." Slide No. 5188. One specimen is missing.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha insidiata (Meyrick), new combination

Plate 36, Figures 2–2b

Stenoma insidiata Meyrick, 1916, Exotic Microlepidoptera, 1: 512.

"♂. 19–22 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: One of the males denoted above with "LeM. .15" at the bottom of the pin-label. Slide No. 5186.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha insimulata Meyrick

Plate 36, Figures 3–3a

Antaeotricha insimulata Meyrick, 1926, Exotic Microlepidoptera, 3: 237.

"♂. 26 mm. . . . Colombia, San Antonio, 6,600 feet, February; 1 ex. Next *helicias*."

Type: The specimen indicated above dated "2.20". Slide No. 4911.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha intersecta (Meyrick), new combination

See Antaeotricha disjecta (Zeller), p. 48.

Antaeotricha iopetra (Meyrick)

Plate 36, Figures 4–4b

Stenoma iopetra Meyrick, 1932, Exotic Microlepidoptera, 4: 295.

"♂. 24 mm. . . . Guatemala, Cayuga, August (*A. Busck*); 1 ex."

Type: The above indicated specimen dated ".8.20". Slide No. 5118.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA


1

1a


1b

1c

2

2a


2b


3

3a


3b

4


4a


4b

ANTAEOTRICHA

Antaeotricha ioptila (Meyrick), new combination

Plate 37, Figures 1–1c

Stenoma ioptila Meyrick, 1915, Exotic Microlepidoptera, 1: 433.

"♀. 23–24 mm. . . . British Guiana, Bartica, in December and January (*Parish*); three specimens."

Lectotype: The female dated "12.12". Slide No. 4997.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Antaeotricha iras Meyrick

Plate 37, Figures 2–2b

Antaeotricha iras Meyrick, 1926, Exotic Microlepidoptera, 3: 237.

"♂. 18 mm. . . . Peru, Cocapata, 12,000 feet; 1 ex. Nearest *cleopatra*."

Type: The above indicated specimen with ".20" in the lower right corner of the pin-label. Slide No. 4822.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha irenias (Meyrick), new combination

Plate 37, Figures 3–3b

Stenoma irenias Meyrick, 1916, Exotic Microlepidoptera, 1: 537.

"♂. 16 mm. . . . French Guiana, St. Jean, R. Maroni, in July; one specimen."

Type: The specimen denoted above with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5483.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha isochyta (Meyrick), new combination

Plate 37, Figures 4–4b

Stenoma isochyta Meyrick, 1915, Exotic Microlepidoptera, 1: 420.

"♂. 17 mm. . . . British Guiana, Bartica, in December (*Parish*); one specimen."

Type: The specimen indicated above, dated "12.12". Slide No. 5132.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha isomeris (Meyrick), new combination

Plate 38, Figures 1–1b

Stenoma isomeris Meyrick, 1912, Trans. Ent. Soc. London, 1911: 711.

"♂. 19 mm. . . . Brazil, Tijuco, in December; one specimen."

Type: The above indicated male labelled "Tijuco, Brazil. R. .12.06." Slide No. 5136.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha isoplintha (Meyrick), new combination

Plate 38, Figures 2–2b

Stenoma isoplintha Meyrick, 1925, Exotic Microlepidoptera, 3: 193.

"♂♀. 22–23 mm. . . . Brazil, Parintins, October (*Parish*); 2 ex."

Lectotype: The male dated ".10–19". Slide No. 4989.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha ithytona Meyrick

Plate 38, Figures 3–3b

Antaeotricha ithytona Meyrick, 1929, Trans. Ent. Soc. London, 76: 514.

"♂♀. 15–16 mm. . . . Colombia, Gorgona I., 200 feet, July, October, at light; 7 ex. Allied to *ovatella* and *tractrix*."

Type: ♂, so marked in the British Museum, "Gorgona Island, Colombia. At light, 200 ft. 7.7.24. St. George Expedn. C. L. Collenette." Slide No. 4945.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha lacertosa Meyrick

See Antaeotricha pallulella (Busck), p. 100.

Antaeotricha lathiptila (Meyrick)

Plate 38, Figures 4–4b

Stenoma lathiptila Meyrick, 1915, Exotic Microlepidoptera, 1: 425.

"♂♀. 17–18 mm. . . . British Guiana, Bartica, in December and January (*Parish*); five specimens."

Lectotype: ♂, dated "1.13". Slide No. 5187.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1
1a
1b
2
2a
2b
3
3a
3b
4
4a
4b

1



1a

1b

2

2a



2b

3

3c

3a

3b



4

4a

4b

ANTAEOTRICHA

Antaeotricha laudata Meyrick

Plate 39, Figures 1–1b

Antaeotricha laudata Meyrick, 1916, Exotic Microlepidoptera, 1: 496.

"♂. 21–23 mm. . . . French Guiana, St. Jean and Godebert, R. Maroni; fourteen specimens. Much like *basalis*. . . ."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 4842.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha laxa (Meyrick), new combination

Plate 39, Figures 2–2b

Stenoma laxa Meyrick, 1915, Exotic Microlepidoptera, 1: 428.

"♂. 17 mm. . . . Venezuela, Ciudad Bolivar, in July; one specimen."

Type: The above indicated specimen with "R. .7.04" at the bottom of the pin-label. Slide No. 4987.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha lebetias (Meyrick)

Plate 39, Figures 3–3c

Stenoma lebetias Meyrick, 1915, Exotic Microlepidoptera, 1: 433.

"♀. 23 mm. . . . French Guiana, S. Laurent; one specimen."

Type: The above indicated specimen, without abdomen, labelled "St. Laurient, Cayenne. R. .11". Male genitalia figured from a specimen from "Teffé, Brazil, Parish. 1.20" which I have compared carefully with the type. Slide No. 4999.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed; 3c, sketch of lateral aspect of gnathos.

Antaeotricha lecithaula Meyrick

Plate 39, Figures 4–4b

Antaeotricha lecithaula Meyrick, 1915, Exotic Microlepidoptera, 1: 401.

"♂. 13 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen."

Type: The above specimen dated "1.13". Slide No. 4923.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha leucocryptis (Meyrick), new combination

Plate 40, Figures 1–1b

Stenoma leucocryptis Meyrick, 1932, Exotic Microlepidoptera, 4: 295.

"♂. 28 mm. . . . Colombia, Medina, 1,650 feet; 1 ex. Probably allied to *subita*."

Type: The specimen denoted above with ".20" in the lower right of the pin-label. Slide No. 5500.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha leucogramma Meyrick

See Antaeotricha fascicularis Zeller.

Antaeotricha lophoptycha (Meyrick), new combination

Plate 40, Figures 2–2b

Stenoma lophoptycha Meyrick, 1925, Exotic Microlepidoptera, 3: 188.

"♂. 17 mm. . . . Brazil, Parintins, October (*Parish*); 1 ex."

Type: The above indicated specimen dated "10–19". Slide No. 5160.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha lophosaris (Meyrick), new combination

Plate 40, Figures 3–3b

Stenoma lophosaris Meyrick, 1925, Exotic Microlepidoptera, 3: 186.

"♂. 14 mm. . . . Brazil, R. Trombetas, September (*Parish*); 1 ex."

Type: The specimen denoted above dated "9.19". Slide No. 5150.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha lucrosa (Meyrick), new combination

Plate 40, Figures 4–4b

Stenoma lucrosa Meyrick, 1925, Exotic Microlepidoptera, 3: 184.

"♂ 14–15 mm., ♀ 16 mm. . . . Brazil, Obidos, Parintins, August, October (*Parish*); 4 ex."

Lectotype; ♂, "Obidos, Brazil, Parish. .8–19." Slide No. 5161. Second segment of abdomen with ventro-lateral pocket of hair-like scales.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA



ANTAEOTRICHA

Antaeotricha lysimeris Meyrick

Plate 41, Figures 1–1b

Antaeotricha lysimeris Meyrick, 1915, Exotic Microlepidoptera, 1: 391.

"♂. 26 mm. . . . Peru, Pacaya, in July (*Mounsey*); one specimen."

Type: The above indicated specimen dated ".7.12". Slide No. 4824.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha macronota (Meyrick), new combination

Plate 41, Figures 2–2b

Stenoma macronota Meyrick, 1912, Trans. Ent. Soc. London, 1911: 716.

"♂♀. 24–27 mm. . . . Colombia, Naranjito, R. Dagua, at 3,900 feet, in June, 1 ♂; Dutch Guiana, Paramaribo, in December, 1 ♀."

Lectotype: The male dated ".6.08". Slide No. 5111. The female probably represents a different species.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha malachita Meyrick

Plate 41, Figures 3–3b

Antaeotricha malachita Meyrick 1915, Exotic Microlepidoptera, 1: 404.

"♂. 22 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated male dated "3.13". Slide No. 4799.

Figure 3, left wings; 3a, ventral aspect of aedeagus to show spinous process; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha manceps Meyrick

Plate 41, Figures 4–4b

Antaeotricha manceps Meyrick, 1925, Exotic Microlepidoptera, 3: 172.

"♂. 13 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex."

Type: The male indicated above dated "3.20". Slide No. 4928.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha melanarma Meyrick

Plate 42, Figures 1–1b

Antaeotricha melanarma Meyrick, 1916, Exotic Microlepidoptera, 1: 500.

"♂. 20–22 mm. . . . French Guiana, St. Jean, R. Maroni; two specimens. . . . The black markings of the ♂ are a striking distinctive character."

Lectotype: ♂, with "LeM. .16" at the bottom of the pin-label. Neither specimen is marked "St. Jean" but both are from "R. Maroni". Slide No. 4906.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha melanopis Meyrick

Plate 42, Figures 2–2a

Antaeotricha melanopis Meyrick, 1909, Trans. Ent. Soc. London, 1909: 31.

"♂. 36 mm. . . . Peru, Huancabamba; one specimen."

Type: The male indicated above with "R. .08" at the bottom of the pin-label. Slide No. 4801.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha melinopa (Meyrick), new combination

Plate 42, Figures 3–3b

Stenoma melinopa Meyrick, 1925, Exotic Microlepidoptera, 3: 222.

"♂. 18–20 mm. . . . Brazil, Obidos, September (*Parish*); 2 ex."

Lectotype: The specimen measuring 18 mm., dated ".9–19". Slide No. 5480.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha mentigera Meyrick

Plate 42, Figures 4–4b

Antaeotricha mentigera Meyrick, 1926, Exotic Microlepidoptera, 3: 236.

"♂. 24–25 mm. . . . Bolivia, Prov. del Sara, Dept. Santa Cruz, 1,500 feet, November; 2 ex."

Lectotype: ♂, "Bolivia, Prov. del Sara, Dept. Santa Cruz, 1,500 feet, November. J. Steinbach." Slide No. 4877.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA

1

1a

1b

2



2a



3

3a

3b



4

4a

4b

ANTAEOTRICHA

Antaeotricha meridiana (Meyrick), new combination

Plate 43, Figures 1–1b

Stenoma meridiana Meyrick, 1915, Exotic Microlepidoptera, 1: 435.

"♂. 24 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated ".3.13". Slide No. 5454.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha mesosaris (Meyrick), new combination

Plate 43, Figures 2–2a

Stenoma mesosaris Meyrick, 1925, Exotic Microlepidoptera, 3: 178.

"♂. 10 mm. . . . French Guiana, R. Maroni; 1 ex."

Type: The above specimen with "LeM. .16" at the bottom of the pin-label.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha mesostrota Meyrick

Plate 43, Figures 3–3b

Antaeotricha mesostrota Meyrick, 1912, Trans. Ent. Soc. London, 1911: 708; 1915, Exotic Microlepidoptera, 1: 393.

"♂. 17 mm. . . . Venezuela, Carupano, in December; one specimen."

Type: The male denoted above, without abdomen. Male genitalia figured from a specimen "Mallali, Brit. Guiana, Parish. .3.13" referred to by Meyrick in his second reference. Slide No. 4817.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha microtypa (Meyrick), new combination

Plate 43, Figures 4–4b

Stenoma microtypa Meyrick, 1915, Exotic Microlepidoptera, 1: 422.

"♂♀. 14–15 mm. . . . British Guiana, Bartica, in January (*Parish*); two specimens."

Lectotype: ♂, dated "1.13". The female is missing. Slide No. 5154.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha milictis Meyrick

Plate 44, Figures 1–1b

Antaeotricha milictis Meyrick, 1925, Exotic Microlepidoptera, 3: 163.

"♂. 21 mm. . . . Brazil, Teffé, December (*Parish*); 2 ex."

Lectotype: The male with "HP. .12.19" at the bottom of the pin-label. Slide No. 4832.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha modulata (Meyrick), new combination

Plate 44, Figures 2–2b

Stenoma modulata Meyrick, 1915, Exotic Microlepidoptera, 1: 436.

"♂. 17 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen."

Type: The above indicated specimen dated "1.13". Slide No. 5397.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha monosaris (Meyrick), new combination

Plate 44, Figures 3–3a

Stenoma monosaris Meyrick, 1915, Exotic Microlepidoptera, 1: 419.

"♂♀. 15 mm. . . . British Guiana, Bartica and Mallali, from January to March (*Parish*); seven specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. 2.13." Slide No. 5130.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha neocrossa (Meyrick), new combination

Plate 44, Figures 4–4b

Stenoma neocrossa Meyrick, 1925, Exotic Microlepidoptera, 3: 193.

"♂. 13 mm. . . . Peru, R. Napo, May (*Parish*); 1 ex."

Type: The above indicated specimen dated "5.20". Slide No. 4988. The hair-pencils from base of sacculus are unique.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA



ANTAEOTRICHA

Antaeotricha nerteropa Meyrick

Plate 45, Figures 1–1b

Antaeotricha nerteropa Meyrick, 1915, Exotic Microlepidoptera, 1: 395.

"♂. 18 mm. . . . Peru, Pacaya, in July (*Mounsey*); one specimen."

Type: The male indicated above dated ".7.12". Slide No. 4821.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha nictitans (Zeller)

Plate 45, Figures 2–2c

Mesoptycha nictitans Zeller, 1854, Linnaea Entomologica, 9: 389, pl. 3, figs. 26, 28.
Stenoma heteropa Meyrick, 1913, Trans. Ent. Soc. London, 1913: 187.

[heteropa]

"♂♀. 27–30 mm. . . . British Guiana, R. Demerara; Dutch Guiana, Paramaribo; French Guiana, St. Laurient; Paraguay; seven specimens."

Lectotype: ♀, "R. Demerara, Brit. Guiana, .06." Slide No. 5629. The lectotype is the only one of the original specimens remaining in the Meyrick collection. I have not seen Zeller's type so must accept Meyrick's synonymy.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signum.

Antaeotricha nimbata Meyrick

Plate 45, Figures 3–3b

Antaeotricha nimbata Meyrick, 1925, Exotic Microlepidoptera, 3: 175.

"♂. 14 mm. . . . Peru, Iquitos, March (*Parish*); 1 ex."

Type: The above indicated specimen dated "3.20". Slide No. 4940.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha nitescens Meyrick

Plate 45, Figures 4–4a

Antaeotricha nitescens Meyrick, 1925, Exotic Microlepidoptera, 3: 171.

"♂. 17 mm. . . . Brazil, Pará, July (*Parish*); 1 ex."

Type: The above indicated specimen dated "7–19". Slide No. 4925.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha nitrota Meyrick

Plate 46, Figures 1–1b

Antaeotricha nitrota Meyrick, 1916, Exotic Microlepidoptera, 1: 497.

"♂. 22 mm. . . . French Guiana, Godebert, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 4863.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha notosaris (Meyrick), new combination

Plate 46, Figures 2–2a

Stenoma notosaris Meyrick, 1925, Exotic Microlepidoptera, 3: 181.

"♂♀. 14–15 mm. . . . Brazil, Pará, July (*Parish*); 3 ex."

Lectotype: ♂, the more strongly marked of the two remaining specimens. Slide No. 5122. The long, slender, curved process from sacculus is unique.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha nuclearis Meyrick

Plate 46, Figures 3–3a

Antaeotricha nuclearis Meyrick, 1913, Trans. Ent. Soc. London, 1913: 181.

"♂. 25–26 mm. . . . Peru, Chanchamayo; two specimens."

Lectotype: ♂, 25 mm., "Chanchamayo, Peru. R. .11." Slide No. 4823.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha obtusa (Meyrick), new combination

Plate 46, Figures 4–4b

Stenoma obtusa Meyrick, 1916, Exotic Microlepidoptera, 1: 513.

"♂. 19 mm. . . . French Guiana, St. Jean, R. Maroni, in July; one specimen."

Type: The specimen denoted above with "LeM. .7.15" at the bottom of the pin-label.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a


1b

2


2a


3


3a


4

4a

4b

1

1a

1b


2

2a

2b


3

3a


3b


4


4a

4b

ANTAEOTRICHA

Antaeotricha ogmolopha (Meyrick), new combination

Plate 47, Figures 1–1b

Stenoma ogmolopha Meyrick, 1930, Exotic Microlepidoptera, 3: 557.

"♂. 13 mm. . . . Brazil, Santarem, August (*Parish*); 1 ex. Allied to *colposaris*."

Type: The above indicated specimen dated "8.19". Slide No. 5151.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha ogmosaris (Meyrick), new combination

Plate 47, Figures 2–2b

Stenoma ogmosaris Meyrick, 1916, Exotic Microlepidoptera, 1: 415.

"♂. 15 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated ".3.13". Slide No. 5164.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha ophrysta Meyrick

Plate 47, Figures 3–3b

Antaeotricha ophrysta Meyrick, 1912, Trans. Ent. Soc. London, 1911: 708.

"♂. 25 mm. . . . Dutch Guiana, Onoribo, in January and March; two specimens."

Lectotype: The specimen dated "1.93". Slide No. 4899.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed (right cucullus broken off).

Antaeotricha orgadopa (Meyrick) new combination

Plate 47, Figures 4–4b

Stenoma orgadopa Meyrick, 1925, Exotic Microlepidoptera, 3: 182.

"♂. 13 mm. . . . Brazil, Pará, Parintins, August, October (*Parish*); 2 ex."

Lectotype: ♂, "Parintins, Brazil, Parish. 10–19." Slide No. 5140. There are ventro-lateral hair-pencils from between the first and second and the seventh and eighth segments as in *triplecta*.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha orthotona Meyrick

Plate 48, Figures 1–1c

Antaeotricha orthotona Meyrick, 1916, Exotic Microlepidoptera, 1: 495.

"♀. 19–21 mm. . . . British Guiana, Bartica, in February (*Parish*); French Guiana, R. Maroni; three specimens."

Lectotype: A female, "R. Maroni, Fr. Guiana. LeM. .16." Slide No. 4813.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Antaeotricha orthriopa Meyrick

Plate 48, Figures 2–2a

Antaeotricha orthriopa Meyrick, 1925, Exotic Microlepidoptera, 3: 166.

"♂. 16 mm. . . . Brazil, Parintins, October (*Parish*); 1 ex. Allied to *inquinula*."

Type: The above indicated specimen dated "10.19". Slide No. 4807.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha ovulifera (Meyrick), new combination

Plate 48, Figures 3–3b

Stenoma ovulifera Meyrick, 1925, Exotic Microlepidoptera, 3: 182.

"♂. 12 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex."

Type: The male denoted above dated "3.20". Slide No. 5141.

Figure 3, left wings; 3a, sketch of lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha oxycentra Meyrick

Plate 48, Figures 4–4b

Antaeotricha oxycentra Meyrick, 1916, Exotic Microlepidoptera, 1: 497.

"♂. 16 mm. . . . French Guiana, St. Jean, R. Maroni, in July; two specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 4853.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA


1


1a

1b


2

2c

2a

2b

3

3a

3b


4

4a


4b

ANTAEOTRICHA

Antaeotricha oxydecta (Meyrick)

Plate 49, Figures 1–1b

Stenoma oxydecta Meyrick, 1915, Exotic Microlepidoptera, 1: 426.

"♂. 19–20 mm. . . . British Guiana, Bartica and Mallali, from January to March (*Parish*); three specimens."

Lectotype: ♂, 19 mm., "Mallali, Brit. Guiana, Parish. .3.13." Slide No. 8571.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha oxyschista (Meyrick), new combination

Plate 49, Figures 2–2c

Stenoma oxyschista Meyrick, 1925, Exotic Microlepidoptera, 3: 187.

"♂. 11–13 mm. . . . Brazil, Teffé, December, January (*Parish*); 3 ex."

Lectotype: ♂, dated "1.20". Slide No. 5155.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed; 2c, denuded abdomen to show abdominal fold and hair-pencil.

Antaeotricha pactota Meyrick

Plate 49, Figures 3–3b

Antaeotricha pactota Meyrick, 1915, Exotic Microlepidoptera, 1: 391; 1925, 3: 164.

"♂. 17–18 mm. . . . British Guiana, Bartica, in December and January (*Parish*); four specimens."

Lectotype: A male dated "12.12". Slide No. 4808.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha palaestrias Meyrick

Plate 49, Figures 4–4b

Antaeotricha palaestrias Meyrick, 1916, Exotic Microlepidoptera, 1: 502.

"♂. 13–15 mm. . . . French Guiana, St. Jean, R. Maroni; eight specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 4924.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha pallulella (Busck)

Plate 50, Figures 1–1b

Stenoma pallulella Busck, 1914, Proc. U.S. Nat. Mus., 47: 47.
Antaeotricha lacertosa Meyrick, 1915, Exotic Microlepidoptera, 1: 404.

[lacertosa]

"♂. 21–22 mm. . . . British Guiana, Bartica, in December and January (*Parish*); French Guiana, S. Laurient; eight specimens. Allied to *Zelleri*."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. 12.12." Slide No. 4803.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha paracrypta Meyrick

Plate 50, Figures 2–2b

Antaeotricha paracrypta Meyrick, 1915, Exotic Microlepidoptera, 1: 405.

"♂♀. 15 mm. . . . British Guiana, Bartica, in January and February (*Parish*); three specimens."

Lectotype: The male dated "2.13". Slide No. 4944.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha paracta (Meyrick), new combination

Plate 50, Figures 3–3b

Stenoma paracta Meyrick, 1915, Exotic Microlepidoptera, 1: 425.

"♂. 26 mm. . . . Peru, Chanchamayo and Porvenir; Colombia, San Antonio, 5,800 feet, in November; three specimens."

Lectotype: ♂, "Chanchamayo, Peru. R. .11." Slide No. 4986.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha pellocoma (Meyrick)

See Aphanoxena Meyrick, p. 140.

Antaeotricha percnocarpa (Meyrick), new combination

Plate 50, Figures 4–4b

Stenoma percnocarpa Meyrick, 1925, Exotic Microlepidoptera, 3: 181.

"♂. 17 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex."

Type: The above indicated specimen dated "1.20". Slide No. 5178.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA


1

1a

1b

2

2a

2b

3

3a


3b

4

4a

4b

4c

ANTAEOTRICHA

Antaeotricha percnogona Meyrick

Plate 51, Figures 1–1b

Antaeotricha percnogona Meyrick, 1925, Exotic Microlepidoptera, 3: 170.

"♂. 13 mm. . . . Peru, Iquitos, March (*Parish*); 1 ex."

Type: The above indicated specimen dated "3.20". Slide No. 4936.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha perfusa Meyrick

Plate 51, Figures 2–2b

Antaeotricha perfusa Meyrick, 1916, Exotic Microlepidoptera, 1: 504.

"♂. 18–19 mm. . . . French Guiana, St. Jean, R. Maroni; four specimens."

Lectotype: A specimen with "LeM. .7–15" at the bottom of the pin-label. Slide No. 4915.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha periphrictis (Meyrick), new combination

Plate 51, Figures 3–3b

Stenoma periphrictis Meyrick, 1915, Exotic Microlepidoptera, 1: 451.

"♀. 18 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen. Probably allied rather to the *murinella* group."

Type: The female indicated above dated ".1.13". Slide No. 5343.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

Antaeotricha phaeoneura (Meyrick), new combination

Plate 51, Figures 4–4c

Stenoma phaeoneura Meyrick, 1913, Trans. Ent. Soc. London, 1913: 187.

"♀. 29 mm. . . . British Guiana; one specimen."

Type: The female denoted above labelled "British Guiana, R. .06". Slide No. 5101.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum; 4c, detail of genital plate and ostium.

Antaeotricha phaeoplintha (Meyrick), new combination

Plate 52, Figures 1–1a

Stenoma phaeoplintha Meyrick, 1915, Exotic Microlepidoptera, 1: 424.

"♂♀. 13–14 mm. . . . British Guiana, Bartica, from December to February (*Parish*); twenty-eight specimens."

Lectotype: A male dated "12.12". Slide No. 5162. Of the original series only 14 remain in the Meyrick collection.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha phaeosaris Meyrick

Plate 52, Figures 2–2b

Antaeotricha phaeosaris Meyrick, 1915, Exotic Microlepidoptera, 1: 394; 1916, 1: 492.

"♂. 21 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated "3.13". Slide No. 4820.

Figure 2, left wings; 2a, left harpe; 2b, lateral aspect of male genitalia with left harpe removed and aedeagus *in situ*.

Antaeotricha phollicodes (Meyrick), new combination

Plate 52, Figures 3–3a

Stenoma phollicodes Meyrick, 1916, Exotic Microlepidoptera, 1: 533.

"♂ 19 mm., ♀ 22 mm. . . . French Guiana, St. Jean, R. Maroni; two specimens."

Lectotype: The male labelled "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 5388.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha phryactis Meyrick

Plate 52, Figures 4–4b

Antaeotricha phryactis Meyrick, 1925, Exotic Microlepidoptera, 3: 167.

"♂. 25 mm. . . . Peru, R. Napo, May (*Parish*); 1 ex. Very distinct, perhaps related to *staurota* group."

Type: The specimen indicated above dated "5.20". Slide No. 4888.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a


2


2a

2b


3

3a

4

4a

4b

ANTAEOTRICHA

1

1a

1b


2

2c

2b

3

2d


4

4d

4b

ANTAEOTRICHA

Antaeotricha planicoma (Meyrick), new combination

Plate 53, Figures 1–1b

Stenoma planicoma Meyrick, 1925, Exotic Microlepidoptera, 3: 189.

"♂. 18 mm. . . . Brazil, Santarem, August (*Parish*); 1 ex."

Type: The above indicated specimen dated "8–19". Slide No. 5173.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha platydesma Meyrick

Plate 53, Figures 2–2c

Antaeotricha platydesma Meyrick, 1915, Exotic Microlepidoptera, 1: 405.

"♂♀. 20–23 mm. . . . British Guiana, Bartica, from December to February (*Parish*); seven specimens."

Lectotype: A male dated ".1.13". Slide No. 4903.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed; 2c, sketch of anellus, lateral aspect.

Antaeotricha plerotis Meyrick

Plate 53, Figure 3

Antaeotricha plerotis Meyrick, 1925, Exotic Microlepidoptera, 3: 165.

"♂. 17 mm. . . . Peru, Pacaya, August (*Mounsey*); 1 ex. Very similar to *cathagnista*. . . ."

Type: The above indicated specimen dated ".8.12". The type is without abdomen.

Figure 3, left wings.

Antaeotricha plesistia (Meyrick), new combination

Plate 53, Figures 4–4b

Stenoma plesistia Meyrick, 1930, Exotic Microlepidoptera, 4: 24.

"♂♀. 21–23 mm. . . . Brazil, Petropolis (*Germain*); 2 ex. (Coll. Paravicini). Perhaps allied to *carbasea*."

Type: The male in the British Museum labelled "Estado Rio de Janeiro, Petropolis, P. Germain, 1888". A small white label bears the inscription "M844". Slide No. 5591.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha polyglypta (Meyrick), new combination

Plate 54, Figures 1–1c

Stenoma polyglypta Meyrick, 1915, Exotic Microlepidoptera, 1: 427.

"♀. 18–20 mm. . . . British Guiana, Mallali, in March (*Parish*); four specimens."

Lectotype: One of the above females dated ".3.13". Slide No. 5102.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Antaeotricha praecisa Meyrick

Plate 54, Figures 2–2b

Antaeotricha praecisa Meyrick, 1912, Trans. Ent. Soc. London, 1911: 709.

"♂♀. 20–25 mm. . . . Brazil, Rio de Janeiro, São Paulo; three specimens. Near *bicolor*, Zell."

Lectotype: ♂, "Rio de Janeiro, Brazil, R. 06." Slide No. 4829.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha praerupta Meyrick

Plate 54, Figures 3–3a

Antaeotricha praerupta Meyrick, 1915, Exotic Microlepidoptera, 1: 394.

"♂. 20 mm. . . . British Guiana, Bartica, in February (*Parish*); one specimen."

Type: The above indicated specimen dated "2.13". Slide No. 4819.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha pratifera (Meyrick), new combination

Plate 54, Figures 4–4b

Stenoma pratifera Meyrick, 1925, Exotic Microlepidoptera, 3: 222.

"♂. 20 mm. . . . Costa Rica; 1 ex. Near *viridis*."

Type: The above indicated specimen with "HS. .20" at the bottom of the pin-label. Slide No. 5481.

Figure 4, left wings; 4a, sketch of anellus and aedeagus, lateral aspect; 4b, ventral view of male genitalia with aedeagus *in situ*.

1

1a

1b


2


2a

2b


3

1c

3a


4

4a

4b



1

1a

1b

2

2a

2b



3

3a

4

4a

4b

ANTAEOTRICHA

Antaeotricha protosaris Meyrick

Plate 55, Figures 1–1b

Antaeotricha protosaris Meyrick, 1915, Exotic Microlepidoptera, 1: 406.

"♂♀. 13–15 mm. . . . British Guiana, Bartica, in January (*Parish*); three specimens."

Lectotype: The male dated "1.13". Slide No. 4943.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha pseudochyta Meyrick

Plate 55, Figures 2–2b

Antaeotricha pseudochyta Meyrick, 1915, Exotic Microlepidoptera, 1: 393.

"♂ 13–15 mm., ♀ 16–19 mm. . . . British Guiana, Bartica, from December to February (*Parish*); twenty-two specimens. The characteristic stalking of veins 8 and 9 is invariable and strongly pronounced, but cannot be esteemed a ground for generic separation. . . ."

Lectotype: A male dated ".2.13". Slide No. 4816.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha pyrobathra (Meyrick), new combination

Plate 55, Figures 3–3a

Stenoma pyrobathra Meyrick, 1931, Exotic Microlepidoptera, 4: 33.

"♂. 16 mm. . . . Brazil, Bahia, S. Antonio da Barra, December; 1 ex. (Coll. Paravicini)."

Type: The specimen referred to above. In addition to the data recorded above there is "Gounelle 1888" on the pin-label. Slide No. 5686.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha pythonaea Meyrick

Plate 55, Figures 4–4b

Antaeotricha pythonaea Meyrick, 1916, Exotic Microlepidoptera, 1: 491.

"♂. 22 mm. . . . French Guiana, R. Maroni; one specimen. Intermediate between *malachita* and *Zelleri*; . . ."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 4872. This is not a synonym of *A. fumipennis* (Busck).

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha quiescens (Meyrick), new combination

Plate 56, Figures 1–1c

Stenoma quiescens Meyrick, 1916, Exotic Microlepidoptera, 1: 514.

"♀. 23–24 mm. . . . French Guiana, Godebert, R. Maroni; four specimens."

Lectotype: ♀, "Godebert, R. Maroni, Fr. Guiana, LeM. .15." Slide No. 4996.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Antaeotricha reprehensa Meyrick

Plate 56, Figures 2–2b

Antaeotricha reprehensa Meyrick, 1926, Exotic Microlepidoptera, 3: 238.

"♂. 18 mm. . . . Brazil, Santa Cruz, Rio Grande do Sul; 1 ex. Nearest *oxycentra* which has grey hindwings."

Type: The above indicated specimen without date or collector shown. Slide No. 4856.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha resiliens Meyrick

Plate 56, Figures 3–3b

Antaeotricha resiliens Meyrick, 1925, Exotic Microlepidoptera, 3: 173.

"♂. 13 mm. . . . Brazil, Parintins, October (*Parish*); 1 ex."

Type: The male denoted above dated ".10–19". Slide No. 4937.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha rhipidaula (Meyrick), new combination

Plate 56, Figures 4–4b

Stenoma rhipidaula Meyrick, 1915, Exotic Microlepidoptera, 1: 431.

"♀. 27 mm. . . . British Guiana, Bartica, in February (*Parish*); one specimen."

Type: The specimen referred to above dated ".2.13". Slide No. 5504.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum.


1

1a

1b

1c

2


2a

2b


3

3a

3b

4

4a

4b

ANTAEOTRICHA



ANTAEOTRICHA

Antaeotricha rostriformis (Meyrick), new combination

Plate 57, Figures 1–1b

Stenoma rostriformis Meyrick, 1916, Exotic Microlepidoptera, 1: 532.

"♂. 18–19 mm. . . . French Guiana, St. Jean, R. Maroni, in July; three specimens."

Lectotype: One of the three specimens indicated above, all of which bear identical data with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5387.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha sana Meyrick

Plate 57, Figures 2–2b; 4a

Antaeotricha sana Meyrick, 1926, Exotic Microlepidoptera, 3: 235.

"♀. 25 mm. . . . Colombia, Sosomoko, 2,650 feet, March; 1 ex."

Type: The above indicated specimen with ".20" in the lower right of the pin-label. Slide No. 4868. This may be the female of *fulta*.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 4a, signum.

Antaeotricha sarcinata Meyrick

Plate 57, Figures 3–3b

Antaeotricha sarcinata Meyrick, 1918, Exotic Microlepidoptera, 2: 200.

"♂. 17 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .16" at the bottom of the pin-label. Slide No. 4933.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha sardania Meyrick

Plate 57, Figures 4, 4b

Antaeotricha sardania Meyrick, 1925, Exotic Microlepidoptera, 3: 168.

"♂. 14–15 mm. . . . Brazil, Pará, Teffé; Peru, Jurimaguas (*Parish*); 3 ex. A distinct form perhaps related to *campylodes*. . . ."

Lectotype: ♂, "Pará, Brazil, Parish. .6–19." Slide No. 4893.

Figure 4, left wings; 4b, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha scapularis (Meyrick), new combination

Plate 58, Figures 1–1c

Stenoma scapularis Meyrick, 1918, Exotic Microlepidoptera, 2: 209.

"♀. 32 mm. . . . French Guiana, R. Maroni; one specimen. Nearly allied to *glaphyrodes*. . . ."

Type: The specimen denoted above with "LeM. .15" at the bottom of the pin-label. Slide No. 5487.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Antaeotricha segmentata (Meyrick), new combination

Plate 58, Figures 2–2b

Stenoma segmentata Meyrick, 1915, Exotic Microlepidoptera, 1: 423.

"♂. 18 mm. . . . British Guiana, Bartica, in March (*Parish*); one specimen."

Type: The specimen indicated above dated "3.13". Slide No. 4998.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha sellifera Meyrick

Plate 58, Figures 3–3b

Antaeotricha sellifera Meyrick, 1925, Exotic Microlepidoptera, 3: 163.

"♂. 15 mm. . . . Brazil, Parintins, Manaos, October, November (*Parish*); 2 ex. Allied to *praecisa*."

Lectotype: The specimen from "Manaos, Brazil, Parish. 11.19." Slide No. 4827.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha semiovata Meyrick

Plate 58, Figures 4–4b

Antaeotricha semiovata Meyrick, 1926, Exotic Microlepidoptera, 3: 234.

"♂. 29 mm. . . . Colombia, Alto de las Anecs, 7,250 feet; 1 ex. Nearest *Thammi*."

Type: The above indicated specimen with ".20" in the lower right of the pin-label. Slide No. 4866.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA



ANTAEOTRICHA

Antaeotricha serangodes Meyrick

Plate 59, Figures 1–1b

Antaeotricha serangodes Meyrick, 1915, Exotic Microlepidoptera, 1 : 400.

"♂. 26 mm. . . . Panama, Chiriqui; one specimen."

Type: The above indicated specimen with "R. .99" at the bottom of the pin-label. Slide No. 4847.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia (right harpe broken off) with aedeagus removed.

Antaeotricha serarcha Meyrick

Plate 59, Figures 2–2c

Antaeotricha serarcha Meyrick, 1930, Exotic Microlepidoptera, 3 : 556.

"♀. 24 mm. . . . Brazil, Caraça; 1 ex. (Coll. Paravicini). Belongs to the *basalis* group. . . ."

Type: The above indicated specimen "Brésil, Caraça. P. Germain. 2e Semestre. 1884." Slide No. 4946.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Antaeotricha smileuta Meyrick

Plate 59, Figures 3–3b

Antaeotricha smileuta Meyrick, 1915, Exotic Microlepidoptera, 1 : 397.

"♂. 21–22 mm. . . . British Guiana, Bartica, in December (*Parish*); French Guiana. S. Laurient; two specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. 12.12." Slide No. 4901.

Figure 3, left wings; 3a, lateral aspect of aedeagus ; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha sortifera Meyrick

Plate 59, Figures 4–4c

Antaeotricha sortifera Meyrick, 1930, Exotic Microlepidoptera, 3 : 557.

"♀. 22 mm. . . . Bolivia, Cochabamba; 1 ex. (Coll. Paravicini). Extremely near *adjunctella* Walk. . . ."

Type: The female denoted above, "Bolivie, Cochabamba (Yunga del Esperitu Santo), P. Germain. 1888–89." Slide No. 4947.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum; 4c, detail of genital plate and ostium.

Antaeotricha sparganota Meyrick

Plate 60, Figures 1–1b

Antaeotricha sparganota Meyrick, 1915, Exotic Microlepidoptera, 1: 389.

"♂. 20–21 mm. . . . British Guiana, Bartica, in January and February (*Parish*); two specimens."

Lectotype: The specimen dated "1.13". Slide No. 4887.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha spermolitha (Meyrick)

Plate 60, Figures 2–2c

Stenoma spermolitha Meyrick, 1915, Exotic Microlepidoptera, 1: 432.

"♀. 23 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen.

Type: The above indicated specimen dated ".1.13". Slide No. 5503.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signum.

Antaeotricha staurota Meyrick

Plate 60, Figures 3–3c

Antaeotricha staurota Meyrick, 1916, Exotic Microlepidoptera, 1: 493.

"♀. 23 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen denoted above with "LeM. .15" at the bottom of the pin-label. Slide No. 4892.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium; 3c, signum.

Antaeotricha sterrhomitra (Meyrick), new combination

Plate 60, Figure 4

Stenoma sterrhomitra Meyrick, 1925, Exotic Microlepidoptera, 3: 178.

"♂♀. 12–13 mm. . . . Brazil, Teffé, December, January (*Parish*); 3 ex."

Lectotype: ♂, dated "1.20". The abdomens of the three specimens are missing.

Figure 4, left wings.



ANTAEOTRICHA


1

1a


1b


2

2a


2b


3

3a


3b


4

4a

4b

4c

ANTAEOTRICHA

Antaeotricha stringens Meyrick

Plate 61, Figures 1–1b

Antaeotricha stringens Meyrick, 1925, Exotic Microlepidoptera, 3: 164.

"♂. 21 mm. . . . Brazil, Teffé, December (*Parish*); 2 ex. Extremely similar to *smileuta*, . . ."

Lectotype: The specimen dated "12.19". Slide No. 4907.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha stygeropa (Meyrick), new combination

Plate 61, Figures 2–2b

Stenoma stygeropa Meyrick, 1925, Exotic Microlepidoptera, 3: 187.

"♂. 14–15 mm. . . . Brazil, Manaos, November (*Parish*); 2 ex."

Lectotype: One of the above males both of which have identical data and are dated "11.19". Slide No. 5148.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha subdulcis (Meyrick), new combination

Plate 61, Figures 3–3b

Stenoma subdulcis Meyrick, 1925, Exotic Microlepidoptera, 3: 195.

"♂. 16 mm. . . . Brazil, Santarem, August (*Parish*); 1 ex."

Type: The above indicated specimen dated "8.19". Slide No. 4992.

Figure 3, left wings; 3a, lateral aspect of aedeagus (much extraneous matter is present in the vesica and must not be confused with the elongate apical patch of cornuti); 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha submersa (Meyrick), new combination

Plate 61, Figures 4–4c

Stenoma submersa Meyrick, 1915, Exotic Microlepidoptera, 1: 429.

"♀. 23 mm. . . . British Guiana, Bartica, in February (*Parish*); one specimen."

Type: The above indicated specimen dated "2.13". Slide No. 4985.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum; 4c, detail of genital plate (deformed?) and ostium.

Antaeotricha substricta Meyrick

Plate 62, Figures 1–1b

Antaeotricha substricta Meyrick, 1918, Exotic Microlepidoptera, 2: 200.

"♂. 15 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen indicated above with "LeM. .15" at the bottom of the pin-label. Slide No. 4934.

Figure 1, right wings with image reversed; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha superciliosa Meyrick

Plate 62, Figures 2–2b

Antaeotricha superciliosa Meyrick, 1918, Exotic Microlepidoptera, 2: 198.

"♂. 23 mm. . . . French Guiana, R. Maroni; two specimens. Differs from all others known to me by the peculiar patch of rough scales in the disc of forewings."

Lectotype: The specimen with "LeM. .16" at the bottom of the pin-label. Slide No. 4900.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha synercta Meyrick

Plate 62, Figures 3–3b

Antaeotricha synercta Meyrick, 1925, Exotic Microlepidoptera, 3: 173.

"♂. 17 mm. . . . Brazil, Parintins, October (*Parish*); 1 ex."

Type: The specimen indicated above dated "10–19". Slide No. 4913.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha tanysta (Meyrick)

Plate 62, Figures 4–4b

See p. 127.


1


1a

1b

2

2a


2b


3

3a


3b

4

4a

4b



1

1a

1b



2

2a



2b



3

3a

3b



4

2c

4a

ANTAEOTRICHA

Antaeotricha tanysta (Meyrick)

Plate 62, Figures 4–4b; Plate 63, Figures 1–1b

Antaeotricha tanysta Meyrick, 1915, Exotic Microlepidoptera, 1: 402.
Stenoma anticharis Meyrick, 1916, Exotic Microlepidoptera, 1: 512.

[tanysta]

Plate 63, Figures 1–1b

"♂. 18–19 mm. . . . British Guiana, Bartica and Mallali, in February and March (*Parish*); three specimens."

Lectotype: ♂, "Mallali, Brit. Guiana, Parish. .3.13." Slide No. 5174.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

[anticharis]

Plate 62, Figures 4–4b

"♂. 18–19 mm. . . . French Guiana, St. Jean, R. Maroni; three specimens."

Lectotype: One of the above indicated specimens all of which bear the same data with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5190.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha tectoria (Meyrick), new combination

Plate 63, Figures 2–2c

Stenoma tectoria Meyrick, 1915, Exotic Microlepidoptera, 1: 429.

"♀. 32–33 mm. . . . British Guiana, Bartica, in January and February (*Parish*); two specimens."

Lectotype: The specimen dated "2–13". Slide No. 4995.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signum.

Antaeotricha teleosema Meyrick

Plate 63, Figures 3–3b

Antaeotricha teleosema Meyrick, 1925, Exotic Microlepidoptera, 3: 165.

"♂. 16–17 mm. . . . Brazil, Parintins, Teffé, October, January (*Parish*); 4 ex."

Lectotype: ♂, "Parintins, Brazil, Parish. 10–19." Slide No. 4811.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha tempestiva (Meyrick), new combination

Plate 63, Figures 4–4a

Stenoma tempestiva Meyrick, 1916, Exotic Microlepidoptera, 1: 529.

"♂. 22 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5367.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha tephrodesma (Meyrick), new combination

Plate 64, Figures 1–1b

Stenoma tephrodesma Meyrick, 1916, Exotic Microlepidoptera, 1: 511.

"♂. 14–15 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: ♂, with "LeM. .15" at the bottom of the pin-label. Slide No. 5168. Two of the original three males have neither hindwings nor abdomens. I have selected the third male as lectotype. Abdomen with latero-longitudinal furrow containing hair-pencil.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha tetrapetra (Meyrick), new combination

Plate 64, Figures 2–2c

Stenoma tetrapetra Meyrick, 1925, Exotic Microlepidoptera, 3: 194.

"♀. 25 mm. . . . Brazil, Teffé, December (*Parish*); 1 ex."

Type: The type, indicated above, is dated "12.19". Slide No. 4990.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Antaeotricha thapsinopa Meyrick

Plate 64, Figures 3–3b

Antaeotricha thapsinopa Meyrick, 1916, Exotic Microlepidoptera, 1: 498.

"♂. 18–19 mm. . . . French Guiana, St. Jean, R. Maroni, in July; two specimens."

Lectotype: The specimen labelled "St. Jean, R. Maroni, Fr. Guiana, LeM. .7.15." Slide No. 4870.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha thesmophora Meyrick

Plate 64, Figures 4–4b

Antaeotricha thesmophora Meyrick, 1915, Exotic Microlepidoptera, 1: 392.

"♀. 18–20 mm. . . . British Guiana, Bartica, in January (*Parish*); two specimens. At first sight much like *inquinula*, . . ."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. .1.13." Both specimens, erroneously recorded by Meyrick as females, are in the Meyrick collection. Slide No. 4818.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

1

1a


1b

2

2a

2b

2c


3

3a


3b


4

4a

4b

ANTAEOTRICHA

1

1a


1b


2

2a

2b

3


3a

3b


4

4a

4b

ANTAEOTRICHA

Antaeotricha thylacosaris (Meyrick), new combination

Plate 65, Figures 1–1b

Stenoma thylacosaris Meyrick, 1915, Exotic Microlepidoptera, 1: 419.

"♂. 17–21 mm. . . . British Guiana, Bartica, from December to February (*Parish*); four specimens."

Lectotype: The specimen dated "2.13". Slide No. 5131.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha thysanodes (Meyrick), new combination

Plate 65, Figures 2–2b

Stenoma thysanodes Meyrick, 1915, Exotic Microlepidoptera, 1: 434.

"♂. 31 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated "3.13". Slide No. 5191.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha tinactis (Meyrick), new combination

Plate 65, Figures 3–3b

Stenoma tinactis Meyrick, 1915, Exotic Microlepidoptera, 1: 414.

"♂. 13 mm. . . . British Guiana, Bartica, in February (*Parish*); one specimen."

Type: The above indicated specimen dated ".2.13". Slide No. 5165.

Figure 3, left wings; 3a, lateral aspect of aedeagus (there are two long cornuti, one of which appears as a triangular spot in photograph due to foreshortening); 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha tornogramma Meyrick

Plate 65, Figures 4–4b

Antaeotricha tornogramma Meyrick, 1925, Exotic Microlepidoptera, 3: 171.

"♂. 15 mm. . . . Brazil, Parintins, October (*Parish*); 1 ex."

Type: The above indicated specimen dated "10–19". Slide No. 4926.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha tractrix Meyrick

Plate 66, Figures 1–1b

Antaeotricha tractrix Meyrick, 1925, Exotic Microlepidoptera, 3: 172.

"♂. 14 mm. . . . Brazil, Obidos, September (*Parish*); 1 ex."

Type: The above indicated specimen dated ".9–19". Slide No. 4932.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha tribomias (Meyrick), new combination

Plate 66, Figures 2–2c

Stenoma tribomias Meyrick, 1915, Exotic Microlepidoptera, 1: 417.

"♀. 19 mm. . . . British Guiana, Bartica, in December (*Parish*); one specimen."

Type: The above indicated specimen dated "12.12". Slide No. 5170.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signum.

Antaeotricha trichonota Meyrick

Plate 66, Figures 3–3b

Antaeotricha trichonota Meyrick, 1926, Exotic Microlepidoptera, 3: 235.

"♂. 21–24 mm. . . . Brazil, Santa Cruz, Rio Grande do Sul; Paraguay; 2 ex. A peculiar form but allied to *dissimilis*."

Lectotype: ♂, 24 mm., "Paraguay, .20". Slide No. 4865.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha triplectra (Meyrick), new combination

Plate 66, Figures 4–4b

Stenoma triplectra Meyrick, 1915, Exotic Microlepidoptera, 1: 423.

"♂♀. 14–17 mm. . . . British Guiana, Bartica, in January and February (*Parish*); three specimens. Allied to *murinella*, . . ."

Lectotype: The male dated "1.13". Slide No. 5144. In the male there are ventrolateral tufts from the second segment and from between the seventh and eighth abdominal segments.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



ANTAEOTRICHA



ANTAEOTRICHA

Antaeotricha triplintha (Meyrick)

Plate 67, Figures 1–1a

Aphanoxena triplintha Meyrick, 1916, Exotic Microlepidoptera, 1: 490.

"♂. 15 mm. . . . French Guiana, St. Jean, R. Maroni, in July; one specimen."

Type: The specimen denoted above with "LeM. .7.15" at the bottom of the pin-label. Slide No. 4930. In this species the harpe is fused to the vinculum and tegumen, and the aedeagus is fused solidly with the base of anellus.

Figure 1, left wings; 1a, lateral view of male genitalia with aedeagus *in situ.*

Antaeotricha trisinuata Meyrick

Plate 67, Figure 2

Antaeotricha trisinuata Meyrick, 1930, Exotic Microlepidoptera, 4: 21.

"♂. 18 mm. . . . Brazil, 1 ex. (Coll. Paravicini). A peculiar form."

Type: The unique specimen in the British Museum, from the Paravicini collection. is a female and was erroneously recorded as a male by Meyrick. On a small white label appears the inscription "M898". The type is without abdomen.

Figure 2, left wings.

Antaeotricha tritogramma Meyrick

Plate 67, Figures 3–3b

Antaeotricha tritogramma Meyrick, 1925, Exotic Microlepidoptera, 3: 176.

"♂♀. 13–15 mm. . . . Brazil, Parintins, Teffé, October, January (*Parish*); 6 ex."

Lectotype: ♂, "Parintins, Brazil, Parish. .10–19." Five of the original specimens, all from Parintins and with identical data remain in the Meyrick collection. Slide No. 4938.

Figure 3, left wings; 3a, aedeagus, lateral aspect; 3b, ventral view of male genitalia with aedeagus removed.

Antaeotricha trochoscia Meyrick

Plate 67, Figures 4–4b

Antaeotricha trochoscia Meyrick, 1915, Exotic Microlepidoptera, 1: 396.

"♂. 20 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated "3.13". Slide No. 4895.

Figure 4, left wings; 4a, aedeagus, lateral aspect; 4b, ventral view of male genitalia with aedeagus removed.

Antaeotricha tumens (Meyrick), new combination

Plate 68, Figures 1–1b

Stenoma tumens Meyrick, 1916, Exotic Microlepidoptera, 1: 511.

"♂. 17 mm. . . . French Guiana, St. Jean, R. Maroni, in July; one specimen."

Type: The male denoted above with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5179.

Figure 1, left wings; 1a, aedeagus, lateral aspect; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha vacata Meyrick

Plate 68, Figures 2–2b

Antaeotricha vacata Meyrick, 1925, Exotic Microlepidoptera, 3: 170.

"♂. 20 mm. . . . Grenada, St. George; 1 ex. Probably allied to *platydesma*."

Type: The above indicated specimen with "E. .12.91" at the bottom of the pin-label. Slide No. 4908. The wings on the right side of the type are missing; those on the left appear badly discoloured.

Figure 2, left wings; 2a, aedeagus, lateral aspect; 2b, ventral view of male genitalia with aedeagus removed.

Antaeotricha vannifera (Meyrick), new combination

Plate 68, Figures 3–3a; 4, 4b

Stenoma vannifera Meyrick, 1915, Exotic Microlepidoptera, 1: 416.
Stenoma asphalopis Meyrick, 1925, Exotic Microlepidoptera, 3: 188 (new synonymy).

[vannifera]

"♂. 15 mm. . . . Peru, Pacaya, in July (*Mounsey*); one specimen."

Type: The above indicated specimen with "M. .7.12" at the bottom of the pin-label. Slide No. 5163.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

[asphalopis]

"♂. 14 mm. . . . French Guiana, St. Jean, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5166.

Figure 4, left wings; 4b, ventral view of male genitalia with aedeagus *in situ*.

Antaeotricha xanthoptila Meyrick

See Antaeotricha herilis (Felder and Rogenhofer).

1

1a

1b

2


2a

2b


3

3a

4

4b

ANTAEOTRICHA

1

1a



1b



2

2a



3



4c

2b



4

4a

4b

ANTAEOTRICHA

Antaeotricha xuthosaris Meyrick

Plate 69, Figures 1–1b

Antaeotricha xuthosaris Meyrick, 1925, Exotic Microlepidoptera, 3: 175.

"♂. 19 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex."

Type: The above indicated specimen dated "1.20". Slide No. 4910.

Figure 1, left wings; 1a, aedeagus, lateral view; 1b, ventral view of male genitalia with aedeagus removed.

Antaeotricha xylocosma Meyrick

Plate 69, Figures 2–2b

Antaeotricha xylocosma Meyrick, 1916, Exotic Microlepidoptera, 1: 491.

"♂. 27 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated male with "LeM. .15" at the bottom of the pin-label. Slide No. 4798. Two slender processes of the anellus lie dorsad to the aedeagus. Aedeagus with serrate distal edge.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*; 2b, enlarged view to show serrate distal edge of aedeagus.

Antaeotricha xylurga (Meyrick), new combination

Plate 69, Figure 3

Stenoma xylurga Meyrick, 1913, Trans. Ent. Soc. London, 1913: 188.

"♂. 17 mm. . . . Peru, Chanchamayo; one specimen."

Type: The above indicated specimen labelled "Chanchamayo, Peru, R. 11". The type is without abdomen.

Figure 3, left wings.

Antaeotricha zanclogramma (Meyrick), new combination

Plate 69, Figures 4–4c

Stenoma zanclogramma Meyrick, 1915, Exotic Microlepidoptera, 1: 417.

"♀. 17 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen. . . ."

Type: The above indicated specimen dated ".1.13". Slide No. 5171.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, detail of genital plate and ostium; 4c, signum.

APHANOXENA Meyrick

Aphanoxena Meyrick, 1915, Exotic Microlepidoptera, 1: 386.

(*Typus generis: Aphanoxena pellocoma* Meyrick, l.c., 387 [by original designation].)

This genus is a synonym of *Antaeotricha*.

Antaeotricha pellocoma (Meyrick)

Plate 70, Figures 1–1d

Aphanoxena pellocoma Meyrick, 1915, Exotic Microlepidoptera, 1: 387.

"♂. 16 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen, without abdomen, dated ".3.13".

Male genitalia figured from a specimen from Obidos, Brazil, Parish. "9–19". Slide No. 4805. Although there is a chance that the Obidos specimen may not be the same as the type, it matches the type perfectly in colour and markings.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, aedeagus, lateral aspect; 1d, ventral view of male genitalia with aedeagus removed.

Continued from p. 11.

Although Walker's type is a female and Meyrick's type of *pleximorpha* is a male I do not hesitate to unite them. In 1932 Meyrick described the species again under the name of *dictyogramma*, from "Brazil, Teffé (*Plason*); 2 ex." Both specimens, one of which is in the Meyrick collection, are females. I have made a genitalic preparation from the Meyrick specimen of *dictyogramma* (Slide No. 5333) and one from Walker's type of *venosella* (Slide No. 5334). The genitalia agree in all details.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Anadasmus germinans (Meyrick), new combination

Plate 5, Figures 4–4b

Stenoma germinans Meyrick, 1925, Exotic Microlepidoptera, 3: 211.

"♂. 27 mm. . . . Colombia, locality uncertain; 1 ex. The tufts of forewings are unique, but the normal palpi and other characters show that it is not a *Ptilogenes*. Allied to *pleurotricha*."

Type: The above indicated specimen with ".20" in the lower right of the pin-label. The "tufts" appear on the left forewing only and were presumably caused by faulty spreading technique. Slide No. 5287.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.









APHANOXENA



1

1a



1b

1c

1d

BAEONOMA

BAEONOMA Meyrick

Baeonoma Meyrick, 1916, Exotic Microlepidoptera, 1: 507.

(*Typus generis: Baeonoma mastodes* Meyrick, l.c.: 508 [by original designation].)

Baeonoma mastodes Meyrick

Plate 71, Figures 1–1d

Baeonoma mastodes Meyrick, 1916, Exotic Microlepidoptera, 1: 508.

"♂♀. 15–19 mm. . . . French Guiana, R. Maroni; seven specimens. The variation in colour of thorax is curious; it is not sexual, but the two specimens with wholly white thorax are both ♀."

Lectotype: A male labelled "R. Maroni, Fr. Guiana, LeM. .15." Slide No. 5517.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral view of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

Baeonoma euphanes Meyrick

Plate 72, Figure 1

Baeonoma euphanes Meyrick, 1916, Exotic Microlepidoptera, 1: 507.

"♂. 9–10 mm. . . . French Guiana, St. Jean, R. Maroni; two specimens. Very similar to *leucodelta*, but in that species the white apical patch and the dorsal blotch are larger, and veins 7 and 8 are stalked."

Lectotype: One of the above indicated males both of which are without abdomens.

Figure 1, left wings.

Baeonoma favillata (Meyrick), new combination

Plate 72, Figures 2–2b

Stenoma favillata Meyrick, 1915, Exotic Microlepidoptera, 1: 430.

"♂. 16 mm. . . . Peru, Pacaya, in July (*Mounsey*); one specimen."

Type: The specimen indicated above, with "M. .7.12" at the bottom of the pin-label. Slide No. 5461.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Baeonoma helotypa Meyrick

Plate 72, Figures 3–3b

Baeonoma helotypa Meyrick, 1916, Exotic Microlepidoptera, 1: 508.

"♂♀. 14–18 mm. . . . French Guiana, R. Maroni; six specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana, LeM. .15." Slide No. 5520.

Figure 3, right wings (image reversed); 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Baeonoma holarga Meyrick

Plate 72, Figures 4–4b

Baeonoma holarga Meyrick, 1916, Exotic Microlepidoptera, 1: 509.

"♂. 19 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen indicated above, with "LeM. .15" at the bottom of the pin-label. Slide No. 5521.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

2b


2

2a

3b


3

3a


4

4a

4b

BAEONOMA



BAEONOMA

Baeonoma infamis Meyrick

Plate 73, Figures 1–1b

Baeonoma infamis Meyrick, 1925, Exotic Microlepidoptera, 3: 160.

"♀. 19–22 mm. . . . Brazil, Teffé, December (*Parish*); 2 ex."

Lectotype: The specimen measuring 22 mm. Slide No. 5524.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum.

Baeonoma leucodelta (Meyrick)

Plate 73, Figures 2–2a

Machimia leucodelta Meyrick, 1914, Exotic Microlepidoptera, 1: 184.

"♂. 7 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The specimen referred to above dated ".3.13". Slide No. 5519.

Figure 2, left wings; 2a, ventral view of male genitalia (right harpe out of focus) with aedeagus *in situ*.

Baeonoma orthozona Meyrick

Plate 73, Figures 3–3b

Baeonoma orthozona Meyrick, 1916, Exotic Microlepidoptera, 1: 508.

"♀. 13–14 mm. . . . ♂. 11–12. . . . French Guiana, R. Maroni; seven specimens (4♂, 3♀). I believe these, though at first sight very different looking, to be the sexes of the same species, but in case of error I constitute the ♀s type."

Lectotype: One of the above indicated females, all of which have the same data, with "LeM. .15" at the bottom of the pin-label. Slide No. 5518.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

Baeonoma suavis Meyrick

Plate 73, Figures 4–4b

Baeonoma suavis Meyrick, 1916, Exotic Microlepidoptera, 1: 541.

"♂. 15 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen indicated above with "LeM. .16" at the bottom of the pin-label. Slide No. 5523.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

CERCONOTA Meyrick

Cerconota Meyrick, 1915, Exotic Microlepidoptera, 1: 385. (*Typus generis: Cerconota tridesma* Meyrick, l.c., 386 [monotypy].)

Cerconota tridesma Meyrick

Plate 74, Figures 1–1d

Cerconota tridesma Meyrick, 1915, Exotic Microlepidoptera, 1: 386.

"♂. 22 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated ".3.13". Slide No. 4965.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral view of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.



1



1a

1b



1c



1d

CERCONOTA



1

1a



1b



2

2a

2b



3

3a

3b



4

4a

4b

CERCONOTA

Cerconota agraria (Meyrick), new combination

Plate 75, Figures 1–1b

Stenoma agraria Meyrick, 1925, Exotic Microlepidoptera, 3: 210.

"♂♀. 20–22 mm. . . . Bolivia, Prov. del Sara, Dept. Santa Cruz, 1,500 feet; 2 ex."

Lectotype: ♂, 22 mm. Slide No. 5203. Meyrick recorded a female but both specimens are males.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Cerconota atricassis (Meyrick), new combination

Plate 75, Figures 2–2b

Stenoma atricassis Meyrick, 1916, Exotic Microlepidoptera, 1: 530.

"♂. 19–20 mm. . . . French Guiana, R. Maroni; four specimens."

Lectotype: One of the three remaining males with "LeM. .15" at the bottom of the pin-label. Slide No. 5372.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Cerconota brachyplaca (Meyrick)

Plate 75, Figures 3–3b

Ptilogenes brachyplaca Meyrick, 1926, Exotic Microlepidoptera, 3: 228.

"♂. 20–21 mm. . . . Brazil, Teffé, January (*Parish*); French Guiana, R. Maroni; 2 ex. Excessively like *palliata*, but certainly distinct by absence of subcostal hairr pencil of hindwings; . . ."

Lectotype: ♂, "Teffé, Brazil, Parish. 1.20." Slide No. 5531.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Cerconota bythochroa (Meyrick)

See Cerconota nimbosa (Zeller), p. 160.

Cerconota capnosphaera (Meyrick)

Plate 75, Figures 4–4b

Stenoma capnosphaera Meyrick, 1916, Exotic Microlepidoptera, 1: 516.

"♂. 17–18 mm. . . . French Guiana, Godebert, R. Maroni; two specimens."

Lectotype: ♂, "Godebert, R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5104.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Cerconota censoria (Meyrick), new combination

Plate 76, Figures 1–1b

Stenoma censoria Meyrick, 1915, Exotic Microlepidoptera, 1: 460.

Stenoma congressella (Walker) Meyrick *not* Walker, 1925, Exotic Microlepidoptera, 3: 192.

"♀. 20–21 mm. . . . British Guiana, Bartica, in January and February (*Parish*); two specimens."

Lectotype: The female dated ".1.13". Slide No. 5280. The second specimen is a male.

Meyrick (1925) suppressed his *censoria* in favour of *congressella* but his synonymy is based on a misdetermination. The signum of *censoria* is dumb-bell shaped and that of *congressella* is a small tri-lobed plate. I hereby remove *censoria* from synonymy.

Figure 1, left wings; 1a, detail of genital plate and ostium; 1b, dorsal view of eighth segment collar to show large median protuberance.

Cerconota consobrina (Meyrick), new combination

Plate 76, Figures 2–2c

Stenoma consobrina Meyrick, 1915, Exotic Microlepidoptera, 1: 460.

"♀. 20–21 mm. . . . British Guiana, Bartica, in January (*Parish*); two specimens."

Lectotype: One of the above indicated females both of which bear identical data and are dated "1.13". Slide No. 5279.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Cerconota curiata (Meyrick), new combination

Plate 76, Figures 3–3c

Stenoma curiata Meyrick, 1929, Trans. Ent. Soc. London, 76: 515

"♀. 22 mm. . . . Panama, Taboga I., 200–1,000 feet, September, at light; 3 ex. Allied to *satyropa*, . . ."

Type: The specimen designated type in the British Museum, dated "15.9.24". A small white label bears the inscription "P825". Slide No. 5582.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Cerconota ebenocista (Meyrick)

Plate 76, Figures 4–4b

Ptilogenes ebenocista Meyrick, 1928, Exotic Microlepidoptera, 3: 433.

"♂. 23–24 mm. . . . French Guiana, R. Maroni; 8 ex. Also 1 from Brazil, Nova Paraigo, February (*Parish*), forewings brownish-tinged, hindwings ochreous-grey-whitish, terminal third grey, but not specifically distinct."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .16." Slide No. 5530.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia (uncus broken) with aedeagus removed.



CERCONOTA



CERCONOTA

Cerconota eriacma (Meyrick), new combination

Plate 77, Figures 1–1b

Stenoma eriacma Meyrick, 1915, Exotic Microlepidoptera, 1: 456.

"♂. 15 mm. . . . British Guiana, Bartica, in December (*Parish*); one specimen."

Type: The above indicated specimen dated "12.12". Slide No. 5232.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Cerconota erotarcha (Meyrick), new combination

See Cerconota rosacea (Butler), p. 167.

Cerconota fermentata (Meyrick), new combination

Plate 77, Figures 2–2b

Stenoma fermentata Meyrick, 1916, Exotic Microlepidoptera, 1: 539.

"♂♀. 29–32 mm. . . . British Guiana, R. Maroni; four specimens."

Lectotype: A male with "LeM. .16" at the bottom of the pin-label. Slide No. 5513.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Cerconota figularis (Meyrick), new combination

Plate 77, Figures 3–3b

Stenoma figularis Meyrick, 1918, Exotic Microlepidoptera, 2: 204.

"♂. 17 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .16" at the bottom of the pin-label. Slide No. 5270.

Figure 3, right wings (image reversed); 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Cerconota flexibilis (Meyrick), new combination

Plate 77, Figures 4–4b

Stenoma flexibilis Meyrick, 1916, Exotic Microlepidoptera, 1: 538.

"♂. 23–24 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, 23 mm., with "LeM. .15" at the bottom of the pin-label. Slide No. 5511.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Cerconota fulminata (Meyrick), new combination

Plate 78, Figures 1–1b

Stenoma fulminata Meyrick, 1916, Exotic Microlepidoptera, 1: 539.

"♂ 19 mm., ♀ 22 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂. Both specimens bear identical data with "LeM. .16" at the bottom of the pin-label. Slide No. 5509.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Cerconota fusigera (Meyrick)

Plate 78, Figures 2–2b

Stenoma fusigera Meyrick, 1915, Exotic Microlepidoptera, 1: 476.

"♂♀. 24–27 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: The male with "LeM. .15" at the bottom of the pin-label. Slide No. 5208.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Cerconota hexascia (Meyrick), new combination

Plate 78, Figures 3–3b

Stenoma hexascia Meyrick, 1925, Exotic Microlepidoptera, 3: 215.

"♂. 15 mm. . . . Brazil, Teffé, December, January (*Parish*); 2 ex."

Lectotype: ♂, "Teffé, Brazil, Parish. 1.20." Slide No. 5434. This species is atypical for *Cerconota* but is better placed here than elsewhere.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Cerconota horometra (Meyrick), new combination

Plate 78, Figures 4–4b

Stenoma horometra Meyrick, 1925, Exotic Microlepidoptera, 3: 222.

"♂. 25 mm. . . . Peru, Rio Napo, May (*Parish*); 2 ex."

Lectotype: One of the above indicated males both of which bear identical data and are dated "5.20". Slide No. 5485.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a


1b


2

2a


2b


3

3a

3b

4

4a


4b

1


1a


1b


2

2a


2b

2c

3

3a

3b

3c


4


4a


4b

CERCONOTA

Cerconota hydrelaeas (Meyrick), new combination

Plate 79, Figures 1–1b

Stenoma hydrelaeas Meyrick, 1931, Exotic Microlepidoptera, 4: 33.

"♂. 18 mm. . . . French Guiana, R. Maroni; 1 ex. (Coll. Paravicini). Possibly allied to *porphyrastis*."

Type: The above indicated specimen with "M905" on a small white label. Slide No. 5585.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Cerconota impressella (Walker)

Plate 79, Figures 2–2c

Cryptolechia impressella Walker, 1864, List of the specimens of lepidopterous insects in the collection of the British Museum, 29: 720.

Agriophara prasoleuca Meyrick, 1916, Exotic Microlepidoptera, 1: 483.

[prasoleuca]

"♂♀. 23–25 mm. . . . French Guiana, R. Maroni; six specimens."

Lectotype: ♀, "R. Maroni, Fr. Guiana, LeM. .16." Slide No. 5649.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and signum.

Cerconota languescens (Meyrick)

Plate 79, Figures 3–3c

Stenoma languescens Meyrick, 1915, Exotic Microlepidoptera, 1: 458.

"♀. 20 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen."

Type: The above indicated specimen dated ".1.13". Slide No. 5369.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Cerconota lembifera (Meyrick), new combination

Plate 79, Figures 4–4b

Stenoma lembifera Meyrick, 1915, Exotic Microlepidoptera, 1: 440.

"♂♀. 16–17 mm. . . . British Guiana, Mallali, in March (*Parish*); Dutch Guiana, Onoribo, in March; two specimens. These appear to be truly sexes of the same species, though from different localities."

Lectotype: ♂, "Onoribo, Dutch Guiana. E. .3.93." Slide No. 5394.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Cerconota leucosaris (Meyrick), new combination

See Cerconota siraphora (Meyrick), p. 168.

Cerconota minna (Busck)

Plate 80, Figures 1–1b

Stenoma minna Busck, 1914, Proc. U.S. Nat. Mus., 47: 52.
Stenoma orthridia Meyrick, 1916, Exotic Microlepidoptera, 1: 520.

[orthridia]

"♂♀. 23–25 mm. . . . French Guiana, St. Jean, R. Maroni; British Guiana, R. Demerara; eighteen specimens. The very similar *collybista* from Peru has no subdorsal hair-pencil on forewings of ♂, . . ."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 5636.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Cerconota myrodora (Meyrick), new combination

Plate 80, Figures 2–2b

Stenoma myrodora Meyrick, 1925, Exotic Microlepidoptera, 3: 197.

"♂♀. 21–22 mm. . . . Brazil, Teffé, January (*Parish*); French Guiana, R. Maroni; 4 ex."

Lectotype: ♂, "Teffé, Brazil, Parish. 1.20." Slide No. 5209.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Cerconota nimbosa (Zeller)

Plate 80, Figures 3–3b; 4c

Cryptolechia nimbosa Zeller, 1877, Hor. Soc. Ent. Ross., 13: 271.
Gonioterma bythochroa Meyrick, 1915, Exotic Microlepidoptera, 1: 385.

[bythochroa]

"♀. 29 mm. . . . British Guiana, Bartica, in February (*Parish*); one specimen."

Type: The above indicated specimen dated ".2.13". Slide No. 5631.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium; 4c, signum.

Cerconota noverca (Meyrick), new combination

Plate 80, Figures 4–4b

Agriophara noverca Meyrick, 1916, Exotic Microlepidoptera, 1: 484.

"♀. 30 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen denoted above with "LeM. .15" at the bottom of the pin-label. Slide No. 5528. The middle portion of the ductus bursae contains foreign matter.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum.



CERCONOTA

1

1a

1b



2

2a

2b



3

3a

3b



4

4a

4b

CERCONOTA

Cerconota nymphas (Meyrick), new combination

Plate 81, Figures 1–1b

Stenoma nymphas Meyrick, 1916, Exotic Microlepidoptera, 1: 534.

"♂♀. 14–16 mm. . . . French Guiana, R. Maroni; four specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5390.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Cerconota omphacopa (Meyrick), new combination

Plate 81, Figures 2–2b

Stenoma omphacopa Meyrick, 1931, Exotic Microlepidoptera, 4: 39.

"♂. 18 mm. . . . Bolivia, Cochabamba (*Germain*); 1 ex. (Coll. Paravicini). Perhaps related to *camarodes*."

Type: The specimen indicated above dated "1888–89" and with a small white label with the inscription "M808". Slide No. 5589.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Cerconota oceanitis (Meyrick), new combination

Plate 81, Figures 3–3b

Stenoma oceanitis Meyrick, 1916, Exotic Microlepidoptera, 1: 519.

"♂♀. 20–21 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: ♂. All specimens have identical data and are labelled "R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5201.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Cerconota orthridia (Meyrick)

See Cerconota minna (Busck), p. 160.

Cerconota phaeophanes (Meyrick), new combination

Plate 81, Figures 4–4b

Stenoma phaeophanes Meyrick, 1912, Trans. Ent. Soc. London, 1911: 714.

"♂♀. 24–27 mm. . . . Colombia, San Antonio, 5,800 feet, in November and December; two specimens."

Lectotype: The male dated ".11.07". Slide No. 5282.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Cerconota phyllocosma (Meyrick), new combination

Plate 82, Figures 1–1b

Stenoma phyllocosma Meyrick, 1916, Exotic Microlepidoptera, 1: 520.

"♂. 20–21 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: ♂, with "LeM. .16" at the bottom of the pin-label. Slide No. 5231. One of the original specimens is a female.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Cerconota physotricha (Meyrick), new combination

Plate 82, Figures 2–2b

Stenoma physotricha Meyrick, 1915, Exotic Microlepidoptera, 1: 455.

"♂. 23 mm. . . . Venezuela, Carupano, in December; one specimen."

Type: The above indicated specimen with "E. .12.91" at the bottom of the pin-label. Slide No. 5207.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Cerconota prasoleuca (Meyrick)

See Cerconota impressella (Walker), p. 159.

Cerconota pseudacma (Meyrick)

See Cerconota recurvella (Walker), p. 167.

Cerconota ptilosema Meyrick

Plate 82, Figures 3–3b

Cerconota ptilosema Meyrick, 1918, Exotic Microlepidoptera, 2: 201.

"♂. 18 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated male with "LeM. .16" at the bottom of the pin-label. Slide No. 4966.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Cerconota punicea (Meyrick), new combination

Plate 82, Figures 4–4b

Stenoma punicea Meyrick, 1916, Exotic Microlepidoptera, 1: 518.

"♂. 16–17 mm. . . . French Guiana, Godebert; two specimens."

Lectotype: The one remaining male in the Meyrick collection labelled "Godebert, R. Maroni, Fr. Guiana. LeM. .15". Slide No. 5199.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1


1a


1b

2

2a

2b


3

3a


3b


4

4a

4b

CERCONOTA

1

4c

1a



2

2a

2b



3

3a

3b



4

4a



4b

CERCONOTA

Cerconota recurrens (Meyrick), new combination

Plate 83, Figures 1–1a

Stenoma recurrens Meyrick, 1925, Exotic Microlepidoptera, 3: 210.

"♂♀. 20–22 mm. . . . Bolivia, Prov. del Sara, Dept. Santa Cruz, 1,500 feet; 4 ex."

Lectotype: ♂. Three of the original specimens are in the Meyrick collection and all bear identical data. Slide No. 5202.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Cerconota recurvella (Walker)

Plate 83, Figures 2–2b

Cryptolechia recurvella Walker, 1864, List of the specimens of lepidopterous insects in the collection of the British Museum, 29: 717.

Stenoma pseudacma Meyrick, 1918, Exotic Microlepidoptera, 2: 203.

[pseudacma]

"♂. 28 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .16" at the bottom of the pin-label. Slide No. 5672. The abdomen of Walker's type is missing but the two species agree in all other details.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Cerconota robiginosa (Meyrick), new combination

Plate 83, Figures 3–3b

Stenoma robiginosa Meyrick, 1925, Exotic Microlepidoptera, 3: 215.

"♂. 16 mm. . . . Brazil, Manaos, November (*Parish*); 1 ex."

Type: The specimen indicated above dated "11.19". Slide No. 5437.

Figure 3, right wings (image reversed); 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Cerconota rosacea (Butler), new combination

Plate 83, Figures 4–4c

Cryptolechia rosacea Butler, 1877, Cistula Entomologica, 2: 192.

Stenoma erotarcha Meyrick, 1915, Exotic Microlepidoptera, 1: 471 (new synonymy).

[erotarcha]

"♀. 29–31 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: ♀, 30 mm. All specimens have identical data with "LeM. .15" at the bottom of the pin-label. Slide No. 5681. The types of both species agree in all details including the genitalia.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, detail of genital plate and ostium; 4c, signum.

Cerconota siraphora (Meyrick), new combination

Plate 84, Figures 1–1b; 2–2b

Stenoma siraphora Meyrick, 1915, Exotic Microlepidoptera, 1: 475.
Stenoma leucosaris Meyrick, 1925, Exotic Microlepidoptera, 3: 196 (new synonymy).

[siraphora]

"♂. 20 mm. . . . French Guiana, R. Maroni; one specimen. Allied to *trichocrossa* and *emma*, which have similar fringes."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5194.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

[leucosaris]

"♂. 20 mm. . . . Peru, Papitinga, June (*Parish*); 1 ex. Extremely similar to *siraphora*, but apex of forewings more rounded. . . ."

Type: The above indicated specimen dated ".6.20". Slide No. 5193.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Cerconota sphragidopis (Meyrick), new combination

Plate 84, Figures 3–3c

Stenoma sphragidopis Meyrick, 1915, Exotic Microlepidoptera, 1: 431.

"♀. 25–26 mm. . . . British Guiana, Bartica, in December (*Parish*); three specimens."

Lectotype: ♀, dated "12.12". Only one of the original specimens, the lectotype, remains in the Meyrick collection. Slide No. 5506.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Cerconota stylonota (Meyrick), new combination

Plate 84, Figures 4–4b

Stenoma stylonota Meyrick, 1915, Exotic Microlepidoptera, 1: 447.

"♂. 17 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The specimen indicated above dated ".3.13". Slide No. 5381.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

1

1a



1b

2

2a

2b

3

3a

3b

3c

4

4a

4b

CERCONOTA



1



1a

1b



2

2a



2b

3

3a

3b



4



4a

4b

CERCONOTA

Cerconota tholodes (Meyrick), new combination

Plate 85, Figures 1–1b

Stenoma tholodes Meyrick, 1915, Exotic Microlepidoptera, 1: 457.

"♀. 22 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated "3.13". Slide No. 5210.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium.

Cerconota tricharacta (Meyrick), new combination

Plate 85, Figures 2–2b

Stenoma tricharacta Meyrick, 1925, Exotic Microlepidoptera, 3: 199.

"♂♀. 19–21 mm. . . . Brazil, Teffé, Obidos, August, September, January; French Guiana, R. Maroni; 23 ex."

Lectotype: ♂, "Teffé, Brazil, Parish. 1.20." Slide No. 5239.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Cerconota trichoneura (Meyrick), new combination

Plate 85, Figures 3–3b

Stenoma trichoneura Meyrick, 1913, Trans. Ent. Soc. London, 1913: 189.

"♂♀. 18–21 mm. . . . Venezuela, Ciudad Bolivar, in May and June; eleven specimens."

Lectotype: ♂, dated "6.04". All specimens bear identical data except date. Slide No. 5192.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Cerconota trochistis (Meyrick), new combination

Plate 85, Figures 4–4b

Stenoma trochistis Meyrick, 1916, Exotic Microlepidoptera, 1: 518.

Stenoma conturbatella (Walker) Meyrick, *not* Walker 1925, Exotic Microlepidoptera, 3: 192.

"♂. 23–28 mm. . . . French Guiana, St. Jean, R. Maroni; three specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana." Only the lectotype bears the designation "St. Jean". Slide No. 5253.

Meyrick suppressed his name for *conturbatella* Walker, but a close examination of Walker's type (also that of *illucidella* Walker [=*conturbatella* Walker]), reveals that the two are abundantly distinct. The largest of Walker's specimens measures only 19 mm. I hereby remove Meyrick's *trochistis* from synonymy.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Cerconota trymalopa (Meyrick), new combination

Plate 86, Figures 1–1b

Stenoma trymalopa Meyrick, 1925, Exotic Microlepidoptera, 3: 211.

"♂. 30 mm. . . . Colombia, San Antonio, 6,600 feet, February, 1 ex."

Type: The above indicated specimen dated ".2.20". Slide No. 5288.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Cerconota tumulata (Meyrick), new combination

Plate 86, Figures 2–2b

Stenoma tumulata Meyrick, 1916, Exotic Microlepidoptera, 1: 538.

"♂. 22 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5510.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.



CERCONOTA



1

1a



1b

1c



1d

CHLAMYDASTIS

CHLAMYDASTIS Meyrick

Plate 87, Figures 1–1d

Chlamydastis Meyrick, 1916, Exotic Microlepidoptera, 1: 481. (*Typus generis: Stenoma lactis* Busck, 1911, Proc. U.S. Nat. Mus., 40: 216 [by original designation].)

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed. Slide No. 9700.

Chlamydastis anamochla (Meyrick)

Plate 88, Figures 1–1b

Ptilogenes anamochla Meyrick, 1929, Trans. Ent. Soc. London, 76: 517.

"♂. 11 mm. . . . Colombia, Gorgona I., 200 feet, November, at light; 1 ex. Allied to *dryosphaera*."

Type: The specimen so marked in the British Museum, labelled "Gorgona Island, Colombia, at light, 200 ft., 19.11.24. St. George Expedn. C. L. Collenette." A small white label bears the inscription "P987". Slide No. 5607.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Chlamydastis ancalota (Meyrick)

Plate 88, Figures 2–2b

Agriophara ancalota Meyrick, 1916, Exotic Microlepidoptera, 1: 486.

"♂ 19 mm., ♀ 23 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, with "LeM. .15" at the bottom of the pin-label. Slide No. 5550.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Chlamydastis apoclina (Meyrick)

Plate 88, Figures 3–3b

Ptilogenes apoclina Meyrick, 1929, Trans. Ent. Soc. London, 76: 516.

"♂. 18–19 mm. . . . Colombia, Gorgona I., 200 feet, October, November, at light; 5 ex. Near *lichenias*."

Type: The specimen so designated in the British Museum, labelled "Gorgona Island, Colombia, at light, 200 ft., 19.11.24. St. George Expedn. C. L. Collenette." A small white label bears the inscription "P898". Slide No. 5608.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed; and section of denuded abdomen to show tuft from intersegmental membrane between segments seven and eight.

Chlamydastis argocymba (Meyrick)

Plate 88, Figures 4–4b

Ptilogenes argocymba Meyrick, 1926, Exotic Microlepidoptera, 3: 232.

"♂♀. 16–19 mm. . . . Brazil, Parintins, October (*Parish*); 4 ex."

Lectotype: ♂, 16 mm., dated "10.19". Three of the four original specimens remain in the Meyrick collection and all bear identical data. Slide No. 5600.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a

1b


2

2a

2b


3

3a

3b

4

4a

4b

CHLAMYDASTIS


1

1a


1b

1c


2


2a

2b


3

3a


3b

4

4a


4b

CHLAMYDASTIS

Chlamydastis aphrogenes (Meyrick)

Plate 89, Figures 1–1c

Agriophara aphrogenes Meyrick, 1915, Exotic Microlepidoptera, 1: 407.

"♂. 20–22 mm. . . . British Guiana, Bartica and Mallali, in February and March (*Parish*); two specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. .2.13." Slide No. 5531. This species has appeared as a synonym of *C. trastices* (Busck) from which it is amply distinct. I hereby remove *aphrogenes* from synonymy.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed; 1c, denuded abdomen to show modification of seventh and eighth segments.

Chlamydastis batrachopis (Meyrick)

Plate 89, Figures 2–2b

Agriophara batrachopis Meyrick, 1913, Trans. Ent. Soc. London, 1913: 182.

"♂. 34 mm. . . . Peru, Chanchamayo, 3,500 feet; one specimen."

Type: The specimen indicated above dated ".11". Slide No. 5534. This species appears to be only a race of *smodicopa*.

Figure 2, left wings; 2a, ventral view of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Chlamydastis bifida (Meyrick)

Plate 89, Figures 3–3b

Agriophara bifida Meyrick, 1916, Exotic Microlepidoptera, 1: 541.

"♂. 17 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen indicated above labelled "R. Maroni, Fr. Guiana. LeM. .16." Slide No. 5558.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Chlamydastis chionoptila (Meyrick)

Plate 89, Figures 4–4b

Ptilogenes chionoptila Meyrick, 1926, Exotic Microlepidoptera, 3: 229.

"♂. 19 mm. . . . Brazil, Teffé, December (*Parish*); 1 ex."

Type: The specimen indicated above dated "12.19". Slide No. 5537.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Chlamydastis byssophanes (Meyrick)

Plate 90, Figures 1–1b

Ptilogenes byssophanes Meyrick, 1926, Exotic Microlepidoptera, 3: 230.

"♂♀. 16–21 mm. . . . Brazil, Teffé, December, January; Peru, Rio Napo, May (*Parish*); 7 ex."

Lectotype: ♂, "Teffé, Brazil, Parish. 1.20." Slide No. 5556.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Chlamydastis caecata (Meyrick)

Plate 90, Figure 2

Agriophara caecata Meyrick, 1916, Exotic Microlepidoptera, 1: 487.

"♂. 24 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen denoted above with "LeM. .15" at the bottom of the pin-label. The type is without abdomen.

Figure 2, left wings.

Chlamydastis chionosphena (Meyrick)

Plate 90, Figures 3–3c

Ptilogenes chionosphena Meyrick, 1931, Exotic Microlepidoptera, 4: 45.

"♀. 17 mm. . . . French Guiana, R. Maroni, 1 ex. (Coll. Paravicini). Between *mysticopis* and *leucoplasta*."

Type: The specimen indicated above marked "Holotype" in the British Museum. A small white label bears the inscription "M823". Slide No. 5609.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium; 3c, signa.

Chlamydastis chloroloba (Meyrick)

Plate 90, Figures 4–4b

Stenoma chloroloba Meyrick, 1915, Exotic Microlepidoptera, 1: 439.

"♂. 18 mm. . . . Peru, Contamano, R. Ucayali, in December (*Mounsey*); one specimen."

Type: The above indicated specimen dated "12.12". Slide No. 5559.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

1

1a


1b

2


3b


3

3a


3c

4

4a

4b

CHLAMYDASTIS


1

2c


2

2a

2b


3

3a

3b


4

4a

4b

CHLAMYDASTIS

Chlamydastis chlorosticta (Meyrick)

Plate 91, Figure 1

Agriophara chlorosticta Meyrick, 1913, Trans. Ent. Soc. London, 1913: 184.

"♂. 30 mm. . . . Peru, Chanchamayo; one specimen."

Type: The above indicated specimen with "R. .11" at the bottom of the pin-label. The abdomen of the type is missing.

Figure 1, left wings.

Chlamydastis deflua (Meyrick)

Plate 91, Figures 2–2c

Ptilogenes deflua Meyrick, 1918, Exotic Microlepidoptera, 2: 210.

"♂♀. 16–18 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: ♂, 16 mm., with "LeM. .15" at the bottom of the pin-label. Slide No. 5564.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed; 2c, denuded abdomen to show abdominal tufts.

Chlamydastis diorista (Meyrick)

Plate 91, Figures 3–3b

Ptilogenes diorista Meyrick, 1929, Trans. Ent. Soc. London, 76: 517.

"♂. 12 mm. . . . Colombia, Gorgona I., 200 feet, July, November, at light; 2 ex."

Type: The specimen in the British Museum so marked, and labelled "Gorgona Island, Colombia, at light, 200 ft. 19.11.24. St. George Expedn. C. L. Collenette." Slide No. 5610.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Chlamydastis disticha (Meyrick)

Plate 91, Figures 4–4b

Agriophara disticha Meyrick, 1916, Exotic Microlepidoptera, 1: 488.

"♀. 15 mm. . . . French Guiana, St. Jean, R. Maroni, in July; one specimen."

Type: The above indicated specimen with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5557.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum.

Chlamydastis complexa (Meyrick)

Plate 92, Figures 1–1c

Agriophara complexa Meyrick, 1916, Exotic Microlepidoptera, 1: 542.

"♂. 24–26 mm. . . . French Guiana, R. Maroni; three specimens. Very similar to *ponderata*, but larger, . . ."

Lectotype: ♂, 25 mm. "R. Maroni; Fr. Guiana. LeM. .15." Slide No. 5568.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed; 1c, section of denuded abdomen to show posterior tufts.

Chlamydastis crateroptila (Meyrick)

Plate 92, Figures 2–2b

Ptilogenes crateroptila Meyrick, 1918, Exotic Microlepidoptera, 2: 211.

"♂. 14 mm. . . . French Guiana, St. Jean, R. Maroni, July; one specimen."

Type: The above indicated specimen dated ".7.15". Slide No. 5565. Each harpe with two or three strong, hooked setae from base, but only one on each side remained attached.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Chlamydastis cystiodes (Meyrick)

Plate 92, Figures 3–3b

Agriophara cystiodes Meyrick, 1916, Exotic Microlepidoptera, 1: 489.

"♂♀. 16–18 mm. . . . French Guiana, R. Maroni; five specimens."

Lectotype: ♂, 16 mm. The five specimens of the original series bear identical data with "LeM. .15" at the bottom of the pin-label. Slide No. 5596.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Chlamydastis deflexa (Meyrick)

Plate 92, Figures 4–4b

Agriophara deflexa Meyrick, 1916, Exotic Microlepidoptera, 1: 489.

"♂. 15–16 mm. . . . French Guiana, St. Jean, R. Maroni; six specimens."

Lectotype: ♂, 15 mm. "St. Jean, R. Maroni, Fr. Guiana, LeM. .4.15." Slide No. 5572.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a

1b

1c


2

2a


2b

3

3a

3b


4

4a

4b

CHLAMYDASTIS


1


1b

1a

1c


2

2a

2 b


3

3a


3 b


4

4a

4 b

CHLAMYDASTIS

Chlamydastis dryosphaera (Meyrick)

Plate 93, Figures 1–1c

Ptilogenes dryosphaera Meyrick, 1926, Exotic Microlepidoptera, 3: 231.

"♂. 15 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex."

Type: The specimen indicated above dated "1.20". Slide No. 5566.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed; 1c, section of denuded abdomen to show posterior tuft.

Chlamydastis forcipata (Meyrick)

Plate 93, Figures 2–2b

Agriophara forcipata Meyrick, 1913, Trans. Ent. Soc. London, 1913: 183.

"♂. 23 mm. . . . Colombia, Naranjito, R. Dagua, 3,900 feet; one specimen."

Type: The specimen indicated above dated ".6.08". Slide No. 5545.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Chlamydastis ichthyodes (Meyrick)

Plate 93, Figures 3–3b

Ptilogenes ichthyodes Meyrick, 1926, Exotic Microlepidoptera, 3: 231.

"♂. 19 mm. . . . Peru, Rio Napo, May (*Parish*); 1 ex."

Type: The specimen referred to above dated "5.20". Slide No. 5570.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Chlamydastis illita (Meyrick)

Plate 93, Figures 4–4b

Ptilogenes illita Meyrick, 1926, Exotic Microlepidoptera, 3: 232.

"♂. 14 mm. . . . Peru, Rio Napo, May (*Parish*); 1 ex."

Type: The above indicated specimen dated "5.20". Slide No. 5598.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Chlamydastis epophrysta (Meyrick)

Plate 94, Figures 1–1c

Stenoma epophrysta Meyrick, 1909, Trans. Ent. Soc. London, 1909: 33.

"♂. 23 mm. . . . Peru, Pozuzo; one specimen."

Type: The specimen indicated above with "D. 2,500′ .06" at the bottom of the pin-label. Slide No. 5547.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed; 1c, section of denuded abdomen to show posterior tufts.

Chlamydastis funicularis (Meyrick)

Plate 94, Figures 2–2b

Ptilogenes funicularis Meyrick, 1926, Exotic Microlepidoptera, 3: 228.

"♀. 23 mm. . . . French Guiana, R. Maroni; 1 ex."

Type: The specimen indicated above with "LeM. .16" at the bottom of the pin-label. Slide No. 5527.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum.

Chlamydastis galeomorpha (Meyrick)

Plate 94, Figures 3–3b

Ptilogenes galeomorpha Meyrick, 1931, Exotic Microlepidoptera, 4: 44.

"♀. 24 mm. . . . Brazil, Serra de Communaty, Pernambuco; 1 ex. (Coll. Paravicini). Allied to *ommatopa*."

Type: The above indicated specimen dated "1.2.1893" and with a small label bearing the inscription "M355". Slide No. 5611.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum.

Chlamydastis hemichlora (Meyrick)

Plate 94, Figures 4–4b; 3c

Agriophara hemichlora Meyrick, 1916, Exotic Microlepidoptera, 1: 487.

Ptilogenes scutellata (Meyrick), Meyrick, *not* Meyrick, 1926, Exotic Microlepidoptera, 3: 230.

"♂ 26–28 mm., ♀ 38 mm. . . . French Guiana, R. Maroni; twelve specimens."

Lectotype: ♂, 26 mm., with "LeM. .15" at the bottom of the pin-label. Slide No. 5554.

Meyrick suppressed *hemichlora* as a synonym of *scutellata* from which it is amply distinct as a comparison of the genitalia of *hemichlora* and *scutellata* (Plate 102, 1a, 1b) will demonstrate. I hereby remove *hemichlora* from synonymy.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed; 3c, eighth sternite with posterior tuft.


1

1a


1b


1c


2

2a


2b


3

3a

3b

3c


4

4a

4b



1

1a

1b



2

2a

2b



3

3a

3b



4

4a



4b

CHLAMYDASTIS

Chlamydastis inspectrix (Meyrick)

Plate 95, Figures 1–1b

Agriophara inspectrix Meyrick, 1916, Exotic Microlepidoptera, 1: 542.

"♂♀. 17–18 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 5594.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Chlamydastis leptobelisca (Meyrick)

Plate 95, Figures 2–2b

Ptilogenes leptobelisca Meyrick, 1929, Trans. Ent. Soc. London, 76: 518.

"♂. 20–22 mm. . . . Colombia, Gorgona I., 200 feet, July, October, November, at light; 13 ex. Allied to *complexa*, but not very closely."

Type: The specimen in the British Museum so marked, labelled "Gorgona Island, Colombia, at light, 200 ft., 15.10.24. St. George Expedn. C. L. Collenette." A small white label bears the inscription "P844". Slide No. 5612.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Chlamydastis melanometra (Meyrick)

Plate 95, Figures 3–3b

Ptilogenes melanometra Meyrick, 1926, Exotic Microlepidoptera, 3: 229.

"♂. 23 mm. . . . Colombia, San Antonio, 6,600 feet, February; 1 ex."

Type: The above indicated specimen dated "2.20". Slide No. 5541.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Chlamydastis melanonca (Meyrick)

Plate 95, Figures 4–4b

Agriophara melanonca Meyrick, 1915, Exotic Microlepidoptera, 1: 408.

"♀. 21 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The specimen indicated above dated ".3.13". Slide No. 5560.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum.

Chlamydastis leucoplasta (Meyrick)

Plate 96, Figures 1–1b

Ptilogenes leucoplasta Meyrick, 1926, Exotic Microlepidoptera, 3: 232.

"♂♀. 13 mm. . . . Brazil, Teffé, December; Peru, Iquitos, May (*Parish*); 2 ex."

Lectotype: ♂, "Teffé, Brazil, Parish. 12.19." Slide No. 5601.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Chlamydastis leucoptila (Meyrick)

Plate 96, Figures 2–2b

Ptilogenes leucoptila Meyrick, 1918, Exotic Microlepidoptera, 2: 210.

"♀. 21 mm. . . . French Guiana, St. Jean, R. Maroni, July; two specimens."

Lectotype: ♀, 21 mm., with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5551.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum.

Chlamydastis lichenias (Meyrick)

Plate 96, Figures 3–3c

Agriophara lichenias Meyrick, 1916, Exotic Microlepidoptera, 1: 486.

"♂ 16–17 mm., ♀ 20–21 mm. . . . French Guiana, R. Maroni; four specimens."

Lectotype: ♂, 16 mm., with "LeM. .15" at the bottom of the pin-label. Slide No. 5349. The larger male is missing.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed; 3c, denuded abdomen to show posterior tufts.

Chlamydastis mysticopis (Meyrick)

Plate 96, Figures 4–4b

Ptilogenes mysticopis Meyrick, 1926, Exotic Microlepidoptera, 3: 233.

"♂. 20–21 mm. . . . Peru, Oconeque, Carabaya, 7,000 feet, February (*Ockenden*); 2 ex."

Lectotype: ♂, 21 mm. Slide No. 5602.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



CHLAMYDASTIS


1

1a

1b

1c

2

2a


2b


3

3a

3b


4

4a

4b

CHLAMYDASTIS

Chlamydastis metacymba (Meyrick)

Plate 97, Figures 1–1c

Agriophara metacymba Meyrick, 1916, Exotic Microlepidoptera, 1: 489.

"♂. 15 mm. . . . French Guiana, St. Jean, R. Maroni, in July; one specimen."

Type: The above indicated specimen with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5595.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed; 1c, denuded abdomen to show internal eversible and posterior tufts.

Chlamydastis metacystis (Meyrick)

Plate 97, Figures 2–2b

Ptilogenes metacystis Meyrick, 1918, Exotic Microlepidoptera, 2: 211.

"♂♀. 14–17 mm. . . . French Guiana, R. Maroni; three specimens. Appears to differ essentially from *cystiodes* only in the colour of the abdomen and hindwings, . . ."

Lectotype: ♂, 15 mm., with "LeM. .15" at the bottom of the pin-label. One of the original specimens is missing. Slide No. 5597.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Chlamydastis mochlopa (Meyrick)

Plate 97, Figures 3–3b

Agriophara mochlopa Meyrick, 1915, Exotic Microlepidoptera, 1: 409.

"♀. 40 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen. Allied to *mendoron* and *acronitis*."

Type: The specimen indicated above dated "1.13". Slide No. 5538.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum.

Chlamydastis monastra (Meyrick)

Plate 97, Figures 4–4b

Stenoma monastra Meyrick, 1909, Trans. Ent. Soc. London, 1909: 33.

"♂. 28 mm. . . . Peru, Pozuzo; one specimen."

Type: The above indicated specimen with "D. 2,500′ .06" at the bottom of the pin-label. Slide No. 5546.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Chlamydastis niphochlaena (Meyrick)

Plate 98, Figures 1–1b

Ptilogenes niphochlaena Meyrick, 1926, Exotic Microlepidoptera, 3: 233.

"♂. 25 mm. . . . Peru, Rio Napo, May (*Parish*); 2 ex."

Lectotype: One of the above indicated specimens both of which bear identical data and are dated "5.20". Slide No. 5603.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Chlamydastis obnupta (Meyrick)

Plate 98, Figures 2–2a

Agriophara obnupta Meyrick, 1916, Exotic Microlepidoptera, 1: 483.

"♂ 21 mm., ♀ 24 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, with "LeM. .15" at the bottom of the pin-label. Slide No. 5529.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ* (composite photograph).

Chlamydastis ophiopa (Meyrick)

Plate 98, Figures 3–3c

Agriophara ophiopa Meyrick, 1916, Exotic Microlepidoptera, 1: 484.

"♂♀. 26–28 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, 26 mm., with "LeM. .15" at the bottom of the pin-label. Slide No. 5532. There are no females in the Meyrick collection.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed; 3c, expansible posterior tuft.

Chlamydastis oxyplaca (Meyrick)

Plate 98, Figures 4–4c

Ptilogenes oxyplaca Meyrick, 1929, Trans. Ent. Soc. London, 76: 518.

"♀. 18 mm. . . . Colombia, Gorgona I., 200 feet, July, at light; 1 ex. Next *argocymba*."

Type: The specimen so marked in the British Museum labelled "Gorgona Island, Colombia, at light, 200 ft., 10.7.24. St. George Expedn. C. L. Collenette." Slide No. 5613.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, inception of ductus seminalis; 4c, signum; a lightly sclerotised plate with a sharp, longitudinal, median keel the anterior point of which is greatly produced.

1

1a

1b

2

3a

2a


3


3c


3b


4

4a

4b

4c

CHLAMYDASTIS



1

1a

1b



2

2a



2b



3

3a

3b



4

4a

4b

CHLAMYDASTIS

Chlamydastis ommatopa (Meyrick)

Plate 99, Figures 1–1b

Ptilogenes ommatopa Meyrick, 1926, Exotic Microlepidoptera, 3: 230.

"♂. 26–29 mm. . . . Colombia, Rio Siato, 5,200 feet, and San Antonio, 6,600 feet; Bolivia, Andes, 10,000 feet; 4 ex."

Lectotype: ♂, "Andes, Bolivia, 10,000'." Slide No. 5553. Only the lectotype and one specimen from R. Siato remain in the Meyrick collection.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Chlamydastis paradromis (Meyrick)

Plate 99, Figures 2–2b

Agriophara paradromis Meyrick, 1915, Exotic Microlepidoptera, 1: 408.

"♂. 27–29 mm. . . . Colombia, San Antonio, 5,800 feet, in November and December; three specimens."

Lectotype: ♂, 29 mm., dated ".12.07". Slide No. 5542.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Chlamydastis perducta (Meyrick)

Plate 99, Figures 3–3b

Agriophara perducta Meyrick, 1916, Exotic Microlepidoptera, 1: 484.

"♂ 24–26 mm., ♀ 34–36 mm. . . . French Guiana, St. Jean, R. Maroni; five specimens. Intermediate between *acronitis* and *nestes*, . . ."

Lectotype: ♂, with "LeM. .16" at the bottom of the pin-label. Slide No. 5539.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Chlamydastis plocogramma (Meyrick)

Plate 99, Figures 4–4b

Agriophara plocogramma Meyrick, 1915, Exotic Microlepidoptera, 1: 407.

"♂ 15 mm., ♀ 18 mm. . . . British Guiana, Bartica and Mallali, in February and March (*Parish*); two specimens. Allied to *morbida*."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. .2.13." Slide No. 5563.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Chlamydastis poliopa (Meyrick)

Plate 100, Figures 1–1b

Agriophara poliopa Meyrick, 1916, Exotic Microlepidoptera, 1: 485.

"♂. 22–23 mm. . . . Colombia, San Antonio, 5,800 feet, in November; four specimens. Allied to *ptilopa*."

Lectotype: One of the above indicated specimens all of which bear identical data and are dated ".11.07". Slide No. 5543.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Chlamydastis ponderata (Meyrick)

Plate 100, Figures 2–2b

Agriophara ponderata Meyrick, 1916, Exotic Microlepidoptera, 1: 488.

"♂. 18–23 mm. . . . French Guiana, St. Jean, R. Maroni; eleven specimens."

Lectotype: ♂, 19 mm., "R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5573.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed (cucullus of left harpe missing).

Chlamydastis praenubila (Meyrick)

Plate 100, Figures 3–3b

Ptilogenes praenubila Meyrick, 1926, Exotic Microlepidoptera, 3: 231.

"♀. 14 mm. . . . Brazil, Teffé, December (*Parish*); 1 ex."

Type: The specimen indicated above dated "12.19". Slide No. 5569.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of ovipositor from ventral side and signum below.

Chlamydastis prudentula (Meyrick)

Plate 100, Figures 4–4b

Ptilogenes prudentula Meyrick, 1926, Exotic Microlepidoptera, 3: 232.

"♂. 14 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex."

Type: The specimen denoted above dated "3.20". Slide No. 5599.

Figure 4, right wings (image reversed); 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a

1b


2

2a

2b


3

3a

3b


4

4a



4b

CHLAMYDASTIS



1

1a

1b



2

2a

2c



2b



3



3a



3b



4

4a

4b

CHLAMYDASTIS

Chlamydastis ptilopa (Meyrick)

Plate 101, Figures 1–1b

Agriophara ptilopa Meyrick, 1913, Trans. Ent. Soc. London, 1913: 183.

"♂♀. 22–26 mm. . . . Colombia, San Antonio, 5,000 feet, in November; five specimens."

Lectotype: ♂, 23 mm., dated ".11.07". Meyrick records the altitude as "5,000 feet", an obvious misprint, since all five specimens are labelled "5,800'" as are other species from this locality. Slide No. 5544.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Chlamydastis scutellata (Meyrick)

Plate 101, Figures 2–2c

Agriophara scutellata Meyrick, 1916, Exotic Microlepidoptera, 1: 487.

"♂. 27–28 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, 27 mm. Both specimens have identical data with "LeM. .15" at the bottom of the pin-label. Slide No. 5552.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed; 2c, eighth tergite and sternite.

Chlamydastis smodicopa (Meyrick)

Plate 101, Figures 3–3b

Agriophara smodicopa Meyrick, 1915, Exotic Microlepidoptera, 1: 409.

"♂. 30 mm. . . . Brazil, Fonte Boa, in July; one specimen."

Type: The above indicated specimen with "K. .7.96" at the bottom of the pin-label. Slide No. 5536.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Chlamydastis stagnicolor (Meyrick)

Plate 101, Figures 4–4b

Ptilogenes stagnicolor Meyrick, 1926, Exotic Microlepidoptera, 3: 229.

"♂. 18–19 mm. . . . Brazil, Teffé, December, January (*Parish*); 2 ex."

Lectotype: One of the above mentioned males, dated "12.19". Slide No. 5540.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Chlamydastis steloglypta (Meyrick)

Plate 102, Figures 1–1b

Ptilogenes steloglypta Meyrick, 1931, Exotic Microlepidoptera, 4: 44.

"♂. 23–25 mm. . . . French Guiana, R. Maroni; 2 ex. Also 1 from same locality in Coll. Paravicini. Nearest *batrachopis*."

Lectotype: ♂, 23 mm., without date. Slide No. 5535.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Chlamydastis synedra (Meyrick)

Plate 102, Figures 2–2b

Agriophara synedra Meyrick, 1916, Exotic Microlepidoptera, 1: 485.

"♂. 18 mm. . . . Paraguay; one specimen."

Type: The specimen referred to above with "B. .06" at the bottom of the pin-label. Slide No. 5548.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Chlamydastis tritypa (Meyrick)

Plate 102, Figures 3–3a

Stenoma tritypa Meyrick, 1909, Trans. Ent. Soc. London, 1909: 32.

"♂. 25–28 mm. . . . Peru, Pozuzo; two specimens."

Lectotype: ♂, 28 mm., with "D. 2,500′ .06" at the bottom of the pin-label. Slide No. 5533.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.


1

1a

1b


2

2a

2b


3

3a

CHLAMYDASTIS



1

1a



1b



2

2a

2b

CHLAMYDASTIS

Chlamydastis truncatula (Meyrick)

Plate 103, Figures 1–1b

Agriophara truncatula Meyrick, 1913, Trans. Ent. Soc. London, 1913: 185.

"♂♀. 15–17 mm. . . . Venezuela, Ciudad Bolivar, in May; four specimens."

Lectotype: ♂, with "R. .5.04" at the bottom of the pin-label. Three of the four original specimens remain in the Meyrick collection. Slide No. 5396. This is not properly referable to *Chlamydastis* and probably belongs in *Orphnolechia*.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia (damaged) with aedeagus removed.

Chlamydastis ungulifera (Meyrick)

Plate 103, Figures 2–2b

Ptilogenes ungulifera Meyrick, 1929, Trans. Ent. Soc. London, 76: 517.

"♂. 21–23 mm. . . . Colombia, Gorgona I., 200 feet, October, November, at light; 2 ex. Next *ommatopa*."

Type: The specimen in the British Museum so marked, labelled "Gorgona Island, Colombia, at light, 200 ft., 20.11.24. St. George Expedn. C. L. Collenette." A small white label bears the inscription "P857". Slide No. 5614.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

DINOTROPA Meyrick

See Oecophoridae, to be dealt with in a later volume.

EUMITURGA Meyrick

Eumiturga Meyrick, 1925, Exotic Microlepidoptera, 3: 177. (*Typus generis: Eumiturga flocculosa* Meyrick, l.c.; 178 [monotypy].)

Eumiturga flocculosa Meyrick

Plate 104, Figures 1–1c

Eumiturga flocculosa Meyrick, 1925, Exotic Microlepidoptera, 3: 178.

"♂♀. 15–17 mm. . . . Brazil, Parintins, Teffé, October, December (*Parish*); 4 ex."

Lectotype: ♂, "Parintins, Brazil, Parish. .10–19." Slide No. 4976.

Figure 1, left wings; 1a, venation of right wings; 1b, head to show palpus; 1c, ventral view of male genitalia with aedeagus *in situ*.

Eumiturga promotella (Zeller)

Plate 104, Figures 2–2a

Cryptolechia promotella Zeller, 1877, Hor. Soc. Ent. Ross., 13: 294, pl. 3, fig. 84.
Eumiturga commutata Meyrick, 1926, Exotic Microlepidoptera, 3: 233 (new synonymy).

[commutata]

"♂♀. 15–17 mm. . . . Brazil, Manaos, Teffé, November, December (*Parish*); 8 ex."

Lectotype: ♂, "Teffé, Brazil, Parish. .12–19." Slide No. 5410.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.



1

1a



1b

1c

2

2a

EUMITURGA



FALCULINA

FALCULINA Zeller

Falculina antitypa Meyrick

Plate 105, Figures 1–1b

Falculina antitypa Meyrick, 1917, Exotic Microlepidoptera, 2: 58.

"♂ 24–30 mm., ♀ 36 mm. . . . French Guiana, R. Maroni; five specimens."

Lectotype: ♂, with "LeM. .16" at the bottom of the pin-label. Slide No. 4958.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Falculina lepidota Meyrick

Plate 105, Figures 2–2a

Falculina lepidota Meyrick, 1916, Exotic Microlepidoptera, 1: 482.

"♂ 28–29 mm., ♀ 36 mm. . . . French Guiana, R. Maroni; four specimens. The species of this genus (which have an aspect of reptilian mimicry) can be distinguished by the difference in form of forewings and colour of hindwings."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .15." Slide No. 4957.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

GONIOTERMA Walsingham

Gonioterma advocata (Meyrick), new combination

Plate 106, Figures 1–1a

Stenoma advocata Meyrick, 1916, Exotic Microlepidoptera, 1: 525; 1925, 3: 192 (as synonym of *pauperatella* Walker).

"♂. 25–26 mm. . . . French Guiana, R. Maroni; two specimens. Very similar indeed to *linteata*, but larger. . . ."

Lectotype: ♂, 25 mm., with "LeM. .15" at the bottom of the pin-label. Slide No. 5266. Tergites 2, 3 and 4 are densely spinulate and there are fine, deciduous hair-pencils between segments 6 and 7 and 7 and 8. The species is not a synonym of *pauperatella* (Walker) [=*linteata* Meyrick] as the genitalia amply demonstrate, and I hereby remove *advocata* from synonymy.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Gonioterma asthenopa (Meyrick), new combination

Plate 106, Figures 2–2b

Stenoma asthenopa Meyrick, 1916, Exotic Microlepidoptera, 1: 517.

"♂. 27 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5196. There is a strong, eversible, forked hair-pencil from first segment contained in an elongate cup-like pocket.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Gonioterma bolistis (Meyrick), new combination

Plate 106, Figures 3–3b

Stenoma bolistis Meyrick, 1925, Exotic Microlepidoptera, 3: 203.

"♂. 19 mm. . . . Brazil, Teffé, December (*Parish*); 1 ex."

Type: The above indicated specimen dated "12.19". Slide No. 5262.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Gonioterma choleroptila (Meyrick), new combination

Plate 106, Figures 4–4b

Stenoma choleroptila Meyrick, 1915, Exotic Microlepidoptera, 1: 467.

"♂♀. 16–21 mm. . . . British Guiana, Bartica, from December to February (*Parish*); four specimens."

Lectotype: The male dated "12.12". One specimen is missing. Slide No. 5264.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



GONIOTERMA

1



1a



1b

2

2a

2b



3



3a

3b



4



3c

4a

GONIOTERMA

Gonioterma crocoptila (Meyrick), new combination

Plate 107, Figures 1–1b

Stenoma crocoptila Meyrick, 1915, Exotic Microlepidoptera, 1: 466.

"♂. 16 mm. . . . British Guiana, Mallali, in March (*Parish*); two specimens. Very like *choleroptila* but easily distinguished by . . . the yellow hair-pencil of hindwings."

Lectotype: One of the above indicated males both of which bear identical data and are dated ".3.13". Slide No. 5268.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Gonioterma expansa (Meyrick), new combination

Plate 107, Figures 2–2b

Stenoma expansa Meyrick, 1915, Exotic Microlepidoptera, 1: 464.

"♂. 21 mm. . . . Brazil, Petropolis; one specimen."

Type: The above indicated specimen with "R. .06" at the bottom of the pin-label Slide No. 5277.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Gonioterma gubernata (Meyrick), new combination

Plate 107, Figures 3–3c

Stenoma gubernata Meyrick, 1915, Exotic Microlepidoptera, 1: 465.

"♂. 20 mm. . . . French Guiana, St. Jean, R. Maroni, in July; two specimens."

Lectotype: One of the above indicated specimens both of which have "LeM. .7.15" at the bottom of the pin-label. Slide No. 5260.

Figure 3, left wings; 3a, detail of sacculus; 3b, ventral view of male genitalia with aedeagus *in situ*; 3c, denuded abdomen to show modified segments.

Gonioterma linteata (Meyrick), new combination

See Gonioterma pauperatella (Walker), below.

Gonioterma pauperatella (Walker), new combination

Plate 107, Figures 4–4a

Cryptolechia pauperatella Walker, 1864, List of the specimens of lepidopterous insects in the collection of the British Museum, 29: 721.

Stenoma linteata Meyrick, 1916, Exotic Microlepidoptera, 1: 525; l.c., 1925, 3: 192 (as synonym of *pauperatella* Walker).

[linteata]

"♂. 21–22 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, 21 mm., with "LeM. .15" at the bottom of the pin-label. Slide No. 5265.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

Gonioterma notifera (Meyrick), new combination

Plate 108, Figures 1–1b

Stenoma notifera Meyrick, 1915, Exotic Microlepidoptera, 1: 465.

"♂. 18 mm. . . . Paraguay; one specimen."

Type: The above indicated specimen with "B. .06" at the bottom of the pin-label. Slide No. 5276.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Gonioterma ochrosaris (Meyrick), new combination

Plate 108, Figures 2–2a

Stenoma ochrosaris Meyrick, 1925, Exotic Microlepidoptera, 3: 199.

"♂♀. 17 mm. . . . Brazil, Teffé, January, ♂ (type); Obidos, September, ♀ (*Parish*); probably correctly associated as sexes, and ♀ easily separated from very similar *tricharacta* by ovipositor."

Type: The male designated above. Slide No. 5237.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Gonioterma periscelta (Meyrick), new combination

Plate 108, Figures 3–3a

Stenoma periscelta Meyrick, 1915, Exotic Microlepidoptera, 1: 457.

"♂. 16–18 mm. . . . Peru, Pacaya, in September (*Mounsey*); two specimens."

Lectotype: ♂, 18 mm., dated ".9.12". Slide No. 5982. The genitalia are badly mildewed.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Gonioterma phortax Meyrick

Plate 108, Figures 4–4a

Gonioterma phortax Meyrick, 1915, Exotic Microlepidoptera, 1: 383.

"♂♀. 15–20 mm. . . . British Guiana, Bartica, in December and January (*Parish*); Venezuela, Carupano, in December; four specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish, 12.12." Slide No. 5235. One specimen is missing. Meyrick transferred this species to *Stenoma*, and I now return it to its original genus.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

Gonioterma platycolpa (Meyrick), new combination

See Gonioterma seppiana (Cramer), p. 219.


1

1a

1b


2

2a

3

3a

4

4a

GONIOTERMA


1

1a


2

2a

3


3a


3b


4

4a

4b

GONIOTERMA

Gonioterma pleonastes (Meyrick), new combination

Plate 109, Figures 1–1a

Stenoma pleonastes Meyrick, 1915, Exotic Microlepidoptera, 1: 475.

"♂. 23 mm. . . . French Guiana, St. Jean, R. Maroni, in July; one specimen."

Type: The above indicated specimen with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5213.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Gonioterma projecta (Meyrick), new combination

Plate 109, Figures 2–2a

Stenoma projecta Meyrick, 1915, Exotic Microlepidoptera, 1: 466.

"♂. 19–20 mm. . . . French Guiana, St. Jean, Godebert, and Nouveau Chantier, R. Maroni, in January and July; five specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 5408.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Gonioterma seppiana (Cramer), new combination

Plate 109, Figures 3–3b; 4–4b

Phalaena Tortrix seppiana Cramer, 1781, Papillons Exotiques, 4: 135, Pl. 359, fig. H.
Stenoma platycolpa Meyrick, 1915, Exotic Microlepidoptera, 1: 465 (new synonymy).
Stenoma sceptrifera Meyrick, 1916, Exotic Microlepidoptera, 1: 524.

[platycolpa]

"♂. 20 mm. . . . French Guiana, St. Jean, R. Maroni, in July; one specimen."

Type: The above indicated male with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5263.

Figure 3, left wings; 3a, lateral aspect of denuded abdomen to show modifications of segments 3 and 4; 3b, ventral view of male genitalia with aedeagus *in situ*.

[sceptrifera]

"♂. 17–19 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, 19 mm., with "LeM. .15" at the bottom of the pin-label. Slide No. 5673.

Figure 4, left wings; 4a, lateral aspect of denuded abdomen to show modifications of segments 3 and 4; 4b, ventral view of male genitalia with aedeagus *in situ*.

Gonioterma sceptrifera (Meyrick), new combination

See Gonioterma seppiana (Cramer) above.

HYALOPSEUSTIS Meyrick

Hyalopseustis Meyrick, 1925, Exotic Microlepidoptera, 3: 157. (*Typus generis: Hyalopseustis vitrea* Meyrick, l.c., 158 [monotypy].)

Hyalopseustis vitrea Meyrick

Plate 110, Figures 1–1a; 2–2a

Hyalopseustis vitrea Meyrick, 1925, Exotic Microlepidoptera, 3: 158.

"♂. 47–53 mm. . . . Peru, Chanchamayo; 4 ex. This very curious and interesting insect, entirely unlike any Stenomid, appears to imitate a form of Danaid butterfly, such as *Ithomia*."

Lectotype: One of the above indicated specimens. Slide No. 4959.

Figure 1, left wings; 1a, venation of right wings; 2, lateral aspect of head to show palpus; 2a, ventral view of male genitalia with aedaegus *in situ*.

HYALOPSEUSTIS



LOXOTOMA

LOXOTOMA Zeller

Loxotoma elegans Zeller

Plate 111, Figures 1–1c

Loxotoma elegans Zeller, 1854, Linnaea Entomologica, 9: 383.
Loxotoma rhodanthes Meyrick, 1915, Exotic Microlepidoptera, 1: 382.

[rhodanthes]

"♀. 34 mm. . . . British Guiana, Bartica, in December (*Parish*); one specimen."

Type: The specimen indicated above dated "12.12". Slide No. 5643.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

LYCHNOCRATES Meyrick

Lychnocrates Meyrick, 1925, Exotic Microlepidoptera, 3: 226. (*Typus generis: Lychnocrates leucocapna* Meyrick, l.c., 227 [monotypy].)

Lychnocrates leucocapna Meyrick

Plate 112, Figures 1–1d

Lychnocrates leucocapna Meyrick, 1925, Exotic Microlepidoptera, 3: 227.

"♂. 46 mm. . . . Colombia, Medina, 1,650 feet; 1 ex."

Type: The above indicated specimen with ".20" in the lower right of the pin-label. Slide No. 4964. I hereby remove *Lychnocrates* from the synonymy of *Timocratica*.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral view of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.



LYCHNOCRATES



MOTHONICA and MYSAROMIMA

MOTHONICA Walsingham

Mothonica fluminata (Meyrick), new combination

Plate 113, Figures 1–1b

Stenoma fluminata Meyrick, 1912, Trans. Ent. Soc. London, 1911: 716.

"♂♀. 18–27 mm. . . . Colombia, San Antonio, 5,800 feet, in November; Dutch Guiana, Paramaribo, in December; five specimens."

Lectotype: ♂, "San Antonio, W. Colombia, 5,800′, .11.07." Slide No. 5415.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

MYSAROMIMA Meyrick

Mysaromima Meyrick, 1926, Exotic Microlepidoptera, 3: 227. (*Typus generis: Mysaromima liquescens* Meyrick, l.c. [monotypy].)

Mysaromima liquescens Meyrick

Plate 113, Figures 2–2a

Mysaromima liquescens Meyrick, 1926, Exotic Microlepidoptera, 3: 227.

"♂. 40 mm. . . . Colombia, Rio Negro, 1,650 feet, April; 1 ex. This striking insect is an interesting example of the type imitative of birds' excrement; it is difficult to believe that the raised white dots and streaks are not fluid."

Type: Lost. For the illustrations of this species I have used a male specimen labelled "Minero, Colombia. 10.20". This, the only specimen I have seen, is without abdomen. Wing slide No. 9303.

Figure 2, left wings; 2a, venation of right wings.

ORPHNOLECHIA Meyrick

Orphnolechia Meyrick, 1909, Trans. Ent. Soc. London, 1909: 28. (*Typus generis: Orphnolechia crypsiphragma* Meyrick, l.c., 29 [by original designation].)

Orphnolechia crypsiphragma Meyrick

Plate 114, Figures 1–1c

Orphnolechia crypsiphragma Meyrick, 1909, Trans. Ent. Soc. London, 1909: 29.

"♂. 13 mm. . . . Bolivia, Songo; one specimen."

Type: The above indicated specimen with "S. .07" at the bottom of the pin-label. Slide No. 4978.

Figure 1, left wings; 1a, venation of right wings; 1b, anterior aspect of head to show palpi; 1c, ventral view of male genitalia with aedeagus *in situ*.



ORPHNOLECHIA



ORPHNOLECHIA

Orphnolechia acridula (Meyrick), new combination

Plate 115, Figures 1–1b

Stenoma acridula Meyrick, 1918, Exotic Microlepidoptera, 2: 201.

"♂ 11 mm., ♀ 13 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: ♂. The three specimens have identical data with "LeM. .16" at the bottom of the pin-label. Slide No. 5158. There is a small, paired, ventral hair-pencil from base of first segment.

Figure 1, right wings (image reversed); 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Orphnolechia anaphanta (Meyrick), new combination

Plate 115, Figures 2–2b

Stenoma anaphanta Meyrick, 1925, Exotic Microlepidoptera, 3: 180.

"♂♀. 11 mm. . . . Brazil, Santarem, R. Trombetas, Teffé, August to December (*Parish*); 4 ex. (2 ♂, 2 ♀). The peculiar modification of the veins of forewing is connected with the development of the equally unique glandular streaks of the male".

Lectotype: ♂, "Santarem, Brazil, Parish. 8.19." Slide No. 5120.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Orphnolechia neastra (Meyrick), new combination

Plate 115, Figures 3–3a

Stenoma neastra Meyrick, 1915, Exotic Microlepidoptera, 1: 412.

"♂. 11 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen. Allied to *chionogramma*. . . ."

Type: The above indicated specimen dated "3.13". Slide No. 5124.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Orphnolechia nuntia (Meyrick), new combination

Plate 115, Figures 4–4c

Stenoma nuntia Meyrick, 1925, Exotic Microlepidoptera, 3: 179.

"♀. 11 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex."

Type: The above indicated specimen dated ".3.20". Slide No. 5119.

Figure 4, left wings; 4a, ventral view of female genitalia (much extraneous matter in ductus bursae); 4b, detail of genital plate and ostium; 4c, signum.

PARASPASTIS Meyrick

Paraspastis Meyrick, 1915, Exotic Microlepidoptera, 1: 479. (*Typus generis: Paraspastis circographa* Meyrick, l.c., 480 [monotypy].)

Paraspastis circographa Meyrick

Plate 116, Figures 1–1e

Paraspastis circographa Meyrick, 1915, Exotic Microlepidoptera, 1: 480.

"♀. 17–18 mm. . . . British Guiana, Mallali, in March (*Parish*); three specimens."

Lectotype: One of the above indicated females all of which bear identical data and are dated "3.13". Slide No. 4968.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, ventral view of female genitalia; 1d, detail of genital plate and ostium; 1e, signum.



PARASPASTIS









PETALOTHYRSA

PETALOTHYRSA Meyrick

Petalothyrsa Meyrick, 1931, Exotic Microlepidoptera, 4: 162. (*Typus generis: Petalothyrsa microphthalma* Meyrick, l.c. [monotypy].)

Petalothyrsa microphthalma Meyrick

Plate 117, Figures 1–1d

Petalothyrsa microphthalma Meyrick, 1931, Exotic Microlepidoptera, 4: 162.

"♂. 8 mm. . . . Brazil, Manaos, November (*Parish*); 1 ex."

Type: The male denoted above dated ".11.19". Slide No. 8042. Although described in the *Epermeniidae* this genus and species clearly belong in the *Stenomidae* and are hereby transferred to that family.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

PETASANTHES Meyrick

Petasanthes Meyrick, 1925, Exotic Microlepidoptera, 3: 158. (*Typus generis: Petasanthes leucactis* Meyrick, l.c. [monotypy].)

Petasanthes leucactis Meyrick

Plate 118, Figures 1–1c

Petasanthes leucactis Meyrick, 1925, Exotic Microlepidoptera, 3: 158.

"♂. 26–29 mm. . . . Ecuador, Huigra, 3,000 feet, February; 2 ex."

Lectotype: ♂, 26 mm., dated "2.13". Slide No. 5116. The lectotype lacks the head, and the second specimen, suffused, has no palpi.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of aedeagus; 1c, ventral view of male genitalia with aedeagus removed.



PETASANTHES









PHELOTROPA

PHELOTROPA Meyrick

Phelotropa Meyrick, 1915, Exotic Microlepidoptera, 1: 411. (*Typus generis: Phelotropa oenodes* Meyrick, l.c. [monotypy].)

Phelotropa oenodes Meyrick

Plate 119, Figures 1–1d

Phelotropa oenodes Meyrick, 1915, Exotic Microlepidoptera, 1: 411.

"♀. 22 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated ".3.13". Slide No. 4967.

Figure 1, left wings; 1a, venation of right wings; 1b, detail of genital plate and ostium; 1c, ventral view of female genitalia; 1d, signum.

PRASOLITHITES Meyrick

Prasolithites Meyrick, 1912, Trans. Ent. Soc. London, 1911: 707. (*Typus generis: Prasolithites virens* Meyrick, l.c. [monotypy].)

Prasolithites virens Meyrick

Plate 120, Figures 1–1d

Prasolithites virens Meyrick, 1912, Trans. Ent. Soc. London, 1911: 707.

"♂. 32 mm. . . . Colombia, at 4,400 feet, in February; one specimen."

Type: The above indicated specimen, labelled "4,400′, W. Colombia. P. .2.08". Slide No. 5482.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.



PRASOLITHITES

PROMENESTA

PROMENESTA Busck

Promenesta autampyx Meyrick

Plate 121, Figures 1–1a

Promenesta autampyx Meyrick, 1925, Exotic Microlepidoptera, 3: 161.

"♂♀. 10–12 mm. . . . Peru, Jurimaguas, March (*Parish*); 2 ex."

Lectotype: The male, dated "3.20". Slide No. 4973.

Figure 1, left wings; 1a, lateral aspect of male genitalia with aedeagus *in situ*.

Promenesta callichlora Meyrick

Plate 121, Figures 2–2b

Promenesta callichlora Meyrick, 1915, Exotic Microlepidoptera, 1: 411.

"♀. 13 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen."

Type: The above indicated specimen dated "1.13". Slide No. 4969.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, genital plate and ostium.

Promenesta chrysampyx Meyrick

Plate 121, Figures 3–3c

Promenesta chrysampyx Meyrick, 1915, Exotic Microlepidoptera, 1: 411.

"♀. 12 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The specimen denoted above dated "3.13". Slide No. 4970.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, genital plate, ostium and posterior section of ductus bursae; 3c, signum.

Promenesta haplodoxa Meyrick

Plate 122, Figure 1

Promenesta haplodoxa Meyrick, 1925, Exotic Microlepidoptera, 3: 162.

"♂. 12 mm. . . . Brazil, Santarem, August (*Parish*); 1 ex."

Type: The above indicated specimen dated "8. 19". Slide No. 4972.

Figure 1, left wings. The genitalia of this species are not illustrated due to the unfortunate loss of the slide.

Promenesta isotrocha Meyrick

Plate 122, Figures 2–2c

Promenesta isotrocha Meyrick, 1918, Exotic Microlepidoptera, 2: 209.

"♀. 16 mm. . . . Argentina, Santiago del Estero; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 4974.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate, ostium and posterior part of ductus bursae; 2c, signum.

Promenesta leucomias Meyrick

Plate 122, Figures 3–3b

Promenesta leucomias Meyrick, 1925, Exotic Microlepidoptera, 3: 162.

"♂♀. 13–14 mm. . . . Brazil, Pará, Obidos, Parintins, July to October (*Parish*); 3 ex."

Lectotype: ♂, "Pará, Brazil, Parish, 7. 19." Slide No. 4971.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.



PROMENESTA



PSEPHOMERES

PSEPHOMERES Meyrick

Psephomeres Meyrick, 1916, Exotic Microlepidoptera, 1 : 505. (*Typus generis: Psephomeres leptogramma* Meyrick, l.c., 506 [monotypy].)

Psephomeres leptogramma Meyrick

Plate 123, Figures 1–1d

Psephomeres leptogramma Meyrick, 1916, Exotic Microlepidoptera, 1 : 506.

"♂ 22 mm., ♀ 30 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, with "LeM. .15" at the bottom of the pin-label. Slide No. 4795. The venational character of veins 11 and 12 of the forewing may be specific since the male genitalia are identical with those of *Antaeotricha*.

Figure 1, left wings; 1a, venation of right wings; 1b, anterior view of head to show palpi; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

PTILOGENES Meyrick

Ptilogenes Meyrick, 1917, Exotic Microlepidoptera, 2: 60. (*Typus generis: Stenoma acronitis* Busck, 1911, Proc. U.S. Nat. Mus., 40: 213 [by original designation].)

Ptilogenes acronitis (Busck)

Plate 124, Figures 1–1c

This genus is a synonym of *Chlamydastis* Meyrick. The species is figured from a specimen in the British Museum. Slide No. 5526.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, ventral view of male genitalia with aedeagus *in situ*.





PTILOGENES


1

1a

1b

1c

1d

RHODANASSA

RHODANASSA Meyrick

Rhodanassa Meyrick, 1915, Exotic Microlepidoptera, 1: 480. (*Typus generis: Rhodanassa callimnestra* Meyrick, ibid. [by original designation].)

Rhodanassa io (Busck)

Plate 125, Figures 1–1d

Stenoma io Busck, 1911, Proc. U.S. Nat. Mus., 40: 213.
Rhodanassa callimnestra Meyrick, 1915, Exotic Microlepidoptera, 1: 480 (new synonymy).

[callimnestra]

"♂. 37 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 4956. Busck's type is a female which probably accounts for the differences which Meyrick thought worthy of distinction. Both are from the same locality.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

SETIOSTOMA Zeller

Setiostoma callidora Meyrick

Plate 126, Figures 1–1b

Setiostoma callidora Meyrick, 1909, Trans. Ent. Soc. London, 1909: 36.

"♂. 11–12 mm. . . . Bolivia, Songo; two specimens."

Lectotype: ♂, 11 mm. Both specimens are labelled "Songo, Bolivia. S. .07". Slide No. 8217.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Setiostoma leuconympha Meyrick

Plate 126, Figures 2–2b

Setiostoma leuconympha Meyrick, 1921, Exotic Microlepidoptera, 2: 477.

"♂. 11 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex. Near *haemotheia*."

Type: The above indicated male dated "1.20". Slide No. 8216.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.



SETIOSTOMA



STENOMA

STENOMA Zeller

Stenoma aconitis Meyrick

Plate 127, Figures 1–1c

Stenoma aconitis Meyrick, 1915, Exotic Microlepidoptera, 1: 414.

"♀. 17 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated "3.13". Slide No. 5341.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Stenoma acratodes Meyrick

Plate 127, Figures 2–2a

Stenoma acratodes Meyrick, 1916, Exotic Microlepidoptera, 1: 536.

"♂. 18–19 mm. . . . French Guiana, St. Jean, R. Maroni; three specimens. Allied to *aphanodesma*."

Lectotype: One of the above indicated males with "LeM. .15" at the bottom of the pin-label. Slide No. 5456.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma adminiculata Meyrick

Plate 127, Figures 3–3a

Stenoma adminiculata Meyrick, 1915, Exotic Microlepidoptera, 1: 472.

"♂♀. 24–26 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: ♂. All specimens bear identical data with "LeM. .15" at the bottom of the pin-label. Slide No. 5228.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma adoratrix Meyrick

Plate 127, Figures 4–4a

Stenoma adoratrix Meyrick, 1925, Exotic Microlepidoptera, 3: 200.

"♂. 25–28 mm. . . . Bolivia, Prov. del Sara, Dept. Sta. Cruz, 1,500 feet, December (*Steinbach*); 1 ex."

Lectotype: ♂, 28 mm., dated December. Meyrick records only "1 ex.", but gives the measurements, "25–28 mm.", indicating he had several specimens. There are three specimens from the above locality, agreeing in size with that described by Meyrick. One specimen is dated "November". Slide No. 5245.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma adulans Meyrick

Plate 128, Figures 1–1b

Stenoma adulans Meyrick, 1925, Exotic Microlepidoptera, 3: 220.

"♂. 19 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex."

Type: The above indicated specimen dated "3.20". Slide No. 5450.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma adytodes Meyrick

Plate 128, Figures 2–2a

Stenoma adytodes Meyrick, 1925, Exotic Microlepidoptera, 3: 196.

"♂. 26–28 mm. . . . Peru, Oconeque, Carabaya, 7,000 feet, February (*Ockenden*); 3 ex."

Lectotype: ♂, 28 mm., dated, "II–1905". Slide No. 5109.

Figure 2, left wings; 2a, lateral aspect of aedeagus (lower figure) and ventral view of male genitalia with aedeagus removed (upper figure).

Stenoma aggerata Meyrick

See Stenoma externella (Walker), p. 311.

Stenoma aggregata Meyrick

Plate 128, Figures 3–3a

Stenoma aggregata Meyrick, 1916, Exotic Microlepidoptera, 1: 527.

"♂. 19 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen indicated above with "LeM. .15" at the bottom of the pin-label. Slide No. 5328.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma algosa Meyrick

Plate 128, Figures 4–4a

Stenoma algosa Meyrick, 1916, Exotic Microlepidoptera, 1: 537.

"♂. 21–23 mm. . . . French Guiana, R. Maroni; four specimens. Allied to *bryophanes*."

Lectotype: ♂, with "LeM. .15" at the bottom of the pin-label. One specimen is a female. Slide No. 5475.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

1

1b

1a



2

2a

3

3a



4

2a

4a

STENOMA


1

1a


1b 1c


2


3b


2a


3


3a

3c


4

4a

4b

Stenoma alligans (Butler)

Plate 129, Figures 1–1c

Cryptolechia alligans Butler, 1877, Cistula Entomologica, 2: 191.
Stenoma obelodes Meyrick, 1915, Exotic Microlepidoptera, 1: 456.

[obelodes]

"♀. 20 mm. . . . British Guiana, R. Demerara; one specimen."

Type: The specimen indicated above with "R. .06" at the bottom of the pin-label. Slide No. 5639.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma alluvialis Meyrick

Plate 129, Figures 2–2a

Stenoma alluvialis Meyrick, 1925, Exotic Microlepidoptera, 3: 204.

"♂. 24–25 mm. . . . Peru, Papitinga, June (*Parish*); French Guiana, R. Maroni; 3 ex."

Lectotype: ♂, "Papitinga, Peru, Parish, 6.20." Slide No. 5246.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma amphiptera Meyrick

Plate 129, Figures 3–3c

Stenoma amphiptera Meyrick, 1913, Trans. Ent. Soc. London, 1913: 185.

"♂. 25 mm. . . . Peru, Chanchamayo; one specimen."

Type: The above indicated specimen with "R. .11" at the bottom of the pin-label. Slide No. 5507.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, distal end of aedeagus to show serrate edge; 3c, ventral view of male genitalia with aedeagus removed.

Stenoma anaxesta Meyrick

Plate 129, Figures 4–4b

Stenoma anaxesta Meyrick, 1915, Exotic Microlepidoptera, 1: 452.

"♂. 23 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen. So extremely similar to *catapsecta* from the same locality that I had unhesitatingly placed them together, but certainly distinct by the absence of dorsal hair-pencil of hind-wings. . . ."

Type: The above indicated specimen dated ".3.13". Slide No. 5313.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma anceps (Butler)

Plate 130, Figures 1–1b

Cryptolechia anceps Butler, 1877, Cistula Entomologica, 2: 163.
Stenoma praeceps Meyrick, 1915, Exotic Microlepidoptera, 1: 453.

"♂. 24 mm. . . . French Guiana, St. Laurent; one specimen."

Type: The specimen indicated above with "R. .11" at the bottom of the pin-label. Slide No. 5637.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma ancillaris Meyrick

Plate 130, Figures 2–2c

Stenoma ancillaris Meyrick, 1916, Exotic Microlepidoptera, 1: 530.

"♀. 16 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5360.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signum.

Stenoma ancylacma Meyrick

Plate 130, Figures 3–3c

Stenoma ancylacma Meyrick, 1925, Exotic Microlepidoptera, 3: 205.

"♀. 24 mm. . . . Peru, Andes, at high elevation; 1 ex."

Type: The specimen referred to above, labelled "Cocapata, Peru, 11,500′, .92." Slide No. 5110.

Figure 3, right wings (image reversed); 3a, detail of genital plate and ostium; 3b, signum; 3c, ventral view of female genitalia.

Stenoma anetodes Meyrick

Plate 130, Figures 4–4b

Stenoma anetodes Meyrick, 1915, Exotic Microlepidoptera, 1: 438.

"♀. 20–21 mm. . . . British Guiana, Bartica, in January (*Parish*); two specimens."

Lectotype: ♀, dated ".1.13". Slide No. 5438. One specimen is missing.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum.


1

1a

1b


2

2a

2b


3

3a

3b

3c

4

4a

4b


2c

STENOMA


1

4c


1a


2

3a

2a


3


3c


3b

4

4a

4b

STENOMA

Stenoma antilyra Meyrick

Plate 131, Figures 1–1a

Stenoma antilyra Meyrick, 1925, Exotic Microlepidoptera, 3: 215.

"♂♀. 15–17 mm. . . . Brazil, Parintins, Manaos, Teffé, October to January; Peru, R. Napo, May (*Parish*); French Guiana, R. Maroni; 6 ex."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5443.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma antitacta Meyrick

Plate 131, Figures 2–2a

Stenoma antitacta Meyrick, 1925, Exotic Microlepidoptera, 3: 219.

"♂. 16 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex."

Type: The specimen indicated above dated "3.20". Slide No. 5449.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma aphrogramma Meyrick

Plate 131, Figures 3–3c

Stenoma aphrogramma Meyrick, 1929, Trans. Ent. Soc. London, 76: 515.

"♀. 27–30 mm. . . . Panama, Taboga I., 200–1,000 feet, September, at light; 3 ex. Near *chlorina* Kearf., but distinguished from this and allied species by whitish hindwings."

Type: The specimen so marked in the British Museum, labelled "Taboga Island, Gulf of Panama, at light, 1,000 ft., 17.9.24. St. George Expedn. C. L. Collenette." Slide No. 5574.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium; 3c, signum.

Stenoma apsorrhoa Meyrick

Plate 131, Figures 4–4c

Stenoma apsorrhoa Meyrick, 1915, Exotic Microlepidoptera, 1: 413.

"♂♀. 13–18 mm. . . . British Guiana, Bartica, in December and January (*Parish*); Peru, Pacaya, in June (*Mounsey*); ten specimens. Near *scitiorella*."

Lectotype: ♀, "Bartica, Brit. Guiana, Parish. .12.12." All specimens in the Meyrick collection are females; three specimens are missing. Slide No. 5347.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, detail of genital plate and ostium; 4c, signum.

Stenoma aptila Meyrick

Plate 132, Figures 1–1b

Stenoma aptila Meyrick, 1915, Exotic Microlepidoptera, 1: 467.

"♂♀. 16–21 mm. . . . British Guiana, Bartica, Mallali, and R. Demerara, from December to March (*Parish*); eight specimens. Excessively like *choleroptila*, but the ♂ is without the dorsal hair-pencil of hindwings; . . ."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. .2.13." Slide No. 5409.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma argicerauna Meyrick

Plate 132, Figures 2–2b

Stenoma argicerauna Meyrick, 1925, Exotic Microlepidoptera, 3: 221.

"♂. 23 mm. . . . Colombia, Cantinero, 1,300 feet; 1 ex. Near *algosa*."

Type: The specimen indicated above with ".20" in the lower right corner of the pin-label. Slide No. 5476.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma argospora Meyrick

Plate 132, Figures 3–3a

Stenoma argospora Meyrick, 1915, Exotic Microlepidoptera, 1: 443.

"♂♀. 15–16 mm. . . . British Guiana, Bartica, from December to February (*Parish*); Dutch Guiana, Onoribo, in March; three specimens."

Lectotype: ♂, 16 mm., "Bartica, Brit. Guiana, Parish. .12.12." Slide No. 5654.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma argotoma Meyrick

Plate 132, Figures 4–4a

Stenoma argotoma Meyrick, 1915, Exotic Microlepidoptera, 1: 412.

"♂. 12–13 mm. . . . British Guiana, Mallali, in March (*Parish*); three specimens."

Lectotype: One of the above indicated males all of which bear identical data and are dated ".3.13". Slide No. 5344.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.



1

1b

1a



2

2b



3

3a

2a

4

4a

STENOMA


1

1a


1b


2


2a

2b


3

3a

3b

4

4a

4b

STENOMA

Stenoma aromatica Meyrick

Plate 133, Figures 1–1b

Stenoma aromatica Meyrick, 1915, Exotic Microlepidoptera, 1: 449.

"♂. 22 mm. . . . Brazil, São Paulo; one specimen."

Type: The specimen indicated above with "R. .06" at the bottom of the pin-label. Slide No. 5463.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed (right harpe broken off).

Stenoma arridens Meyrick

Plate 133, Figures 2–2b

Stenoma arridens Meyrick, 1931, Exotic Microlepidoptera, 4: 36.

"♂. 17 mm. . . . French Guiana, R. Maroni; 1 ex. (Coll. Paravicini). Allied to *ascodes*."

Type: The specimen indicated above. In addition to the locality label (without date) is a small white label bearing the inscription "M806". Slide No. 5575.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma ascodes Meyrick

Plate 133, Figures 3–3b

Stenoma ascodes Meyrick, 1915, Exotic Microlepidoptera, 1: 441.

"♂♀. 17–18 mm. . . . British Guiana, Bartica and Mallali, in January and March (*Parish*); two specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. .1.13." Slide No. 5386.

Figure 3, left wings; 3a, ventral tuft of first segment; 3b, ventral view of male genitalia with aedeagus *in situ*.

Stenoma assignata Meyrick

Plate 133, Figures 4–4b

Stenoma assignata Meyrick, 1918, Exotic Microlepidoptera, 2: 208.

"♂. 18 mm. . . . French Guiana, St. Jean, R. Maroni, July; one specimen. Perhaps allied to *algosa*."

Type: The above indicated specimen dated ".7.15". Slide No. 5474.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma associata Meyrick

Plate 134, Figures 1–1a

Stenoma associata Meyrick, 1925, Exotic Microlepidoptera, 3: 214.

"♂. 16–17 mm. . . . Brazil, Pará, Santarem, August; Peru, Jurimaguas, March (*Parish*); 4 ex."

Lectotype: ♂, "Jurimaguas, Peru, Parish. 3.20." Slide No. 5407.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma astronoma Meyrick

Plate 134, Figures 2–2a

Stenoma astronoma Meyrick, 1909, Trans. Ent. Soc. London, 1909: 34.

"♂. 11 mm. . . . Bolivia, Songo; one specimen."

Type: The above indicated specimen, without abdomen, with "S. .07" at the bottom of the pin-label. Male genitalia figured from a specimen from "Cocapata, Peru. 12,000′ .20". Slide No. 5129.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma atmodes Meyrick

Plate 134, Figures 3–3c

Stenoma atmodes Meyrick, 1915, Exotic Microlepidoptera, 1: 451.

"♀. 18 mm. . . . Peru, Pacaya, in June (*Mounsey*); one specimen."

Type: The above indicated specimen dated ".6.12". Slide No. 5337. The genitalia are damaged by fungus.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Stenoma augescens Meyrick

Plate 134, Figures 4–4c

Stenoma augescens Meyrick, 1925, Exotic Microlepidoptera, 3: 206.

"♀. 39 mm. . . . Peru, Putomayo, R. Ampiyacu; 1 ex."

Type: The above indicated specimen without date. Slide No. 5297.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, detail of genital plate and ostium; 4c, signum.


1
1a
4c
2
2a
3
3a
3b
3c
4
4a
4b



STENOMA

Stenoma avida Meyrick

Plate 135, Figures 1–1b

Stenoma avida Meyrick, 1915, Exotic Microlepidoptera, 1: 435.

"♂♀. 15–16 mm. . . . French Guiana, St. Laurent du Maroni (*LeMoult*); British Guiana, Mallali, in March (*Parish*); three specimens. Allied to *humilis*."

Lectotype: ♀, "Mallali, Brit. Guiana, Parish. .3.13." The St. Laurent specimen and one from Mallali are missing. Slide No. 5628.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium.

Stenoma baliandra Meyrick

Plate 135, Figures 2–2b

Stenoma baliandra Meyrick, 1915, Exotic Microlepidoptera, 1: 442.

"♂♀. 15–17 mm. . . . British Guiana, Bartica and Mallali, from December to March (*Parish*); six specimens."

Lectotype: ♂, "Mallali, Brit. Guiana, Parish. .3.13." Slide No. 5981.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*; 2b, denuded abdomen to show modification of segments 3, 4 and 5.

Stenoma bathrocentra Meyrick

Plate 135, Figures 3–3b

Stenoma bathrocentra Meyrick, 1915, Exotic Microlepidoptera, 1: 462.

"♂. 18 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The specimen indicated above dated ".3.13". Slide No. 5424.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma bathrogramma (Meyrick)

Plate 135, Figures 4–4b

Orphnolechia bathrogramma Meyrick, 1912, Trans. Ent. Soc. London, 1911: 711.

"♀. 12 mm. . . . Venezuela, Ciudad Bolivar, in May; two specimens."

Lectotype: One of the above indicated specimens, both of which bear identical data with "R. .5.04" at the bottom of the pin-label. Slide No. 4979.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum.

Stenoma benigna Meyrick

Plate 136, Figures 1–1c

Stenoma benigna Meyrick, 1916, Exotic Microlepidoptera, 1: 540.

"♀. 24–25 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: One of the above two females, both of which bear the same data with "LeM. .16" at the bottom of the pin-label. Slide No. 5373.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma bicensa Meyrick

Plate 136, Figures 2–2c

Stenoma bicensa Meyrick, 1915, Exotic Microlepidoptera, 1: 464.

"♀. 24 mm. . . . Brazil, São Paulo; one specimen."

Type: The above indicated specimen with "R. .07" at the bottom of the pin-label. Slide No. 5272.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium (black spots in upper part of illustration are extraneous matter).

Stenoma binodis Meyrick

Plate 136, Figures 3–3b

Stenoma binodis Meyrick, 1915, Exotic Microlepidoptera, 1: 472.

"♂. 21 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5222.

Figure 3, left wings; 3a, sketch of lateral aspect of anellus; 3b, ventral view of male genitalia with aedeagus *in situ*.

Stenoma bisignata Meyrick

Plate 136, Figures 4–4b

Stenoma bisignata Meyrick, 1916, Exotic Microlepidoptera, 1: 523.

"♂. 23–24 mm. . . . French Guiana, St. Jean, R. Maroni; two specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 5257.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



STENOMA



STENOMA

Stenoma blandula Meyrick

Plate 137, Figures 1–1b, 4a

Stenoma blandula Meyrick, 1915, Exotic Microlepidoptera, 1: 456.

"♀. 17–18 mm. . . . Venezuela, Ciudad Bolivar, in June; Paraguay; two specimens."

Lectotype: ♀, "Ciudad Bolivar, Venezuela. R. .6.04." The specimen from Paraguay is without abdomen. Slide No. 5368.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 4a, signum.

Stenoma bryocosma Meyrick

Plate 137, Figures 2–2b

Stenoma bryocosma Meyrick, 1916, Exotic Microlepidoptera, 1: 535.

"♂. 35 mm. . . . French Guiana, Godebert, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" on the bottom of the pin-label. Slide No. 5419.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma bryophanes Meyrick

Plate 137, Figures 3–3b, 4b

Stenoma bryophanes Meyrick, 1915, Exotic Microlepidoptera, 1: 477.

"♂. 27–28 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, 28 mm., "R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5477.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed; 4b, lateral aspect of anterior part of denuded abdomen to show modification of segments 2, 3 and 4.

Stenoma bryoxyla Meyrick

Plate 137, Figure 4

Stenoma bryoxyla Meyrick, 1915, Exotic Microlepidoptera, 1: 437.

"♂. 19 mm. . . . Peru, Chanchamayo; one specimen."

Type: The specimen indicated above with "R. .11" at the bottom of the pin-label. The abdomen of the type is missing.

Figure 4, left wings.

Stenoma butyrota Meyrick

Plate 138, Figures 1–1c

Stenoma butyrota Meyrick, 1929, Trans. Ent. Soc. London, 76: 516.

"♂. 29 mm. . . . Colombia, Gorgona I., 200 feet, October, at light; 1 ex."

Type: The specimen so marked in the British Museum, labelled "Gorgona Island, Colombia, at light, 200 ft., 7.10.24. St. George Expedn. C. L. Collenette." Slide No. 5577.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed; 1c, paired, ventral, abdominal tuft from first segment.

Stenoma bythitis Meyrick

Plate 138, Figures 2–2b

Stenoma bythitis Meyrick, 1915, Exotic Microlepidoptera, 1: 443.

"♂♀. 17–18 mm. . . . British Guiana, Bartica, in February (*Parish*); two specimens."

Lectotype: The male, dated ".2.13". Slide No. 5472.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma caesarea Meyrick

Plate 138, Figures 3–3b

Stenoma caesarea Meyrick, 1915, Exotic Microlepidoptera, 1: 450.

"♂♀. 18–25 mm. . . . British Guiana, Bartica, from December to February (*Parish*); sixteen specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. .12.12." Nine specimens are missing. Slide No. 5359.

Figure 3, left wings; 3a, lateral aspect of aedeagus (four cornuti partly obscured in photograph); 3b, ventral view of male genitalia with aedeagus removed.

Stenoma caesia Meyrick

Plate 138, Figures 4–4b

Stenoma caesia Meyrick, 1915, Exotic Microlepidoptera, 1: 449.

"♂♀. 16–18 mm. . . . British Guiana, Mallali, in March (*Parish*); four specimens."

Lectotype: ♂. All bear identical data and are dated ".3.13". Slide No. 5358.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a

1c


1b


2

2a

2b

3

3a


3b

4

4a

4b

STENOMA


1

1a

1b


2

2a

2b


3

3a


4

4a

4b


4c

STENOMA

Stenoma callicoma Meyrick

Plate 139, Figures 1–1b

Stenoma callicoma Meyrick, 1916, Exotic Microlepidoptera, 1: 528.

"♂. 20–22 mm. . . . French Guiana, St. Jean, R. Maroni, in July; four specimens."

Lectotype: One of the above four males all of which bear identical data with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5354.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma camarodes Meyrick

Plate 139, Figures 2–2b

Stenoma camarodes Meyrick, 1915, Exotic Microlepidoptera, 1: 478.

"♂. 21 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen indicated above with "LeM. .15" at the bottom of the pin-label. Slide No. 5469.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma camptospila Meyrick

Plate 139, Figures 3–3a

Stenoma camptospila Meyrick, 1925, Exotic Microlepidoptera, 3: 202.

"♂. 21–22 mm. . . . Brazil, Teffé, December, January (*Parish*); 3 ex."

Lectotype: The male, dated "12.19". Slide No. 5254.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma canonias Meyrick

Plate 139, Figures 4–4c

Stenoma canonias Meyrick, 1913, Trans. Ent. Soc. London, 1913: 187.

"♀. 21 mm. . . . French Guiana, St. Laurient; one specimen."

Type: The above indicated specimen with "R. .11" at the bottom of the pin-label.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, detail of genital plate and ostium; 4c, signum.

Stenoma caryodesma Meyrick

Plate 140, Figures 1–1b

Stenoma caryodesma Meyrick, 1925, Exotic Microlepidoptera, 3: 184.

"♀. 11 mm. . . . Peru, Iquitos, March (*Parish*); 1 ex."

Type: The above indicated specimen dated "3.20". Slide No. 5143.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium.

Stenoma cassigera Meyrick

Plate 140, Figures 2–2b

Stenoma cassigera Meyrick, 1915, Exotic Microlepidoptera, 1: 479.

"♂. 18–23 mm. . . . French Guiana, St. Jean, R. Maroni, in March; three specimens."

Lectotype: One of the above three specimens with "LeM. .15" at the bottom of the pin-label. Slide No. 5484.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma castellana Meyrick

Plate 140, Figures 3–3b

Stenoma castellana Meyrick, 1916, Exotic Microlepidoptera, 1: 516.

"♂. 20–21 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: One of the above indicated specimens, both of which bear identical data with "LeM. .15" at the bottom of the pin-label. Slide No. 5106.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma catapsecta Meyrick

Plate 140, Figures 4–4b

Stenoma catapsecta Meyrick, 1915, Exotic Microlepidoptera, 1: 452.

"♂. 26 mm. . . . British Guiana, Mallali, in March (*Parish*); two specimens."

Lectotype: One of the above indicated specimens, both of which bear identical data, dated "3.13". Slide No. 5317. There is a median, ventral hair-pencil from first and second sternites.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



STENOMA

1

1a

1b



2



3



4a

4b

STENOMA

Stenoma catharmosta Meyrick

Plate 141, Figures 1–1b

Stenoma catharmosta Meyrick, 1915, Exotic Microlepidoptera, 1: 445.

"♂. 16–19 mm. . . . British Guiana, Bartica, in February (*Parish*); Dutch Guiana, Paramaribo, in May; two specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. .2.13." Slide No. 5382.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma cathosiota Meyrick

Plate 141, Figure 2

Stenoma cathosiota Meyrick, 1925, Exotic Microlepidoptera, 3: 200.

"♀. 31 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex."

Type: The above indicated specimen dated "1.20". The abdomen of the type is missing.

Figure 2, left wings.

Stenoma capnobola Meyrick

Plate 141, Figure 3

Stenoma capnobola Meyrick, 1913, Trans. Ent. Soc. London, 1913: 186.

"♀. 26 mm. . . . Dutch Guiana, Paramaribo; one specimen."

Type: The above indicated specimen with "E. .12.92" at the bottom of the pin-label. The abdomen and right forewing are missing.

Figure 3, left wings.

Stenoma centrodina Meyrick

See Stenoma trichocolpa Meyrick, p. 371.

Stenoma charitarcha Meyrick

Plate 141, Figures 4–4b

Stenoma charitarcha Meyrick, 1915, Exotic Microlepidoptera, 1: 438; 1932, 4: 303.

"♂. 16 mm. . . . British Guiana, Bartica, in January and February (*Parish*); two specimens."

Lectotype: ♂, dated ".2.13". Slide No. 5677. The hindwings and abdomen are missing from the second male.

I have compared *charitarcha* with Walsingham's *symmicta* with which Meyrick synonymised it. The aedeagus of *charitarcha* bears two pairs of longitudinal dorsal spines, the anellus possesses paired, long, strongly sclerotised dorso-lateral processes and the harpe extends well beyond the uncus. All of these characters are absent in *symmicta*.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma chionogramma (Meyrick)

Plate 142, Figures 1–1a; 2–2a

Orphnolechia chionogramma Meyrick, 1909, Trans. Ent. Soc. London, 1909: 29.
Stenoma rectificata Meyrick, 1925, Exotic Microlepidoptera, 3: 180 (new synonymy).

[chionogramma]

"♂. 12 mm. . . . Bolivia, Songo; one specimen."

Type: The above indicated specimen with "S. .07" at the bottom of the pin-label. Slide No. 5128.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

[rectificata]

"♂♀. 9–10 mm. . . . Peru, Iquitos, March, 1 ♂ (type); Brazil, Santarem, August, 1 ♀ (*Parish*)."

Type: The male designated by Meyrick, dated "3.20". Slide No. 5121.

Figure 2, left wings; 2a, ventro-lateral view of male genitalia with aedeagus *in situ*.

Stenoma chloroplaca (Meyrick)

Plate 142, Figures 3–3c

Gonioterma chloroplaca Meyrick, 1915, Exotic Microlepidoptera, 1: 383.

"♀. 19 mm. . . . British Guiana, Bartica, in December (*Parish*); one specimen."

Type: The above indicated specimen dated "12.12". Slide No. 5236.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium; 3c, signum.

Stenoma chloroxantha Meyrick

Plate 142, Figures 4–4b

Stenoma chloroxantha Meyrick, 1925, Exotic Microlepidoptera, 3: 216.

"♀. 21 mm. . . . Brazil, Rio Grande do Sul, Santa Cruz; 1 ex."

Type: The specimen indicated above with ".20" in the lower right of the pin-label. The abdomen of the type is missing. I figure the male genitalia of a specimen in the U.S.N.M., "Nova Teutonia, Brazil, 27° 11′ B. 52° 23′ L. viii–1848, Fritz Plaumann." Slide No. 8291.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



1

1a

3c



2

2a



3

3a



3b



4

4a

4b

1

1a

1b


2

2a


2b


3

3a

3b

4

4a

4b

STENOMA

Stenoma chromolitha Meyrick

Plate 143, Figures 1–1b

Stenoma chromolitha Meyrick, 1925, Exotic Microlepidoptera, 3: 203.

"♂. 24 mm. . . . Bolivia, Prov. del Sara, Dept. Sta. Cruz, 1,500 feet; 1 ex."

Type: The above indicated specimen, without date. Slide No. 5267.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma chromotechna Meyrick

Plate 143, Figures 2–2b

Stenoma chromotechna Meyrick, 1925, Exotic Microlepidoptera, 3: 212.

"♂. 17 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex. A singular and elegant form, perhaps allied to *modulata*."

Type: The specimen indicated above dated ".1.20". Slide No. 5399.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma chrysogastra Meyrick

Plate 143, Figures 3–3b

Stenoma chrysogastra Meyrick, 1915, Exotic Microlepidoptera, 1: 476.

"♂. 40 mm. . . . French Guiana, St. Jean, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5490.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma cirrhogramma Meyrick

Plate 143, Figures 4–4b

Stenoma cirrhogramma Meyrick, 1930, Exotic Microlepidoptera, 4: 27.

"♂. 24 mm. . . . Brazil, Bahia, S. Antonio da Barra (*Pujol*); 1 ex. (Coll. Paravicini). Probably allied to *chloromis*."

Type: The specimen indicated above dated "1888". A small white label bears the inscription "M372". Slide No. 5580.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma citroxantha Meyrick

Plate 144, Figures 1–1a

Stenoma citroxantha Meyrick, 1916, Exotic Microlepidoptera, 1: 515.

"♂. 17 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5103.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma clavifera Meyrick

See Stenoma salome Busck, p. 356.

Stenoma clysmographa Meyrick

Plate 144, Figures 2–2a

Stenoma clysmographa Meyrick, 1925, Exotic Microlepidoptera, 3: 197.

"♂. 30 mm."

Type: A male labelled "Pozuzo, E. Peru, .10" answering Meyrick's description and measuring exactly 30 mm. is in the Meyrick collection. I regard this specimen as type. Slide No. 5195. There is a strong, ventral, paired, eversible hair-pencil from first segment.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma codicata Meyrick

Plate 144, Figures 3–3b

Stenoma codicata Meyrick, 1916, Exotic Microlepidoptera, 1: 530.

"♂. 18–20 mm. . . . French Guiana, St. Jean, R. Maroni; five specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 5371.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma colligata Meyrick

Plate 144, Figures 4–4a

Stenoma colligata Meyrick, 1915, Exotic Microlepidoptera, 1: 461.

"♂. 14 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The specimen indicated above dated ".3.13". Slide No. 5427.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

1


1a


2

2a


3

3a

3b

4

4a


1

1a


1b


2

2a


2b


3

3a

3b


4


4a

4b

STENOMA

Stenoma collybista (Meyrick)

Plate 145, Figures 1–1b; 4a

Gonioterma collybista Meyrick, 1915, Exotic Microlepidoptera, 1: 384.

"♀. 21–22 mm. . . . Peru, Contamano, R. Ucuyali, in December (*Mounsey*); three specimens."

Lectotype: One of the above indicated specimens all of which bear identical data and are dated "12.12". Slide No. 5214.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 4a, signum.

Stenoma columbaris Meyrick

Plate 145, Figures 2–2b

Stenoma columbaris Meyrick, 1909, Trans. Ent. Soc. London, 1909: 31.

"♂. 28 mm. . . . Peru, Huancabamba; one specimen."

Type: The above indicated specimen with "R. .08" at the bottom of the pin-label. Slide No. 5320.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma comma Busck

Plate 145, Figures 3–3b

Stenoma comma Busck, 1911, Proc. U.S. Nat. Mus., 40: 218.
Stenoma melanocrypta Meyrick, 1915, Exotic Microlepidoptera, 1: 455.

[melanocrypta]

"♂. 14 mm. . . . British Guiana, R. Essequibo, in June; one specimen."

Type: The above indicated specimen with "D. .6.06" at the bottom of the pin-label. Slide No. 5642.

Figure 3, left wings; 3a, lateral view of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma compsocharis Meyrick

Plate 145, Figures 4, 4b

Stenoma compsocharis Meyrick, 1925, Exotic Microlepidoptera, 3: 218.

"♂. 23 mm. . . . Bolivia, Andes, 12,000 feet; 2 ex. Near *symmicta*."

Lectotype: ♂, "Andes, Bolivia. .20." The second specimen is missing. The left forewing of the lectotype is missing. Slide No. 5436.

Figure 4, right wings (image reversed); 4b, ventral view of male genitalia with aedeagus *in situ*.

Stenoma congrua Meyrick

Plate 146, Figures 1–1a

Stenoma congrua Meyrick, 1925, Exotic Microlepidoptera, 3: 213.

"♂. 15 mm. . . . Peru, Iquitos, May (*Parish*); 2 ex."

Lectotype: One of the above indicated specimens, both of which bear identical data and are dated "5.20". Slide No. 5404.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma contophora Meyrick

Plate 146, Figures 2–2a

Stenoma contophora Meyrick, 1915, Exotic Microlepidoptera, 1: 472.

"♂. 28 mm. . . . French Guiana, Godebert, R. Maroni; two specimens."

Lectotype: One of the above indicated specimens, both of which bear identical data with "LeM. .15" at the bottom of the pin-label. Slide No. 5227.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma contumax Meyrick

See Stenoma thoristes Busck, p. 368.

Stenoma conveniens Meyrick

Plate 146, Figures 3–3c

Stenoma conveniens Meyrick, 1925, Exotic Microlepidoptera, 3: 217.

"♀. 14 mm. . . . Brazil, Matura, June (*Parish*); 1 ex."

Type: The above indicated specimen with "P. .6.20" at the bottom of the pin-label. Slide No. 5441.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, lateral aspect of genital plate and ostium; 3c, signum.

Stenoma crepitans Meyrick

Plate 146, Figures 4–4b

Stenoma crepitans Meyrick, 1918, Exotic Microlepidoptera, 2: 204.

"♂. 17 mm. . . . French Guiana, St. Jean, R. Maroni, July; one specimen."

Type: The above indicated specimen with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5261.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus *in situ*.


1

1a


2

3c

2a

3

3a

3b

4

4a

4b

STENOMA

1

1a

1b



2

2a



2b

3

3a



4

4a

STENOMA

Stenoma crocosticta Meyrick

Plate 147, Figures 1–1b

Stenoma crocosticta Meyrick, 1925, Exotic Microlepidoptera, 3: 219.

"♂. 23 mm. . . . Peru, Iquitos, May (*Parish*); 1 ex."

Type: The specimen indicated above dated "5.20". Slide No. 5447.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed. Note the long, filamentous process from sacculus paralleling ventral edge of harpe.

Stenoma crypsastra Meyrick

Plate 147, Figures 2–2b

Stenoma crypsastra Meyrick, 1915, Exotic Microlepidoptera, 1: 414.

"♂. 10 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen."

Type: The above indicated specimen dated "1.13". Slide No. 5125.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma crypsetaera Meyrick

Plate 147, Figures 3–3a

Stenoma crypsetaera Meyrick, 1925, Exotic Microlepidoptera, 3: 183.

"♂. 13 mm. . . . Brazil, Manaos, November (*Parish*); 1 ex. Perhaps near *scoriodes*."

Type: The above indicated specimen dated "11.19". Slide No. 5429.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma cyanarcha Meyrick

Plate 147, Figures 4–4a

Stenoma cyanarcha Meyrick, 1915, Exotic Microlepidoptera, 1: 414.

"♂. 10 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated "3.13". Slide No. 5123.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*. There is a lateral, spatulate process on each side of the uncus.

Stenoma cycloptila Meyrick

Plate 148, Figures 1–1c

Stenoma cycloptila Meyrick, 1915, Exotic Microlepidoptera, 1: 446.

"♀. 15–17 mm. . . . British Guiana, Bartica, in February (*Parish*); two specimens."

Lectotype: The larger of the two specimens both of which are dated ".2.13". Slide No. 5361.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma cymbalista Meyrick

Plate 148, Figures 2–2c

Stenoma cymbalista Meyrick, 1918, Exotic Microlepidoptera, 2: 204.

"♀. 21 mm. . . . French Guiana, R. Maroni; one specimen. Very similar to *gubernata*. . . ."

Type: The specimen indicated above with "LeM. .15" at the bottom of the pin-label. Slide No. 5440.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Stenoma delenita Meyrick

See Stenoma inflata (Butler), p. 324.

Stenoma delphinodes Meyrick

Plate 148, Figures 3–3b

Stenoma delphinodes Meyrick, 1925, Exotic Microlepidoptera, 3: 222.

"♂. 36–40 mm. . . . Peru, La Oroya, Carabaya, 3,000 feet, March (*Ockenden*); Colombia, R. Tamaua, Choco, 320 feet, February; 3 ex."

Lectotype: ♂, "La Oroya, Carabaya, Peru, 3,000 ft., III–1905. G. Ockenden." Slide No. 5489. The specimen from R. Tamaua, Choco, Colombia, is a female.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed. The uncus is bent down over the gnathos.

Stenoma deltomis Meyrick

Plate 148, Figures 4–4b

Stenoma deltomis Meyrick, 1925, Exotic Microlepidoptera, 3: 206.

"♂. 33 mm. . . . Brazil, Teffé, and Santa Rita, January, February (*Parish*); 2 ex."

Lectotype: ♂, "Santa Rita, Brazil. P. .2.20." Slide No. 5303.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



STENOMA

1

1a

1b

1c


2

2a


2b


3

3a

3b


4

4a


4b

STENOMA

Stenoma desidiosa Meyrick

Plate 149, Figures 1–1c

Stenoma desidiosa Meyrick, 1925, Exotic Microlepidoptera, 3: 201.

"♂♀. 26–27 mm. . . . Colombia, R. Tamaua, Choco, 400 feet, February, and R. Negro, 1,650 feet, April; 3 ex."

Lectotype: ♀, "Choco, R. Tamaua, Colombia, 400 ft., .2.09." Slide No. 5248. The male is missing.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma deuteropa Meyrick

Plate 149, Figures 2–2b; 4a

Stenoma deuteropa Meyrick, 1931, Journ. Linnean Soc. London, 37: 281.

"♀. 16 mm. . . . 1 ex., May. Near *S. notifera*."

Type: The specimen referred to above, labelled "Makthalawaiya, Paraguay, G.S.C. 2.5.27." Slide No. 6243.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 4a, signum.

Stenoma diametrica Meyrick

Plate 149, Figures 3–3b

Stenoma diametrica Meyrick, 1926, Exotic Microlepidoptera, 3: 239.

"♂. 18 mm. . . . Colombia, Mt. Socorro, 12,500 feet, July; 1 ex. Strongly resembles a *Peronea* such as *variegana*, and quite unlike any other *Stenoma*."

Type: The above indicated specimen dated ".7.20". Slide No. 5393.

Figure 3, right wings (image reversed); 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma dilinopa Meyrick

Plate 149, Figures 4, 4b

Stenoma dilinopa Meyrick, 1925, Exotic Microlepidoptera, 3: 198.

"♂♀. 20–24 mm. . . . Brazil, Teffé, January; Peru, R. Napo, May (*Parish*); 4 ex."

Lectotype: ♂, "Teffé, Brazil, Parish. 1.20." Slide No. 5220.

Figure 4, left wings; 4b, ventral view of male genitalia with aedeagus *in situ*.

Stenoma discrepans Meyrick

Plate 150, Figures 1–1c

Stenoma discrepans Meyrick, 1925, Exotic Microlepidoptera, 3: 199.

"♀. 22 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex."

Type: The above indicated specimen dated "3.20". Slide No. 5234.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma dorcadopa Meyrick

Plate 150, Figures 2–2b

Stenoma dorcadopa Meyrick, 1916, Exotic Microlepidoptera, 1: 532.

"♂. 13–14 mm. . . . French Guiana, R. Maroni; four specimens."

Lectotype: ♂, with "LeM. .15" on the bottom of the pin-label. Three of the four original specimens are without abdomens. Slide No. 5384.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma dryaula Meyrick

Plate 150, Figures 3–3c

Stenoma dryaula Meyrick, 1925, Exotic Microlepidoptera, 3: 216.

"♀. 14–15 mm. . . . Brazil, Obidos, August, September (*Parish*); 12 ex., all ♀."

Lectotype: ♀, dated ".8.19". Slide No. 5444.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Stenoma dryoconis Meyrick

Plate 150, Figures 4–4c

Stenoma dryoconis Meyrick, 1930, Exotic Microlepidoptera, 4: 31.

"♀. 34 mm. . . . Bolivia, 1 ex. (Coll. Paravicini). Perhaps nearest *patens*."

Type: The above indicated specimen labelled "Bolivia, Prov. Cochabamba, Y. d. Esperitu Santo, P. Germain." A small white label bears the inscription "M832". Slide No. 5583.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum; 4c, detail of genital plate and ostium.


1
1a
1b
1c
2
2a
2b
3
3a
3b
3c
4
4a
4b
4c



STENOMA

Stenoma dryocosma Meyrick

Plate 151, Figures 1–1c

Stenoma dryocosma Meyrick, 1918, Exotic Microlepidoptera, 2: 206.

"♀. 19–20 mm. . . . French Guiana, R. Maroni; two specimens. Allied to *modulata* and *truncatulata*; it might possibly be the other sex of the former of these species, but the differences are very considerable."

Lectotype: ♀, with "LeM. .16" at the bottom of the pin-label. One specimen is missing. Slide No. 5398.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma dryotechna Meyrick

Plate 151, Figures 2–2c

Stenoma dryotechna Meyrick, 1915, Exotic Microlepidoptera, 1: 421.

"♂♀. 18–19 mm. . . . British Guiana, Bartica, in January and February (*Parish*); Dutch Guiana; four specimens."

Lectotype: ♀, "Bartica, Brit. Guiana, Parish. .2.13." Slide No. 5687. The abdomens of all the males are missing.

This species has been synonymised with *S. ptycta* (Walsingham) but *dryotechna* lacks the black hair-pencil from the upper surface of hindwing at base. The ostium of *ptycta* is nearly twice as wide as that of *dryotechna* and is lined with numerous fine spines. For the above reasons I remove *dryotechna* from synonymy.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Stenoma ebria Meyrick

Plate 151, Figures 3–3c

Stenoma ebria Meyrick, 1915, Exotic Microlepidoptera, 1: 441.

"♀. 16 mm. . . . Venezuela, Ciudad Bolivar, in May; one specimen."

Type: The specimen indicated above with "R. .5.04" at the bottom of the pin-label. Slide No. 5391.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Stenoma elaeurga Meyrick

Plate 151, Figures 4–4a

Stenoma elaeurga Meyrick, 1926, Exotic Microlepidoptera, 3: 226.

"♂. 23 mm. . . . Bolivia, Rio Songo, 2,500 feet; 1 ex. Next *bryocosma*."

Type: The above indicated specimen with ".20" in the lower right corner of the pin-label. Slide No. 5423.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma embythia Meyrick

Plate 152, Figures 1–1b

Stenoma embythia Meyrick, 1916, Exotic Microlepidoptera, 1: 529.

"♂. 13–14 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: The male with "LeM. .16" at the bottom of the pin-label. Slide No. 5362.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma eminens Meyrick

Plate 152, Figures 2–2b

Stenoma eminens Meyrick, 1918, Exotic Microlepidoptera, 2: 206.

"♂. 20 mm. . . . French Guiana, R. Maroni; one specimen. Although very similar to several species of the *sororia* group, it appears to be distinguished from all these by the different form of forewings, of which the apex is not rounded."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5325.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma eminula Meyrick

See Stenoma sericata (Butler), p. 363.

Stenoma emphanes Meyrick

Plate 152, Figures 3–3b

Stenoma emphanes Meyrick, 1917, Exotic Microlepidoptera, 2: 58.

"♀. 20 mm. . . . Burma, Rangoon; one specimen."

Type: The above indicated specimen labelled "Rangoon, Burma. CS. .06." Slide No. 7585. This is not a true Stenomid.

Figure 3, left wings; 3a, ventral view of female genitalia; signum absent; 3b, details of genital plate and ostium.

Stenoma emphatica Meyrick

Plate 152, Figures 4–4a

Stenoma emphatica Meyrick, 1916, Exotic Microlepidoptera, 1: 539.

"♂♀. 22–23 mm. . . . French Guiana, R. Maroni; three specimens. Allied to *melanixa*."

Lectotype: ♂, with "LeM. .15" at the bottom of the pin-label. Slide No. 5217.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.


1

1a

1b

2

2a

2b

3

3a


3b

4

4a

STENOMA



1

3c

1a

2

2a

2b



3



3a

3b

4

4a

4b

STENOMA

Stenoma epicnesta Meyrick

Plate 153, Figures 1–1a

Stenoma epicnesta Meyrick, 1915, Exotic Microlepidoptera, 1: 461.

"♂♀. 14–15 mm. . . . British Guiana, Bartica, in January and February (*Parish*); two specimens. Possibly allied to *avida*."

Lectotype: The male dated ".2.13". Slide No. 5401.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma epipacta Meyrick

Plate 153, Figures 2–2b

Stenoma epipacta Meyrick, 1915, Exotic Microlepidoptera, 1: 444.

"♂. 19 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen. Allied to *adustella* and *melanesia*."

Type: The above indicated specimen dated ".3.13". Slide No. 5377.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma eumenodora Meyrick

Plate 153, Figures 3–3c

Stenoma eumenodora Meyrick, 1937, Exotic Microlepidoptera, 5: 152.

"♂. 15 mm. . . . Argentina, Tigre, February (*Col. F. Bourquin*); 1 ex. Group of *bilinguis*."

Type: The specimen referred to above, erroneously recorded as a male, dated "2.32". Slide No. 5157.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium; 3c, signum.

Stenoma eusticta Meyrick

Plate 153, Figures 4–4b

Stenoma eusticta Meyrick, 1916, Exotic Microlepidoptera, 1: 521.

"♂♀. 18–19 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, with "LeM. .15" at the bottom of the pin-label. Slide No. 5417.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma eva Meyrick

Plate 154, Figures 1–1a

Stenoma eva Meyrick, 1915, Exotic Microlepidoptera, 1: 447.

"♂. 13 mm. . . . British Guiana, Bartica and Mallali, in December and March (*Parish*); four specimens."

Lectotype: ♂, "Mallali, Brit. Guiana, Parish. .3.13." Slide No. 5515.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma exempta Meyrick

Plate 154, Figures 2–2c

Stenoma exempta Meyrick, 1925, Exotic Microlepidoptera, 3: 179.

"♀. 17 mm. . . . Brazil, Obidos, September (*Parish*); 1 ex."

Type: The above indicated specimen dated ".9–19". Slide No. 5342.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signum.

Stenoma exhalata Meyrick

Plate 154, Figures 3–3b

Stenoma exhalata Meyrick, 1915, Exotic Microlepidoptera, 1: 474.

"♂. 21–22 mm. . . . French Guiana, St. Jean, R. Maroni, in July; three specimens."

Lectotype: One of the above indicated males with "LeM. .15" at the bottom of the pin-label. Slide No. 5242.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma expilata Meyrick

Plate 154, Figures 4–4b

Stenoma expilata Meyrick, 1915, Exotic Microlepidoptera, 1: 469.

"♂. 26–31 mm., ♀. 38 mm. . . . British Guiana, R. Demerara; French Guiana, Godebert and Nouveau Chantier; five specimens. Almost indistinguishable in form of wings, colour and markings from *immunda*, but the ♂ has no costal fold."

Lectotype: ♂, "Godebert, R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5306.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

2c


1a

2

2d

2b

3

3a


3b


4


4a

4b

STENOMA



1

1b

1a

2

2a



3



3b

3a



4

3c 1c

Stenoma externella (Walker)

Plate 155, Figures 1–1c

Cryptolechia externella Walker, 1864, List of the specimens of lepidopterous insects in the collection of the British Museum, 29: 732.

Cryptolechia megaspilella Walker, 1864, List of the specimens of lepidopterous insects in the collection of the British Museum, 29: 734.

Stenoma aggerata Meyrick, 1915, Exotic Microlepidoptera, 1: 428.

[aggerata]

"♀. 17 mm. . . . British Guiana, Bartica, in January (*Parish*); one specimen."

Type: The above indicated specimen dated "1.13". Slide No. 5630. Both of Walker's types are males but all names appear to be synonyms.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Stenoma farraria Meyrick

Plate 155, Figures 2–2a

Stenoma farraria Meyrick, 1915, Exotic Microlepidoptera, 1: 462.

"♂. 18 mm. . . . Brazil, São Paulo; one specimen."

Type: The above indicated specimen with "R. .06" in the lower part of the pin-label. Slide No. 5283.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma fastigata Meyrick

Plate 155, Figures 3–3c

Stenoma fastigata Meyrick, 1915, Exotic Microlepidoptera, 1: 430.

"♀. 23–24 mm. . . . British Guiana, Bartica, in February (*Parish*); two specimens."

Lectotype: One of the above indicated females, both of which bear identical data and are dated ".2.13". Slide No. 5473.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium; 3c, signum.

Stenoma finitrix Meyrick

Plate 155, Figure 4

Stenoma finitrix Meyrick, 1925, Exotic Microlepidoptera, 3: 208.

"♂. 18–21 mm. . . . Colombia, San Antonio, 6,600 feet, February; Bolivia, R. Songo, 2,800 feet; 3 ex."

Lectotype: ♂, "S. Antonio, Colombia, 6,600'. .2.20." The abdomens of all three specimens are missing.

Figure 4, left wings.

Stenoma fulcrata Meyrick

Plate 156, Figures 1–1a

Stenoma fulcrata Meyrick, 1915, Exotic Microlepidoptera, 1: 471.

"♂. 23–24 mm. . . . French Guiana, Godebert, R. Maroni; two specimens."

Lectotype: One of the above denoted males with "LeM. .15" at the bottom of the pin-label. Slide No. 5219.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma futura Meyrick

Plate 156, Figures 2–2a

Stenoma futura Meyrick, 1913, Trans. Ent. Soc. London, 1913: 188.

"♂♀. 32–38 mm. . . . Peru, Chanchamayo; three specimens."

Lectotype: ♂. There are three specimens (3 ♂!) under this name in the Meyrick collection agreeing in all details with Meyrick's description, but all are labelled "Huancabamba, Peru. R. .11". There appears to be no doubt that the three specimens are the original types and that Meyrick erred in recording the locality. Slide No. 5293.

Figure 2, right wings (image reversed); 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma gemellata Meyrick

Plate 156, Figures 3–3b

Stenoma gemellata Meyrick, 1916, Exotic Microlepidoptera, 1: 524.

"♂. 17–18 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5258.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma glaucopa Meyrick

Plate 156, Figures 4–4c

Stenoma glaucopa Meyrick, 1912, Trans. Ent. Soc. London, 1911: 712.

"♀. 38 mm. . . . Colombia, San Antonio. 5,800 feet, in November; one specimen."

Type: The female indicated above dated ".11.07". Slide No. 5464.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, detail of genital plate and ostium; 4c, signum.


1

4c

1a


2

2a


3

3a

3b

4

4a


4b

STENOMA

1

1a

1b



2

2a



3



3a

3b

4

4a

4b

STENOMA

Stenoma gymnastis Meyrick

Plate 157, Figures 1–1b

Stenoma gymnastis Meyrick, 1915, Exotic Microlepidoptera, 1: 453.

"♂♀. 21–24 mm. . . . British Guiana, Bartica and Mallali, from December to March (*Parish*); ten specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. 12.12." Slide No. 5314. There is a paired, median-ventral hair-pencil from the first and second sternites.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma halmas Meyrick

Plate 157, Figures 2–2a

Stenoma halmas Meyrick, 1925, Exotic Microlepidoptera, 3: 200.

"♂. 25 mm. . . . Peru, Iquitos, March (*Parish*); 1 ex."

Type: The above indicated specimen dated "3.20". Slide No. 5241.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma haploxyla Meyrick

Stenoma haploxyla Meyrick, 1915, Exotic Microlepidoptera, 1: 420.

"♂. 18 mm. . . . Dutch Guiana, Boschland, in May; one specimen."

Type: The above indicated specimen dated "5.93". The abdomen and forewings are missing and the head is glued to the pin.

Stenoma hectorea Meyrick

Plate 157, Figures 3–3b

Stenoma hectorea Meyrick, 1915, Exotic Microlepidoptera, 1: 384.

"♂. 26–28 mm. . . . Peru, Huancabamba; French Guiana, S. Laurient; two specimens."

Lectotype: ♂, "Huancabamba, Peru. R. .11." The second specimen is missing. Slide No. 5983.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma hemilampra Meyrick

Plate 157, Figures 4–4b

Stenoma hemilampra Meyrick, 1915, Exotic Microlepidoptera, 1: 470.

"♂ 30–34 mm., ♀ 40 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: ♂, "Godebert, R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5218.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma hemiphanta Meyrick

Plate 158, Figures 1–1c

Stenoma hemiphanta Meyrick, 1925, Exotic Microlepidoptera, 3: 198.

"♀. 24–26 mm. . . . Brazil, Teffé, December (*Parish*); 2 ex."

Lectotype: ♀, 26 mm., dated ".12.19". Slide No. 5226.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma herifuga Meyrick

Plate 158, Figures 2–2b

Stenoma herifuga Meyrick, 1932, Exotic Microlepidoptera, 4: 298.

"♀. 11 mm. . . . Peru, Iquitos, March (*Parish*); 1 ex. Allied to group of *communicata*."

Type: The above indicated specimen dated "5.20". Slide No. 5405.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, bursa copulatrix and signum.

Stenoma himerodes Meyrick

Plate 158, Figures 3–3b

Stenoma himerodes Meyrick, 1916, Exotic Microlepidoptera, 1: 535.

"♂♀. 13–15 mm. . . . French Guiana, St. Jean, R. Maroni, in July; three specimens. Allied to *comma*."

Lectotype: ♂, with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5412.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma holcadica Meyrick

Plate 158, Figures 4–4c

Stenoma holcadica Meyrick, 1916, Exotic Microlepidoptera, 1: 531.

"♀. 22–24 mm. . . . French Guiana, R. Maroni; four specimens."

Lectotype: One of the above indicated specimens with "LeM. .15" at the bottom of the pin-label. One specimen is missing.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum; 4c, detail of genital plate and ostium.


1
1a
1b
1c
2
2a
2b
3
3a
3b
4
4a
4b
4c



1

1a

1b



2

2a



2b



3

3a



3b



4



4a



4b

STENOMA

Stenoma holophaea (Meyrick), new combination

Plate 159, Figures 1–1b

Baeonoma holophaea Meyrick, 1916, Exotic Microlepidoptera, 1: 541.

"♂. 18 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5522. This species is not a true *Stenoma*, nor a *Baeonoma*, but I have placed it in this genus because it is closely related to *S. pyrrhonota*, already referred here, and for which there is no described genus which will properly accommodate it.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma hoplitica Meyrick

Plate 159, Figures 2–2b

Stenoma hoplitica Meyrick, 1925, Exotic Microlepidoptera, 3: 216.

"♂. 16–17 mm. . . . Brazil, Teffé, December, January (*Parish*); 2 ex."

Lectotype: The male dated "1.20". Slide No. 5421.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma horocyma Meyrick

Plate 159, Figures 3–3b; 4a

Stenoma horocyma Meyrick, 1925, Exotic Microlepidoptera, 3: 209.

"♂. 10 mm. . . . Brazil, Teffé, December (*Parish*); 1 ex."

Type: The above indicated specimen dated "12.19". Slide No. 5340.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed; 4a, denuded abdomen to show modifications of segments 2 to 5.

Stenoma hospitalis Meyrick

Plate 159, Figures 4, 4b

Stenoma hospitalis Meyrick, 1915, Exotic Microlepidoptera, 1: 463.

"♂. 29 mm. . . . Brazil, Novo Friburgo; one specimen."

Type: The specimen denoted above with "R. .06" at the bottom of the pin-label. Slide No. 5275.

Figure 4, left wings; 4b, ventral view of male genitalia with aedeagus *in situ*.

Stenoma hyacinthitis Meyrick

Plate 160, Figures 1–1b; 4a

Stenoma hyacinthitis Meyrick, 1930, Exotic Microlepidoptera, 3: 559.

"♀. 16 mm. . . . British Guiana, Tumatumari, May (*A. Mackie*); 1 ex. (Brit. Mus.). A distinct species of the *iocoma* group, allied to *icteropis*."

Type: The specimen indicated above dated "29–v–1929". Slide No. 5584.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 4a, signum.

Stenoma hydraena Meyrick

Plate 160, Figures 2–2b

Stenoma hydraena Meyrick, 1916, Exotic Microlepidoptera, 1: 536.

"♂. 16–17 mm. . . . French Guiana, R. Maroni; five specimens."

Lectotype: One of the above five males with "LeM. .15" at the bottom of the pin-label. Slide No. 5471.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma hypocirrha Meyrick

Plate 160, Figures 3–3b

Stenoma hypocirrha Meyrick, 1930, Exotic Microlepidoptera, 4: 30.

"♀. 35 mm. . . . Brazil, S. Paula d'Olivença (*de Mathan*); 1 ex. (Coll. Paravicini). Allied to *leucana*."

Type: The specimen referred to above, labelled "Prov. Amazonas, S. Paulo de Olivença. M. de Mathan." A small white label bears the inscription "M969". Slide No. 5586.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

Stenoma iatma Meyrick

Plate 160, Figures 4, 4b

Stenoma iatma Meyrick, 1915, Exotic Microlepidoptera, 1: 474.

"♂♀. 21–23 mm. . . . French Guiana, St. Jean, R. Maroni, in July; nine specimens."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5243. Eight of the original nine, and three additional specimens, are in the Meyrick collection. All are males.

Figure 4, left wings; 4b, ventral view of male genitalia with aedeagus *in situ*.



STENOMA


1

1b

1a

2

2b

3


3a

2a

4

4a

STENOMA

Stenoma icteropis Meyrick

Plate 161, Figures 1–1b

Stenoma icteropis Meyrick, 1925, Exotic Microlepidoptera, 3: 209.

"♂♀. 13–14 mm. . . . Brazil, Santarem, August (*Parish*); 3 ex."

Lectotype: ♂. The three specimens bear identical data and are dated "8–19". Slide No. 5351. A lateral hair-pencil from metathorax fits into a groove on the abdomen.

Figure 1, right wings (image reversed); 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma immunda (Zeller)

Plate 161, Figures 2–2b

Cryptolechia immunda, ?1854, Linnaea Entomologica, 9: 360.

Cryptolechia tectella Walker, 1864, List of the specimens of lepidopterous insects in the collection of the British Museum, 29: 714.

Stenoma perophora Meyrick, 1915, Exotic Microlepidoptera, 1: 469 (new synonymy).

[perophora]

"♂. 36 mm. . . . Peru; one specimen. Also very close to *immunda*, but distinguished from it and *thylacandra* by more elongate forewings and grey hindwings."

Type: The above indicated specimen with "S,/88" at the bottom of the pin-label. Slide No. 5300.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma impurata Meyrick

Plate 161, Figures 3–3a

Stenoma impurata Meyrick, 1915, Exotic Microlepidoptera, 1: 468.

"♂. 17 mm. . . . Dutch Guiana, Paramaribo, in July; one specimen. Closely allied to *leucana*, in which species also veins 8 and 9 of forewings are stalked, apparently constantly."

Type: The above indicated specimen with "E. .7.92" at the bottom of the pin-label. Slide No. 5278.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma inardescens Meyrick

Plate 161, Figures 4–4a

Stenoma inardescens Meyrick, 1925, Exotic Microlepidoptera, 3: 202.

"♂. 22–23 mm. . . . Brazil, Teffé, January (*Parish*); 2 ex."

Lectotype: One of the above specimens, both of which are dated "1.20". Slide No. 5250.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma inflata (Butler)

Plate 162, Figures 1–1b

Cryptolechia inflata Butler, 1877, Cistula Entomologica, 2: 187.
Stenoma delenita Meyrick, 1915, Exotic Microlepidoptera, 1: 473 (new synonymy).

[delenita]

"♂. 27 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen indicated above with "LeM. .15" at the bottom of the pin-label. Slide No. 5684.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma infusa Meyrick

Plate 162, Figures 2–2a

Stenoma infusa Meyrick, 1916, Exotic Microlepidoptera, 1: 522.

"♂. 22 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5225.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma injucunda Meyrick

Plate 162, Figures 3–3b

Stenoma injucunda Meyrick, 1925, Exotic Microlepidoptera, 3: 219.

"♂♀. 18–19 mm. . . . Peru, R. Napo, Jurimaguas, March, May (*Parish*); 1 ex."

Lectotype: ♂, "Rio Napo, Peru, Parish. 5–20." Meyrick records "1 ex.", an obvious misprint, since both specimens indicated at the beginning of the description are in the Meyrick collection. Slide No. 5448.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma invigilans Meyrick

Plate 162, Figures 4–4b

Stenoma invigilans Meyrick, 1915, Exotic Microlepidoptera, 1: 476.

"♂. 24–26 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, 26 mm., with "LeM. .15" at the bottom of the pin-label. Slide No. 5215.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1


1a

1b

2

2a


3

3a

3b

4

4a

4b

STENOMA


1

1a

1b


2


2b

2a


3


3b

3a

4

3c

STENOMA

Stenoma invulgata Meyrick

Plate 163, Figures 1–1b

Stenoma invulgata Meyrick, 1915, Exotic Microlepidoptera, 1: 458.

"♀. 22–23 mm. . . . Venezuela, Palma Sola; two specimens."

Lectotype: One of the above indicated specimens, both of which bear identical data with "D. .06" at the bottom of the pin-label. Slide No. 5321.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium.

Stenoma iocoma Meyrick

Plate 163, Figures 2–2b

Stenoma iocoma Meyrick, 1915, Exotic Microlepidoptera, 1: 450.

"♂♀. 19–22 mm. . . . British Guiana, Bartica and Mallali, from January to March (*Parish*); French Guiana, St. Laurent; eight specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. .2.13." Slide No. 5350.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma iopercna Meyrick

Plate 163, Figures 3–3c

Stenoma iopercna Meyrick, 1932, Exotic Microlepidoptera, 4: 300.

"♀. 13 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex. Allied to group of *communicata*, but distinguished by its dark, uniform colouring."

Type: The female indicated above dated "3.20". Slide No. 5430.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium; 3c, signum.

Stenoma iostalacta Meyrick

Plate 163, Figure 4

Stenoma iostalacta Meyrick, 1925, Exotic Microlepidoptera, 3: 212.

"♂. 24 mm. . . . Peru, Cocapata, 12,000 feet; 1 ex. Close to *compressa*."

Type: The above indicated specimen, without abdomen, with ".20" in the lower right of the pin-label.

Figure 4, left wings.

Stenoma javarica (Butler)

Plate 164, Figure 1

Cryptolechia javarica Butler, 1877, Cistula Entomologica, 2: 192.

Stenoma paramochla Meyrick, 1918, Exotic Microlepidoptera, 2: 207.

"♀. 19 mm. . . . French Guiana, R. Maroni; one specimen. Allied to *cretifera*."

Type: The above indicated specimen, without abdomen with "LeM. .15" at the bottom of the pin-label.

Figure 1, left wings.

Stenoma jucunda Meyrick

Plate 164, Figures 2–2b; 1b

Stenoma jucunda Meyrick, 1915, Exotic Microlepidoptera, 1: 429.

"♂♀. 18–19 mm. . . . Peru, Contamano, R. Ucayali, in December (*Mounsey*); three specimens."

Lectotype: ♂. All specimens bear identical data and are dated ".12.12". Slide No. 5460. All specimens are male, although one lacks the antennal ciliations, a fact which probably led Meyrick astray. The harpes and uncus of this specimen are clearly visible and the frenulum is single.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed; 1b, lateral aspect of denuded abdomen to show abdominal tufts.

Stenoma lapidea Meyrick

Plate 164, Figures 3–3b

Stenoma lapidea Meyrick, 1916, Exotic Microlepidoptera, 1: 541.

"♂. 38 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen denoted above with "LeM. .16" at the bottom of the pin-label. Slide No. 5653. Meyrick suppressed his name under *S. sesquitertia* (Zeller) from which it is distinguished by the absence of armature on the aedeagus and the harpe is broad with tuberculate surface and inner margin. The harpe of *sesquitertia* is narrow and pointed, with strong setae directed towards sacculus, on inner margin; the aedeagus is armed with two thorn-like spines on distal end ventrally and two distally on dorsal surface. For the above reasons I remove *lapidea* from synonymy.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma leptogma Meyrick

Plate 164, Figures 4–4b; 1a

Stenoma leptogma Meyrick, 1925, Exotic Microlepidoptera, 3: 217.

"♂. 29 mm. . . . Colombia, Monte del Eden, near Haque, 9,600 feet, October; 1 ex."

Type: The specimen indicated above dated ".10.20". The type lacks the right forewing and left hindwing, palpi and most of the legs. Meyrick erroneously recorded the sex as male. Slide No. 5422.

Figure 4, left forewing; 4a, ventral view of female genitalia (damaged); 4b, detail of genital plate and ostium; 1a, the minute signum.

1


1a

1b

2


2a

2b


3

3a

3b

4

4a

4b

STENOMA



STENOMA

Stenoma libertina Meyrick

Plate 165, Figures 1–1b

Stenoma libertina Meyrick, 1916, Exotic Microlepidoptera, 1: 526.

"♂. 26 mm. . . . French Guiana, St. Jean, R. Maroni; one specimen."

Type: The specimen denoted above with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5309.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma licmaea Meyrick

See Stenoma vaga (Butler), p. 375.

Stenoma lithoxesta Meyrick

Plate 165, Figures 2–2b

Stenoma lithoxesta Meyrick, 1915, Exotic Microlepidoptera, 1: 432.

"♂ 22 mm., ♀ 26 mm. . . . British Guiana, Bartica, in February (*Parish*); two specimens. An elegant species suggesting a conglomerate of coloured pebbles."

Lectotype: The male, dated "2.13". Slide No. 5675. A comparison of the types of *lithoxesta* and *nebrita*, with which it has been synonymised, shows them to be closely related but distinct. The uncus of *nebrita* is narrow and broadly dilated distally and the gnathos is wide and lightly sclerotised. The uncus of *lithoxesta* is stout and only moderately dilated distally and the gnathos is pointed and strongly sclerotised. For the above reasons I remove *lithoxesta* from synonymy.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma machinatrix Meyrick

Plate 165, Figure 3

Stenoma machinatrix Meyrick, 1925, Exotic Microlepidoptera, 3: 208.

"♂. 28 mm. . . . Colombia, Minero, 1,650 feet, October; 1 ex."

Type: The above indicated specimen, without abdomen, dated ".10.20".

Figure 3, left wings.

Stenoma melixesta Meyrick

Plate 165, Figure 4

Stenoma melixesta Meyrick, 1925, Exotic Microlepidoptera, 3: 210.

"♀. 42 mm. . . . Colombia, Rio Negro, 1,650 feet, April; 1 ex."

Type: The specimen indicated above, without abdomen, with ".20" in the lower right of the pin-label.

Figure 4, left wings.

Stenoma macroptycha Meyrick

Plate 165, Figures 5–5b

Stenoma macroptycha Meyrick, 1930, Exotic Microlepidoptera, 4: 28.

"♂. 32 mm. . . . Panama, Chiriqui Volcano (*de Mathan*); 1 ex. (Coll. Paravicini). An isolated and peculiar form; perhaps related to *chalybeella* Walk."

Type: The specimen indicated above dated "Mai–Juin 1898". A small white label bears the inscription "M949". Slide No. 5587.

Figure 5, left wings; 5a, lateral aspect of aedeagus; 5b, ventral view of male genitalia with aedeagus removed.

Stenoma melanesia Meyrick

Plate 166, Figures 1–1b

Stenoma melanesia Meyrick, 1912, Trans. Ent. Soc. London, 1911: 712.

"♂. 18 mm. . . . Colombia, at 4,400 feet, in February; one specimen."

Type: The above indicated specimen with "P. .2.08" at the bottom of the pin-label. Slide No. 5378.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma melanixa Meyrick

Plate 166, Figures 2–2a

Stenoma melanixa Meyrick, 1912, Trans. Ent. Soc. London, 1911: 715.

"♂. 26 mm. . . . Colombia, Bogota; one specimen."

Type: The specimen denoted above, without abdomen, with "C. .06" at the bottom of the pin-label. The genitalia are figured from a specimen in the Meyrick collection labelled "Banos, Rio Pastaza, E. Ecuador, 5–7,000 feet. M. G. Palmer." Slide No. 5216.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma melanocrypta Meyrick

See Stenoma comma Busck, p. 291.

Stenoma meligrapta Meyrick

Plate 166, Figures 3–3b

Stenoma meligrapta Meyrick, 1925, Exotic Microlepidoptera, 3: 183.

"♀. 12 mm. . . . Brazil, Parintins (*Parish*); 1 ex."

Type: The above indicated specimen dated ".10–19". Slide No. 5142.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

Stenoma milichodes Meyrick

Plate 166, Figures 4–4c

Stenoma milichodes Meyrick, 1915, Exotic Microlepidoptera, 1: 448.

"♀. 23–24 mm. . . . Colombia, San Antonio, 5,800 feet, in November; two specimens."

Lectotype: ♀, dated ".11.07". Slide No. 5366. Both specimens bear identical data and one, without abdomen, is a male.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, sketch of lateral aspect of genital plate and ostium; 4c, signum.

1

1a

1b


2

4c

2a


3

3a


3b

4

4a

4b



STENOMA

Stenoma mniodora Meyrick

Plate 167, Figures 1–1b

Stenoma mniodora Meyrick, 1925, Exotic Microlepidoptera, 3: 204.

"♂. 22 mm. . . . Colombia, Sosomoko, E. Cordilleras, 2,650 feet, March; 1 ex. Next *byssina*."

Type: The above indicated specimen dated "3.20". Slide No. 5233.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma mundula Meyrick

Plate 167, Figures 2–2b

Stenoma mundula Meyrick, 1916, Exotic Microlepidoptera, 1: 528.

"♂. 16–18 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: One of the above indicated males all of which bear identical data with "LeM. .15" at the bottom of the pin-label. Slide No. 5329.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma myrochroa Meyrick

Plate 167, Figures 3–3c

Stenoma myrochroa Meyrick, 1915, Exotic Microlepidoptera, 1: 448.

"♀. 18 mm. . . . Venezuela, Ciudad Bolivar, in July; one specimen."

Type: The above indicated specimen with "R. .7.04" at the bottom of the pin-label. Slide No. 5364.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Stenoma myrrhinopa Meyrick

Plate 167, Figures 4–4c

Stenoma myrrhinopa Meyrick, 1932, Exotic Microlepidoptera, 4: 300.

"♀, 16–18 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex. Also 1 from Cayuga, Guatemala, in U.S. Nat. Mus."

Type: ♀, "Teffé, Brazil, Parish. .1.20". Slide No. 5338.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signum; 4c, detail of genital plate and ostium.

Stenoma negotiosa Meyrick

Plate 168, Figures 1–1c

Stenoma negotiosa Meyrick, 1925, Exotic Microlepidoptera, 3: 201.

"♂. 30 mm. . . . Brazil, Manaos, November (*Parish*); 1 ex."

Type: The above indicated specimen, erroneously recorded as a male, dated "11.19". Slide No. 5249.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma neoptila Meyrick

Plate 168, Figures 2–2c

Stenoma neoptila Meyrick, 1925, Exotic Microlepidoptera, 3: 195.

"♀. 25 mm. . . . Brazil, R. Trombetas, September (*Parish*); 1 ex."

Type: The above indicated specimen dated "9–19". Slide No. 4994.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signum.

Stenoma neurocentra Meyrick

Plate 168, Figures 3–3a

Stenoma neurocentra Meyrick, 1925, Exotic Microlepidoptera, 3: 179.

"♂. 13 mm. . . . Peru, Rio Napo, May (*Parish*); 1 ex."

Type: The specimen denoted above dated "5.20". Slide No. 4982.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma neurotona (Meyrick)

Plate 168, Figures 4–4c

Athleta neurotona Meyrick, 1915, Exotic Microlepidoptera, 1: 382.

"♀. 14 mm. . . . British Guiana, Bartica, in February (*Parish*); one specimen."

Type: The above indicated specimen dated "2.13". Slide No. 4981.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, detail of genital plate and ostium; 4c, signum.


1

1a

1b

1c

2

2a

2b

3


2c

4c

3a


4

4a


4b

STENOMA

1

1a

1b

1c


2

2a


2b


3

3a


3b


4

5a


5

STENOMA

Stenoma nycteropa Meyrick

Plate 169, Figures 1–1c

Stenoma nycteropa Meyrick, 1915, Exotic Microlepidoptera, 1: 422.

"♀. 14–15 mm. . . . British Guiana, Bartica and Mallali, in December and March (*Parish*); two specimens."

Lectotype: ♀, "Bartica, Brit. Guiana, Parish. 12.12." Slide No. 5139.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma nymphotima Meyrick

Plate 169, Figures 2–2b

Stenoma nymphotima Meyrick, 1931, Exotic Microlepidoptera, 4: 41.

"♂ 17 mm., ♀ 21 mm. . . . Brazil, Petropolis and Cachimbo; 2 ex. (Coll. Paravicini)."

Type: The male so marked in the British Museum labelled "Estado de Bahia, Cachimbo, Pujol. 1890." A small white label bears the inscription "M833". Slide No. 5588.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma obelodes Meyrick

See Stenoma alligans (Butler).

Stenoma ochricollis (Zeller)

Plate 169, Figures 3–3b

Cryptolechia ochricollis Zeller, 1877, Hor. Soc. Ent. Ross., 13: 292, pl. 3, fig. 83.

Stenoma thymiota Meyrick, 1915, Exotic Microlepidoptera, 1: 451.
[thymiota]

"♂♀. 15–16 mm. . . . British Guiana, Bartica and Mallali, from January to March (*Parish*); five specimens."

Lectotype: ♂, "Bartica, Brit. Guiana, Parish. .1.13." One specimen is missing. Slide No. 5339.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma ochrothicta Meyrick

Plate 169, Figure 4

Stenoma ochrothicta Meyrick, 1925, Exotic Microlepidoptera, 3: 180.

"♂. 12 mm. . . . Brazil, Pará, July (*Parish*); 1 ex."

Type: The specimen denoted above, without abdomen, dated "7.19".

Figure 4, left wings.

Stenoma octacentra Meyrick

Plate 169, Figures 5–5a

Stenoma octacentra Meyrick, 1915, Exotic Microlepidoptera, 1: 459.

"♂♀. 17–19 mm. . . . Peru, Pacaya, from June to August (*Mounsey*); three specimens."

Lectotype: ♂, dated ".8.12". Slide No. 5416. One specimen is missing.

Figure 5, left wings; 5a, ventral view of male genitalia with aedeagus *in situ*. There are four cornuti, two on each side, obscured in the illustration.

Stenoma orneopis Meyrick

Plate 170, Figures 1–1b

Stenoma orneopis Meyrick, 1925, Exotic Microlepidoptera, 3: 221.

"♂. 14 mm. . . . Brazil, Teffé, December (*Parish*); 1 ex."

Type: The specimen denoted above dated "12.19". Slide No. 5470.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma orthocapna Meyrick

Plate 170, Figures 2–2b

Stenoma orthocapna Meyrick, 1912, Trans. Ent. Soc. London, 1911: 717.

"♀. 30–31 mm. . . . British Guiana, R. Demerara; two specimens."

Lectotype: ♀, with ".06" at the bottom right of the pin-label. Slide No. 5112.

Figure 2, left wings; 2a, ventral view of female genitalia, except bursa copulatrix; 2b, bursa copulatrix and signum.

Stenoma paracapna Meyrick

Plate 170, Figures 3–3a

Stenoma paracapna Meyrick, 1915, Exotic Microlepidoptera, 1: 444.

"♂. 15 mm. . . . British Guiana, Mallali, in March (*Parish*); three specimens."

Lectotype: ♂, dated ".3.13". Slide No. 5383. One specimen is missing.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma paramochla Meyrick

See Stenoma javarica (Butler), p. 328.

Stenoma paraplecta Meyrick

Plate 170, Figures 4–4a

Stenoma paraplecta Meyrick, 1925, Exotic Microlepidoptera, 3: 214.

"♂. 13 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex. An eccentric species."

Type: The above indicated specimen dated "1.20". Slide No. 5352.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.


1

1a

1b


2


2a

2b


3

3a

4

4a

STENOMA



STENOMA

Stenoma pardalodes Meyrick

Plate 171, Figures 1–1c

Stenoma pardalodes Meyrick, 1918, Exotic Microlepidoptera, 2: 207.

"♀. 16 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen indicated above with "LeM. .15" at the bottom of the pin-label.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma paropta Meyrick

Plate 171, Figures 2–2b

Stenoma paropta Meyrick, 1916, Exotic Microlepidoptera, 1: 536.

"♂♀. 18–19 mm. . . . French Guiana, R. Maroni; in March; two specimens. Allied to *jucunda*."

Lectotype: ♂, "Nouveau Chantier, R. Maroni, Fr. Guiana. LeM. .3.15." Slide No. 5458.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma patens Meyrick

Plate 171, Figures 3–3b

Stenoma patens Meyrick, 1913, Trans. Ent. Soc. London, 1913: 189.

"♂. 34–43 mm. . . . Peru, Huancabamba, Chanchamayo; four specimens. Allied to *immunda*."

Lectotype: ♂, "Huancabamba, Peru, R. .11." Slide No. 5304.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma patula Meyrick

Plate 171, Figures 4–4b

Stenoma patula Meyrick, 1916, Exotic Microlepidoptera, 1: 526.

"♂. 21 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5307.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma periaula Meyrick

Plate 172, Figures 1–1c

Stenoma periaula Meyrick, 1916, Exotic Microlepidoptera, 1: 510.

"♀. 16 mm. . . . French Guiana, St. Jean, R. Maroni, in July; one specimen."

Type: The specimen denoted above with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5346.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Stenoma peridesma Meyrick

Plate 172, Figures 2–2b

Stenoma peridesma Meyrick, 1925, Exotic Microlepidoptera, 3: 220.

"♂. 13 mm. . . . Brazil, Obidos, September (*Parish*); 1 ex."

Type: The specimen indicated above dated ".9–19". Slide No. 5455.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma perirrhoa Meyrick

Plate 172, Figures 3–3a

Stenoma perirrhoa Meyrick, 1930, Exotic Microlepidoptera, 4: 28.

"♀. 26 mm. . . . French Guiana, R. Maroni; 1 ex. (Coll. Paravicini). Nearly related to *adytodes*, but that species has a broader white apical spot, and a dark fuscous spot in the cilia below middle of termen."

Type: The specimen denoted above. A small white label bears the inscription "M362". Slide No. 5590.

Figure 3, left wings; 3a, ventral view of the remains of female genitalia. The abdomen is badly damaged, glued on, and the bursa copulatrix is missing.

Stenoma perophora Meyrick

See Stenoma immunda (Zeller), p. 323.

Stenoma persita Meyrick

Plate 172, Figures 4–4a

Stenoma persita Meyrick, 1915, Exotic Microlepidoptera, 1: 463.

"♂. 19 mm. . . . Peru, Chanchamayo; one specimen."

Type: The above indicated specimen with "R. .11" at the bottom of the pin-label. Slide No. 5269.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ.*



STENOMA


1

1a


1b


2


2a

2b

3


3a


3b 3c


4

4a


4b

STENOMA

Stenoma pertinax Meyrick

Plate 173, Figures 1–1b

Stenoma pertinax Meyrick, 1915, Exotic Microlepidoptera, 1: 459.

"♂. 17 mm. . . . Peru, Pacaya, in July (*Mounsey*); one specimen."

Type: The above indicated specimen dated ".7.12". Slide No. 5370.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma phaeomystis Meyrick

Plate 173, Figures 2–2b; 3c

Stenoma phaeomystis Meyrick, 1925, Exotic Microlepidoptera, 3: 197.

"♂. 27 mm. . . . Peru, R. Ucuyali; 1 ex."

Type: The specimen denoted above with ".13" in the lower right of the pin-label. Slide No. 5197.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed; 3c, left lateral element of anellus.

Stenoma phylloxantha Meyrick

Plate 173, Figures 3–3b

Stenoma phylloxantha Meyrick, 1933, Exotic Microlepidoptera, 4: 435.

"♂. 18 mm. . . . Argentina, Alta Gracia (*C. Bruch*); 1 ex. Perhaps near *cyphoxantha* Meyr., but with peculiar neuration."

Type: The specimen referred to above, erroneously recorded as a male, with "CB. .32" at the bottom of the pin-label. Slide No. 5459.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum.

Stenoma platyphylla Meyrick

Plate 173, Figures 4–4b

Stenoma platyphylla Meyrick, 1916, Exotic Microlepidoptera, 1: 540.

"♂. 17–18 mm. . . . French Guiana, St. Jean, R. Maroni, in July; two specimens."

Lectotype: ♂, with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5514.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma platyterma Meyrick

Plate 174, Figures 1–1c

Stenoma platyterma Meyrick, 1915, Exotic Microlepidoptera, 1: 436.

"♀. 18 mm. . . . British Guiana, Mallali, in March (*Parish*); two specimens. Allied to *truncatula*."

Lectotype: One of the above indicated females, both of which are dated ".3.13". Slide No. 5395.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma porphyrastis Meyrick

Plate 174, Figures 2–2b

Stenoma porphyrastis Meyrick, 1915, Exotic Microlepidoptera, 1: 449.

"♂. 21–22 mm. . . . British Guiana, Mallali, in March (*Parish*); two specimens."

Lectotype: One of the above indicated males dated ".3.13". Slide No. 5355.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma praecauta Meyrick

Plate 174, Figures 3–3b

Stenoma praecauta Meyrick, 1916, Exotic Microlepidoptera, 1: 527; 1925, 3: 192.

"♂. 23–26 mm. . . . French Guiana, St. Jean and Nouveau Chantier, R. Maroni; four specimens. Much like *sororia*, but larger, and readily distinguished by the lighter hindwings."

Lectotype: ♂, "St. Jean, R. Maroni, Fr. Guiana, LeM. .15." Slide No. 5319. Although Meyrick synonymised *praecauta* with *S. nonagriella* (Walker) it is abundantly distinct. The ground colour of Walker's species is whitish-ochreous (badly rubbed) and it probably will be found to be identical with *pantogenes* Meyrick, also from Brazil. *S. praecauta* is very closely related to *S. anceps* (Butler). I hereby remove *praecauta* from synonymy.

There is a median ventral hair-pencil from the first and second sternites.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma praeceps Meyrick

See Stenoma anceps (Butler), p. 260.

Stenoma procritica Meyrick

Plate 174, Figures 4–4b

Stenoma procritica Meyrick, 1925, Exotic Microlepidoptera, 3: 217.

"♂♀. 22–29 mm. . . . Brazil, Teffé, December (*Parish*); S.E. Peru, Santo Domingo, 6,000 feet, November (*Ockenden*); 2 ex."

Lectotype: ♂, "Teffé, Brazil, Parish. 12.19." Slide No. 5420.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a

1b

1c


2

2a


2b

3

3a

3b

4

4a

4b

1

1a

1b


2

2a


2c

2b


3

3a

3b


4


4a

4c

4b

STENOMA

Stenoma psilomorpha Meyrick

Plate 175, Figures 1–1b

Stenoma psilomorpha Meyrick, 1915, Exotic Microlepidoptera, 1: 470.

"♂. 37 mm. . . . Peru; one specimen. Very similar to the preceding species [*expilata*], and also without costal fold, but readily distinguished by the narrower forewings, . . ."

Type: The specimen denoted above with "S./88" at the bottom of the pin-label. Slide No. 5305.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma ptychocentra Meyrick

Plate 175, Figures 2–2c

Stenoma ptychocentra Meyrick, 1916, Exotic Microlepidoptera, 1: 510.

"♀. 13 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 4980.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Stenoma ptychophthalma Meyrick

Plate 175, Figures 3–3b

Stenoma ptychophthalma Meyrick, 1930, Exotic Microlepidoptera, 4: 31.

"♂. 37 mm. . . . Bolivia, R. Songo, 2,500 feet; 1 ex. (Coll. Paravicini). Perhaps allied to *augescens*."

Type: The specimen so marked in the British Museum. A small white label bears the inscription "M959". Slide No. 5592.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma pyrenodes Meyrick

Plate 175, Figures 4–4c

Stenoma pyrenodes Meyrick, 1915, Exotic Microlepidoptera, 1: 448.

"♀. 19 mm. . . . Argentina, Parana; one specimen."

Type: The above indicated specimen with "R. .06" at the bottom of the pin-label. Slide No. 5365.

Figure 4, left wings; 4a, signum; 4b, ventral view of female genitalia; 4c, detail of genital plate and ostium.

Stenoma pyrrhias Meyrick

Plate 176, Figures 1, 1b

Stenoma pyrrhias Meyrick, 1915, Exotic Microlepidoptera, 1: 455.

"♂♀. 18–22 mm. . . . British Guiana, Mallali and Demerara, in March (*Parish*); three specimens."

Lectotype: ♂, "Mallali, Brit. Guiana, Parish. .3.13." Slide No. 5229.

Figure 1, left wings; 1b, lateral aspect of male genitalia with aedeagus *in situ*.

Stenoma pyrrhonota Meyrick

Plate 176, Figures 2–2b

Stenoma pyrrhonota Meyrick, 1915, Exotic Microlepidoptera, 1: 446.

"♂. 13 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The specimen denoted above dated ".3.13". Slide No. 5451.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma recondita Meyrick

Plate 176, Figures 3–3b; 1a

Stenoma recondita Meyrick, 1915, Exotic Microlepidoptera, 1: 439.

"♂♀. 14–17 mm. . . . British Guiana, Bartica, from December to February (*Parish*); three specimens."

Lectotype: ♂, dated "12–12". Slide No. 5411.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed; 1a, lateral aspect of denuded abdomen to show abdominal tufts.

Stenoma rectificata Meyrick

See Stenoma chionogramma (Meyrick), p. 284.

Stenoma redintegrata Meyrick

Plate 176, Figures 4–4b

Stenoma redintegrata Meyrick, 1925, Exotic Microlepidoptera, 3: 218.

"♂. 17 mm. . . . Bolivia, Prov. del Sara, Dept. Santa Cruz, 1,500 feet; 1 ex."

Type: The above indicated specimen without date. Slide No. 5445.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.



STENOMA

1

1a

1b



2

2a



2b



3



3a



3b



4

4a

4b

STENOMA

Stenoma regesta Meyrick

Plate 177, Figures 1–1b

Stenoma regesta Meyrick, 1926, Exotic Microlepidoptera, 3: 225.

"♂♀. 26–29 mm. . . . Peru, Jurimaguas, March (*Parish*); Colombia, Sosomoko, 2,659 feet, March; 4 ex."

Lectotype: ♂, "Jurimaguas, Peru, Parish. 3.20." Slide No. 5508. The Sosomoko specimen is missing.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma relata Meyrick

Plate 177, Figures 2–2b

Stenoma relata Meyrick, 1925, Exotic Microlepidoptera, 3: 207.

"♂♀. 19–21 mm. . . . Brazil, Teffé, January; French Guiana, R. Maroni; Peru, Papitinga (Upper Amazon), January (*Parish*); 5 ex."

Lectotype: ♂, "Teffé, Brazil, Parish. 1.20." Slide No. 5327. One specimen is missing.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma remorsa Meyrick

Plate 177, Figures 3–3b

Stenoma remorsa Meyrick, 1925, Exotic Microlepidoptera, 3: 195.

"♀. 30 mm. . . . Bolivia, R. Suruta, Dept. Santa Cruz, 1,300 feet; 1 ex."

Type: The above denoted specimen without date or collector indicated. Slide No. 4993.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

Stenoma rhodocolpa Meyrick

Plate 177, Figures 4–4b

Stenoma rhodocolpa Meyrick, 1916, Exotic Microlepidoptera, 1: 517.

"♂. 23 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5107.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma rhothiodes Meyrick

Plate 178, Figures 1–1c

Stenoma rhothiodes Meyrick, 1915, Exotic Microlepidoptera, 1: 437.

"♀. 17 mm. . . . Brazil, São Paulo; one specimen."

Type: The specimen indicated above with "R. .07" at the bottom of the pin-label. Slide No. 5433.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Stenoma salome Busck

Plate 178, Figure 2

Stenoma salome Busck, 1911, Proc. U.S. Nat. Mus., 40: 219.

Stenoma clavifera Meyrick, 1912, Trans. Ent. Soc. London, 1911: 715.

[clavifera]

"♀. 23 mm. . . . Brazil, São Paulo; one specimen."

Type: The above indicated specimen, without abdomen, with "R. .06" at the bottom of the pin-label.

Figure 2, left wings.

Stenoma salubris Meyrick

Plate 178, Figures 3–3a

Stenoma salubris Meyrick, 1925, Exotic Microlepidoptera, 3: 208.

"♂. 13–14 mm. . . . Brazil, Parintins, Teffé, October, December (*Parish*); 4 ex."

Lectotype: ♂, "Parintins, Brazil, Parish. 10–19." Slide No. 5330.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma satelles Meyrick

Plate 178, Figures 4–4a

Stenoma satelles Meyrick, 1925, Exotic Microlepidoptera, 3: 202.

"♂. 26 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex."

Type: The above indicated specimen dated "1.20". Slide No. 5255.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

1


1a 1b


2


1c

3


3a

4

4a

STENOMA


1

1a

1b


2

2a

2b

2c

3

3a


3b

3c


4

4a

4b

STENOMA

Stenoma satyropa Meyrick

Plate 179, Figures 1–1b

Stenoma satyropa Meyrick, 1915, Exotic Microlepidoptera, 1: 478.

"♂. 25 mm. . . . French Guiana, Godebert, R. Maroni; one specimen. Allied to *glaucopa*."

Type: The specimen indicated above with "LeM. .15" at the bottom of the pin-label. Slide No. 5462.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma sciaphilina (Zeller)

Plate 179, Figures 2–2c

Cryptolechia sciaphilina Zeller, 1877, Hor. Ent. Soc. Ross., 13: 289, pl. 3, fig. 81.

Stenoma torophragma Meyrick, 1915, Exotic Microlepidoptera, 1: 439.

[torophragma]

"♀. 19–21 mm. . . . British Guiana, Mallali, in March (*Parish*); two specimens."

Lectotype: ♀, 21 mm., dated ".3.13". Slide No. 5331.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Stenoma sciocnesta Meyrick

Plate 179, Figures 3–3c

Stenoma sciocnesta Meyrick, 1925, Exotic Microlepidoptera, 3: 213.

"♂♀. 13–14 mm. . . . Brazil, Teffé, December, January (*Parish*); 3 ex. Close to *epicnesta*. . . ."

Lectotype: ♀, dated "12.19". The abdomen of the male is missing. Slide No. 5402.

Figure 3, left wings; 3a, signum; 3b, ventral view of female genitalia; 3c, detail of genital plate and ostium.

Stenoma sciogama Meyrick

Plate 179, Figures 4–4b

Stenoma sciogama Meyrick, 1930, Exotic Microlepidoptera, 3: 558.

"♂ 18–20 mm., ♀ 22–23 mm. . . . Brazil, Bahia, bred from larvae feeding on leaves of *Anona squamosa* (*Dr. G. Bondar*); 5 ex. (type Brit. Mus.). Allied to *paropta*, which has grey head and palpi."

Type: The specimen designated by Meyrick in the British Museum. Slide No. 5457.

Figure 4, dorsal view of unspread type; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma scoliandra Meyrick

Plate 180, Figures 1–1a

Stenoma scoliandra Meyrick, 1915, Exotic Microlepidoptera, 1: 441.

"♂. 15 mm. . . . British Guiana, Bartica, in February (*Parish*); one specimen. The crooked terminal joint of palpi is quite peculiar, but doubtless only sexual; the species is nearly related to *baliandra*."

Type: The specimen indicated above dated ".2.13". Slide No. 5980.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma scoriodes (Meyrick)

Plate 180, Figures 2–2c

Orphnolechia scoriodes Meyrick, 1915, Exotic Microlepidoptera, 1: 382.

"♀. 13–15 mm. . . . British Guiana, Mallali, in March (*Parish*); three specimens."

Lectotype: One of the females denoted above, all of which bear identical data, and are dated ".3.13". Slide No. 5431. One specimen is missing.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Stenoma scortea Meyrick

Plate 180, Figures 3–3c

Stenoma scortea Meyrick, 1915, Exotic Microlepidoptera, 1: 459.

"♀. 25–26 mm. . . . British Guiana, R. Essequibo and R. Demerara, in June; Brazil, Manaos; seven specimens."

Lectotype: ♀, "Manaos, Brazil. 5.06." Slide No. 5285.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Stenoma secundata Meyrick

Plate 180, Figures 4–4a

Stenoma secundata Meyrick, 1925, Exotic Microlepidoptera, 3: 214.

"♂. 12 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex."

Type: The specimen indicated above dated "3.20". Slide No. 5406.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.


1

2a

1a


2

2b

2c

3

3a

3c


4

3b

4a

STENOMA


1

1a

1b

2a

1c

2

3

2b


3a

4

2c


4a

Stenoma seducta Meyrick

Plate 181, Figures 1–1c

Stenoma seducta Meyrick, 1918, Exotic Microlepidoptera, 2: 206.

"♂♀. 17 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♀, with "LeM. .16" at the bottom of the pin-label. Both specimens are females, the opposite sex having been recorded in error, and both bear identical data. Slide No. 5400.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Stenoma sematopa Meyrick

Plate 181, Figures 2–2c

Stenoma sematopa Meyrick, 1915, Exotic Microlepidoptera, 1: 450.

"♀. 15–16 mm. . . . British Guiana, Mallali, in March (*Parish*); three specimens. Closely allied to the preceding [*iocoma*] and the following [*thymiota*] species, notwithstanding the different neuration."

Lectotype: One of the above indicated females, both of which have identical data, and are dated ".3.13". Slide No. 5349.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signum.

Stenoma sequestra Meyrick

Plate 181, Figures 3–3a

Stenoma sequestra Meyrick, 1918, Exotic Microlepidoptera, 2: 205.

"♂♀. 22 mm. . . . French Guiana, Godebert, R. Maroni; two specimens."

Lectotype: ♂, "R. Maroni, Fr. Guiana. LeM. .16." Slide No. 5271.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma sericata (Butler)

Plate 181, Figures 4–4a

Cryptolechia sericata Butler, 1877, Cistula Entomologica, 2: 163.
Stenoma eminula Meyrick, 1915, Exotic Microlepidoptera, 1: 473.

[eminula]

"♂. 23 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen denoted above with "LeM. .15" at the bottom of the pin-label. Slide No. 5682.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma stephanodes Meyrick

Plate 182, Figures 1–1b

Stenoma stephanodes Meyrick, 1931, Exotic Microlepidoptera, 4: 42.

"♀ 32 mm., ♂ 28 mm. . . . French Guiana, R. Maroni; 2 ex. (Coll. Paravicini). Perhaps allied to *fermentata*. I am satisfied that these two curiously dissimilar specimens are correctly associated as sexes."

Type: ♂. Both specimens are in the British Museum and both are marked "Holotype". I have selected the male as the actual type. Slide No. 5604.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma stolida Meyrick

Plate 182, Figures 2–2c

Stenoma stolida Meyrick, 1911, Ann. Transvaal Mus., 3: 75.

"♀. 25–32 mm. . . . Woodbush Village (*Swiertsra*); in December, two specimens. Type in Coll. Meyrick."

Type: ♀, 25 mm., "Haenertsburg, Transvaal. CIS. 12.12.09." Despite the apparent discrepancy in the localities we must accept the specimen in the Meyrick collection as the actual type. The two localities are very near each other so we can attribute the differences to another case of Meyrick's "dual localities". Slide No. 8524.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Stenoma strophalodes Meyrick

Plate 182, Figures 3–3c

Stenoma strophalodes Meyrick, 1915, Exotic Microlepidoptera, 1: 454.

"♀. 22 mm. . . . Peru, Pacaya, in June (*Mounsey*); one specimen."

Type: The above indicated specimen dated ".6.12".

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Stenoma stupefacta Meyrick

Plate 182, Figures 4–4a

Stenoma stupefacta Meyrick, 1916, Exotic Microlepidoptera, 1: 523.

"♂♀. 23–24 mm. . . . French Guiana, R. Maroni; two specimens."

Lectotype: ♂, with "LeM. .15" on the bottom of the pin-label. Slide No. 5256.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

1

1a

1b



2

2a

2c



3

2b

3a

3b



3c

4

4a


1

1a


2

2a


3


3a

3b


4

4a

STENOMA

Stenoma sustentata Meyrick

Plate 183, Figures 1–1a

Stenoma sustentata Meyrick, 1926, Exotic Microlepidoptera, 3: 225.

"♂. 26 mm. . . . Colombia, La Selve Chaco, 4,000 feet, September; 1 ex."

Type: The specimen denoted above with "P. .9.09" at the bottom of the pin-label. Slide No. 5324.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ.*

Stenoma symphonica Meyrick

Plate 183, Figures 2–2a

Stenoma symphonica Meyrick, 1916, Exotic Microlepidoptera, 1: 522.

"♂. 30–33 mm. . . . French Guiana, R. Maroni; eight specimens."

Lectotype: One of the above examples with "LeM. .15" at the bottom of the pin-label. Slide No. 5244.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ.*

Stenoma symposias (Meyrick)

Plate 183, Figures 3–3b

Gonioterma symposias Meyrick, 1915, Exotic Microlepidoptera, 1: 385.

"♂♀. 22–24 mm. . . . Colombia, San Antonio, 5,800 feet, in November; three specimens."

Lectotype: ♂, dated ".11.07". Slide No. 5200.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma tetrabola Meyrick

Plate 183, Figures 4–4a

Stenoma tetrabola Meyrick, 1913, Trans. Ent. Soc. London, 1913: 189.

"♂. 33 mm. . . . Peru, Huancabamba; one specimen."

Type: The specimen denoted above with "R. .11" at the bottom of the pin-label.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ.*

Stenoma thaleropa Meyrick

Plate 184, Figures 1–1b

Stenoma thaleropa Meyrick, 1916, Exotic Microlepidoptera, 1: 529.

"♂. 16 mm. . . . French Guiana, St. Jean, R. Maroni, in July; two specimens. Apparently allied to *porphyrastis*."

Lectotype: One of the above indicated males with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5256.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma thespia Meyrick

Plate 184, Figures 2–2a

Stenoma thespia Meyrick, 1915, Exotic Microlepidoptera, 1: 474.

"♂♀. 27–28 mm. . . . French Guiana, St. Jean, R. Maroni, in July; five specimens."

Lectotype: ♂, with "LeM. .7.15" at the bottom of the pin-label. Slide No. 5247.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma thoristes Busck

Plate 184, Figures 3–3a

Stenoma thoristes Busck, 1911, Proc. U.S. Nat. Mus., 40: 220.

Stenoma contumax Meyrick, 1916, Exotic Microlepidoptera, 1: 522.

[contumax]

"♂. 30–31 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: ♂, with "LeM. .16" at the bottom of the pin-label. Slide No. 5259.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma thylacandra Meyrick

Plate 184, Figures 4–4b

Stenoma thylacandra Meyrick, 1915, Exotic Microlepidoptera, 1: 468.

"♂. 32 mm. . . . French Guiana, R. Maroni; one specimen. Very close to *immunda*, but differs in form of costal fold, which is longer and less broad. . . ."

Type: The specimen denoted above with "LeM. .15" at the bottom of the pin-label. Slide No. 5301. This species has been synonymised with *S. tectella* (Walker), from which it differs by the elongate patch of stout setae on ventral margin of harpe and the absence of the patch of long, strong setae on the middle of harpe. For the above reasons I remove *thylacandra* from synonymy.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma thymiota Meyrick

See Stenoma ochricollis (Zeller), p. 339.

Stenoma torophragma Meyrick

See Stenoma sciaphilina (Zeller), p. 359.



STENOMA



1

1a



2

2a

3

3a

3b

4

5

STENOMA

Stenoma trichocolpa Meyrick

Plate 185, Figures 1–1a; 2–2a

Stenoma trichocolpa Meyrick, 1915, Exotic Microlepidoptera, 1: 445.
Stenoma centrodina Meyrick, 1916, Exotic Microlepidoptera, 1: 531 (new synonymy).

[trichocolpa]

"♂. 16 mm. . . . French Guiana, La Forestière, in January (*LeMoult*); one specimen."

Type: The above indicated specimen dated ".1.13". Slide No. 5379.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

[centrodina]

"♂♀. 13–15 mm. . . . French Guiana, R. Maroni; five specimens. Belongs to group of *adustella*."

Lectotype: ♂, with "LeM. .16" at the bottom of the pin-label. Slide No. 5376.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma trichorda Meyrick

Plate 185, Figures 3–3b

Stenoma trichorda Meyrick, 1912, Trans. Ent. Soc. London, 1911: 713.

"♂. 35–45 mm. . . . Colombia, San Antonio, 5,800 feet, in November and December; eleven specimens."

Lectotype: ♂, dated ".11.07". Of the entire series only one is a female. Slide No. 5294.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma trirecta Meyrick

Plate 185, Figure 4

Stenoma trirecta Meyrick, 1931, Exotic Microlepidoptera, 4: 38.

"♀. 31 mm. . . . Brazil; 1 ex. (Coll. Paravicini). Perhaps allied to *liniella* Busck."

Type: The specimen indicated above, without abdomen. A small white label bears the inscription "M936".

Figure 4, left wings.

Stenoma subita Meyrick

Plate 185, Figure 5

Stenoma subita Meyrick, 1925, Exotic Microlepidoptera, 3: 224.

"♂. 26 mm. . . . Bolivia, Prov. del Sara, Dept. Santa Cruz, 1,500 feet; 1 ex."

Type: The above indicated specimen, without abdomen, badly covered with mould, and without date or collector's name.

Figure 5, left wings.

Stenoma tyrocrossa Meyrick

Plate 186, Figures 1–1c

Stenoma tyrocrossa Meyrick, 1925, Exotic Microlepidoptera, 3: 183.

"♀. 13–14 mm. . . . Brazil, Santarem, Parintins, August, October (*Parish*); 2 ex. Near *triplectra*."

Lectotype: ♀, "Parintins, Brazil, Parish. 10–19." Slide No. 5145.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Stenoma tyroxesta Meyrick

Plate 186, Figure 2

Stenoma tyroxesta Meyrick, 1925, Exotic Microlepidoptera, 3: 213.

"♂. 14 mm. . . . Brazil, Teffé, January (*Parish*); 1 ex."

Type: The above indicated specimen dated "1.20". Only the basal part of the abdomen remains.

Figure 2, left wings.

Stenoma uncticoma Meyrick

Plate 186, Figures 3–3c

Stenoma uncticoma Meyrick, 1916, Exotic Microlepidoptera, 1: 528.

"♂♀. 13–14 mm. . . . French Guiana, R. Maroni; four specimens. Allied to *iocoma*."

Lectotype: ♂. All specimens bear identical data with "LeM. .15" at the bottom of the pin-label. Slide No. 5348.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed; 3c, anterior section of denuded abdomen to show receptacle, of first and second sternites, for hair-pencil from thorax.

Stenoma uranophanes Meyrick

Plate 186, Figures 4–4b

Stenoma uranophanes Meyrick, 1931, Exotic Microlepidoptera, 4: 41.

"♂. 20 mm. . . . Colombia, Cauca (*de Mathan*); 1 ex. (Coll. Paravicini). Of quite exceptional facies, recalling *Hyalopseustis* superficially, but the palpi and other structures are quite normal for *Stenoma*; perhaps allied to *bryophanes*, which it resembles in form of wing, with some suggestion of similar colouring."

Type: The specimen indicated above dated "fin 1897 1–1898". A small white label bears the inscription "M57". Slide No. 5576.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a

1b


2

1c

3c

3

3a


3b


4

4a

4b

STENOMA



1

1a

1b



2

2a

2b



3

3a



3b

4

4a



4b

STENOMA

Stenoma vacans Meyrick

Plate 187, Figures 1–1b

Stenoma vacans Meyrick, 1916, Exotic Microlepidoptera, 1: 519.

"♂. 38–40 mm. . . . French Guiana, R. Maroni; three specimens."

Lectotype: One of the above indicated males, all of which bear identical data with "LeM. .15" at the bottom of the pin-label. Slide No. 5204.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma vaga (Butler)

Plate 187, Figures 2–2b

Cryptolechia vaga Butler, 1877, Cistula Entomologica, 2: 187.
Stenoma licmaea Meyrick, 1915, Exotic Microlepidoptera, 1: 462.

[licmaea]

"♂♀. 22–23 mm. . . . British Guiana, Bartica and Mallali, in December and March (*Parish*); four specimens."

Lectotype: ♂, "Mallali, Brit. Guiana, Parish. .3.13." Slide No. 5680. Three of the original four specimens are missing. I have compared the genitalia of both types and they are identical.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma vasifera Meyrick

Plate 187, Figures 3–3b

Stenoma vasifera Meyrick, 1925, Exotic Microlepidoptera, 3: 204.

"♂. 27 mm. . . . Colombia, Monte del Eden, near Haque, E. Cordilleras, 9,600 feet, October; 1 ex. Probably allied to *punicea*."

Type: The specimen denoted above dated ".10.20". Slide No. 5198. First sternite with a pair of weak, ventral, eversible hair-pencils.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma ventilatrix Meyrick

Plate 187, Figures 4–4b

Stenoma ventilatrix Meyrick, 1916, Exotic Microlepidoptera, 1: 527.

"♂. 26 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 5318. There is a median-ventral hair-pencil from first and second sternites.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Stenoma vexata Meyrick

Plate 188, Figures 1–1a

Stenoma vexata Meyrick, 1915, Exotic Microlepidoptera, 1: 444.

"♂♀. 17–18 mm. . . . British Guiana, Bartica, in January and February (*Parish*); three specimens."

Lectotype: ♂, dated "2.13". Slide No. 5375.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma vinifera Meyrick

Plate 188, Figures 2–2c

Stenoma vinifera Meyrick, 1916, Exotic Microlepidoptera, 1: 521.

"♀. 20–23 mm. . . . French Guiana, R. Maroni; seven specimens."

Lectotype: One of the above indicated specimens, all of which bear identical data with "LeM. .15" at the bottom of the pin-label.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signum.

Stenoma vitreola Meyrick

Plate 188, Figures 3–3b

Stenoma vitreola Meyrick, 1925, Exotic Microlepidoptera, 3: 209.

"♂. 12 mm. . . . Brazil, Pará, July (*Parish*); 1 ex. Allied to *stylonota*."

Type: The specimen denoted above dated ".7.19". Slide No. 5380.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Stenoma volitans Meyrick

Plate 188, Figures 4–4a

Stenoma volitans Meyrick, 1925, Exotic Microlepidoptera, 3: 220.

"♂. 33 mm. . . . Peru, Yahuas Territory; 1 ex."

Type: The above indicated male without date. Slide No. 5465.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.



1

2c

1a



2

2b

3

2a

3b

4

3a

4a

1

1a

1b


2

2a

2b

3


3a


4

4a


4b

STENOMA

Stenoma xanthobyrsa Meyrick

Plate 189, Figures 1–1b

Stenoma xanthobyrsa Meyrick, 1915, Exotic Microlepidoptera, 1: 447.

"♂. 19 mm. . . . British Guiana, Mallali, in March (*Parish*); one specimen."

Type: The above indicated specimen dated ".3.13". Slide No. 5363.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Stenoma xylinopa Meyrick

Plate 189, Figures 2–2b

Stenoma xylinopa Meyrick, 1925, Exotic Microlepidoptera, 3: 203.

"♂. 28 mm. . . . Bolivia, Prov. del Sara, Dept. Sta. Cruz, 1,500 feet; 1 ex."

Type: The specimen denoted above, without date. Slide No. 5274.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Stenoma zephyritis Meyrick

Plate 189, Figures 3–3a

Stenoma zephyritis Meyrick, 1925, Exotic Microlepidoptera, 3: 218.

"♂. 19 mm. . . . Brazil, Santarem, August (*Parish*); 1 ex."

Type: The specimen denoted above dated "8.19". Slide No. 5446.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Stenoma zobeida Meyrick

Plate 189, Figures 4–4b

Stenoma zobeida Meyrick, 1931, Exotic Microlepidoptera, 4: 33.

"♀. 19 mm. . . . Mexico, Tabasco (*Gugelmann*); 1 ex. (Coll. Paravicini). Perhaps allied to *thaleropa*."

Type: The specimen, indicated above, so marked in the British Museum. A small white label bears the inscription "M824". Slide No. 5605. Signum absent.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, detail of genital plate and ostium.

THIOSCELIS Meyrick

Thioscelis Meyrick, 1909, Trans. Ent. Soc. London, 1909: 29. (*Typus generis: Thioscelis directrix* Meyrick, l.c., 30 [monotypy].)

Thioscelis directrix Meyrick

Plate 190, Figures 1–1c

Thioscelis directrix Meyrick, 1909, Trans. Ent. Soc. London, 1909: 30.

"♂. 62–63 mm. . . . Peru, Pozuzo and El Porvenir; two specimens. A singular insect; the actual length of posterior leg is, femur 3 mm., tibia 14 mm., tarsus 24 mm.; total 41 mm."

Lectotype: The male from "El Porvenir, Peru, R. .08". Slide No. 5117.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, ventral view of male genitalia with aedeagus *in situ*.

THIOSCELIS

THYRSOMNESTIS

THYRSOMNESTIS Meyrick

Thyrsomnestis Meyrick, 1929, Trans. Ent. Soc. London, 76: 514. (*Typus generis: Thyrsomnestis ceramoxantha* Meyrick, ibid. [monotypy].)

Thyrsomnestis ceramoxantha Meyrick

Plate 191, Figures 1–1c

Thyrsomnestis ceramoxantha Meyrick, 1929, Trans. Ent. Soc. London, 76: 514.

"♀. 10 mm. . . . Colombia, Gorgona I., 200 feet, November, at light., 1 ex. The specimen is defective, but the species is easily recognisable."

Type: The above indicated specimen in the British Museum labelled "Gorgona Island, Colombia, at light, 200 ft., 21.11.24. St. George Expedn. C. L. Collenette." A small white label bears the inscription "P984". Slide No. 5615. The palpi, slightly sinuate hindwing and the genitalia strongly suggest that this species should be placed in the *Gelechiidae*, where I believe it belongs. In the absence of more and better material I refrain from making a transfer at this time.

Figure 1, left wings; 1a, lateral aspect of head to show palpus; 1b, detail of genital plate and ostium; 1c, ventral view of female genitalia.

TIMOCRATICA Meyrick

Timocratica Meyrick, 1912, Trans. Ent. Soc. London, 1911: 706. (*Typus generis: Timocratica isographa* Meyrick, l.c., 707 [monotypy].)

Timocratica isographa Meyrick

Plate 192, Figures 1–1d

Timocratica isographa Meyrick, 1912, Trans. Ent. Soc. London, 1911: 707.

"♂. 52 mm. . . . Venezuela, Palma Sola; one specimen. In one forewing 7 and 8 appear to be short-stalked, but on close examination it appears rather that the base of 7 is obsolete and its lower extremity appressed to 8."

Type: The above indicated specimen with "D. .06" at the bottom of the pin-label. Slide No. 4960.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral view of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

Timocratica aphanodesma (Meyrick)

See Timocratica tristrigata (Zeller), p. 392.





TIMOCRATICA







TIMOCRATICA

Timocratica argonias (Meyrick)

Plate 193, Figures 1–1c

Stenoma argonias Meyrick, 1925, Exotic Microlepidoptera, 3: 224.

"♂. 42 mm. . . . Brazil, Fonte Boa, February (*Parish*); 1 ex."

Type: The specimen referred to above, erroneously recorded as a male, dated ".2.20". Slide No. 5497.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Timocratica auxoleuca (Meyrick)

Plate 193, Figures 2–2b

Stenoma auxoleuca Meyrick, 1925, Exotic Microlepidoptera, 3: 223.

"♂♀. 47–53 mm. . . . Brazil, Leopoldina, Espiritu Santo; 2 ex."

Lectotype: ♂, with ".24" in the lower right of the pin-label. Slide No. 5493.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Timocratica cantatrix (Meyrick), new combination

Plate 193, Figures 3–3a

Stenoma cantatrix Meyrick, 1925, Exotic Microlepidoptera, 3: 221.

"♂. 31 mm. . . . Bolivia, Prov. del Sara, Dept. Santa Cruz, 1,500 feet; 1 ex."

Type: The specimen denoted above, labelled "Prov. del Sara, Dep. Sta. Cruz, Bolivia. 450 m., December, J. Steinbach." Slide No. 5468.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Timocratica claudescens Meyrick

Plate 193, Figures 4–4a

Timocratica claudescens Meyrick, 1925, Exotic Microlepidoptera, 3: 177.

"♂. 45–48 mm. . . . Peru, San Gaban, 2,500 feet, April; 2 ex."

Lectotype: ♂, 45 mm. Both specimens bear identical data and are dated "4.13". Slide No. 4963.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

Timocratica crassa Meyrick

Plate 194, Figures 1–1a

Timocratica crassa Meyrick, 1925, Exotic Microlepidoptera, 3: 177.

"♂. 54–56 mm. . . . Brazil, Pará, July (*Parish*); 2 ex."

Lectotype: One of the above indicated specimens, both of which bear identical data and are dated "7.19". Slide No. 4962.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Timocratica effluxa (Meyrick)

Plate 194, Figures 2–2b

Lychnocrates effluxa Meyrick, 1930, Exotic Microlepidoptera, 4: 19.

"♂. 41 mm. . . . Bolivia, R. Songo, 2,500 feet; 1 ex. (Coll. Paravicini)."

Type: The specimen so marked in the British Museum. A small white label bears the inscription "M786". Slide No. 5616.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Timocratica isarga (Meyrick)

Plate 194, Figures 3–3c

Stenoma isarga Meyrick, 1925, Exotic Microlepidoptera, 3: 224.

"♀. 40 mm. . . . Bolivia, Prov. del Sara, Dept. Santa Cruz, 1,500 feet, November; 1 ex."

Type: The above indicated specimen with "J. Steinbach" at the bottom of the pin-label. Slide No. 5498.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium; 3c, signum.

Timocratica leucorectis (Meyrick)

Plate 194, Figures 4–4b

Stenoma leucorectis Meyrick, 1925, Exotic Microlepidoptera, 3: 223.

"♂. 50–60 mm. . . . Brazil, Pará, July (*Parish*); and Leopoldina, Espiritu Santo; 2 ex."

Lectotype: ♂, "Leopoldina, Espiritu Santo, Brazil. .24." Slide No. 5492.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

3c

1a

2

2a

2b

3


3a

3b

4

4a

4b

TIMOCRATICA

1

1a

1b

2

2a


2b

3

3a

3b

4

4a


4b

TIMOCRATICA

Timocratica maturescens (Meyrick)

Plate 195, Figures 1–1b

Stenoma maturescens Meyrick, 1925, Exotic Microlepidoptera, 3: 223.

"♂♀. 42–50 mm. . . . French Guiana, R. Maroni; 10 ex."

Lectotype, ♂, 46 mm., with "LeM. .16" at the bottom of the pin-label. Slide No. 5796.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Timocratica megaleuca (Meyrick)

Plate 195, Figures 2–2b; 4a

Stenoma megaleuca Meyrick, 1912, Trans. Ent. Soc. London, 1911: 711.

"♀. 54 mm. . . . Colombia, Popayan; one specimen. Closely allied to *albella* Zell. In one forewing vein 3 is reduplicated, a curious abnormality."

Type: The above indicated specimen with "L. .06" at the bottom of the pin-label. Slide No. 5496.

Figure 2, left wings; 2a, bursa copulatrix and signum (the black lines on each side of the signum, and the spots above and below, are foreign matter); 2b, detail of genital plate and ostium; 4a, ventral view of female genitalia.

Timocratica philomela (Meyrick)

Plate 195, Figures 3–3b

Stenoma philomela Meyrick, 1925, Exotic Microlepidoptera, 3: 224.

"♂. 22 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex."

Type: The above indicated specimen dated "3.20". Slide No. 5499.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Timocratica pompeiana Meyrick

Plate 195, Figures 4, 4b

Timocratica pompeiana Meyrick, 1925, Exotic Microlepidoptera, 3: 176.

"♂. 65 mm. . . . Peru, La Oroya, Carabaya, 3,000 feet, March (*Ockenden*); 1 ex."

Type: The above indicated specimen dated "III–1905". Slide No. 4961.

Figure 4, left wings; 4b, ventral view of male genitalia with aedeagus *in situ*.

Timocratica sacra (Meyrick)

See Timocratica xanthosoma (Dognin), p. 392.

Timocratica syndicastis (Meyrick), new combination

Plate 196, Figures 1–1b

Stenoma syndicastis Meyrick, 1929, Trans. Ent. Soc. London, 76: 516.

"♂. 23 mm. . . . Colombia, Gorgona I., 200 feet, July, at light; 1 ex."

Type: The specimen so marked in the British Museum, labelled "Gorgona Island, Colombia, at light, 200 ft., 7.7.24. St. George Expedn. C. L. Collenette." A small white label bears the inscription "P830". Slide No. 5593.

Figure 1, left wings; 1 a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Timocratica tristrigata (Zeller)

Plate 196, Figures 2–2a

Cryptolechia tristrigata Zeller, 1854, Linnaea Entomologica, 9: 382, pl. 3, fig. 21.
Stenoma aphanodesma Meyrick, 1915, Exotic Microlepidoptera, 1: 478.

[aphanodesma]

"♂. 34 mm. . . . French Guiana, Godebert, R. Maroni; one specimen."

Type: The specimen denoted above with "LeM. .15" at the bottom of the pin-label. Slide No. 5467.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Timocratica xanthosoma (Dognin)

Plate 196, Figures 3–3c

Stenoma xanthosoma Dognin, 1913, Ann. Ent. Soc. Belgium, 57: 416.
Stenoma sacra Meyrick, 1918, Exotic Microlepidoptera, 2: 209.

[sacra]

"♀. 35 mm. . . . French Guiana, R. Maroni; one specimen."

Type: The specimen indicated above with "LeM. .16" at the bottom of the pin-label. Slide No. 5641.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.









TIMOCRATICA

XYLORYCTIDAE


1

1a

1b


2

2a


2b


3

3a

3b


4

4a

4b

ACRIA Stephens

Acria amphorodes Meyrick

Plate 197, Figures 1–1b

Acria amphorodes Meyrick, 1923, Exotic Microlepidoptera, 2: 612.

"♂♀. 22 mm. . . . Punjab, Murree, 7,500 feet, June (*Dutt*); 2 ex."

Lectotype: ♂. Both specimens are labelled "Murree, Punjab, D. 7,500′, .6.18". Both specimens are males, the opposite sex having been recorded in error. Slide No. 7644.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Acria ceramitis Meyrick

Plate 197, Figures 2–2b

Acria ceramitis Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 636.

"♂♀. 16–19 mm. . . . Palni Hills (*Campbell*); Khasis, in June; seven specimens."

Lectotype: ♀, 16 mm., "Khasi Hills, Assam. D. 1906." Slide No. 7643. The five remaining specimens in the Meyrick collection are females.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium.

Acria eulectra Meyrick

Plate 197, Figures 3–3b

Acria eulectra Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 635.

"♂. 16 mm. . . . Khasis, in December; one specimen."

Type: The above indicated specimen, "Khasi Hills, Assam. 12.1906." Slide No. 7640.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Acria xanthosaris Meyrick

Plate 197, Figures 4–4b

Acria xanthosaris Meyrick, 1908, Journ. Bombay Nat. Hist. Soc. 18: 636.

"♂. 16–18 mm. . . . Khasis, in December; two specimens. Specially distinguished by the yellow pecten of hindwings."

Lectotype: ♂, 16 mm., "Khasi Hills, Assam. 12.1906." Slide No. 7642.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

AEOLANTHES Meyrick

Aeolanthes Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 739. (*Typus generis: Aeolanthes callidora* Meyrick, ibid. [by original designation].)

Aeolanthes callidora Meyrick

Plate 198, Figures 1–1d

Aeolanthes callidora Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 739.

"♂♀. 21–24 mm. . . . Khasi Hills; three specimens (*Swinhoe*)."

Lectotype: ♂, 21 mm. (barely), "Khasi Hills, Assam. .6.1906." Slide No. 7673.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.







AEOLANTHES


1

1a

2

2a


2b

3

3a


3b


4

4a

4b

AEOLANTHES

Aeolanthes ampelurga Meyrick

Plate 199, Figures 1–1a

Aeolanthes ampelurga Meyrick, 1925, Exotic Microlepidoptera, 3: 146.

"♂. 24 mm. . . . Kumaon, Muktesar, 7,000 feet, May (*Fletcher*); 1 ex. Allied to *rhodochrysa*. . . ."

Type: The male denoted above dated ".5.23". Slide No. 7666.

Figure 1, left wings; 1a, caudal view of male genitalia in place in the abdomen, with aedeagus *in situ*.

Aeolanthes brochias Meyrick

Plate 199, Figures 2–2b

Aeolanthes brochias Meyrick, 1938, *in* Caradja and Meyrick, Deuts. Ent. Zeit. Iris, 52: 11.

". . . ♂. 23–24 mm. . . . Yül., 2,800–3,200 m., VII, VIII, 3 ex. Nearly allied to *meniscias* and *diacrita*. . . ."

Type: The single male in the Meyrick collection, "Likiang, China. H. .8.34." Slide No. 7681.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Aeolanthes cianolitha Meyrick

Plate 199, Figures 3–3b

Aeolanthes cianolitha Meyrick, 1938, *in* Caradja and Meyrick, Deuts. Ent. Zeit. Iris, 52: 12.

". . . ♂. 21–24 mm. . . . Yül., 3,200 to 4,000 m., VII–IX, 3 ex. Nearest allied to *erebomicta* Meyr."

Lectotype: ♂, "Likiang, China. H. .9.34." Slide No. 7682. Meyrick intended this name to be *cyanolitha* as indicated in his personal catalogues.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Aeolanthes clinacta Meyrick

Plate 199, Figures 4–4b

Aedanthes [recte *Aeolanthes*] *clinacta* Meyrick, 1925, *in* Caradja, Mem. Sect. sti. Acad. Română, 3: 382.

"♂. 16–18 mm. . . . Lienping, April to July; three examples. Allied to *lychnidias*. . . ."

Lectotype: ♂, "Lienping, China. .4.25." Slide No. 7693.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Aeolanthes coronifera Meyrick

Plate 200, Figures 1–1b

Aeolanthes coronifera Meyrick, 1938, *in* Caradja and Meyrick, Deuts. Ent. Zeit. Iris, 52: 12.

". . . ♂. 18–19 mm. . . . Yül., 3,200–4,000 m., VIII, IX, 3 ex."

Lectotype: ♂, "Likiang, China. H. .8.34." Slide No. 7680.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Aeolanthes cyclantha Meyrick

Plate 200, Figures 2–2d

Aeolanthes cyclantha Meyrick, 1923, Exotic Microlepidoptera, 2: 611.

"♀. 22 mm. . . . S. India, Nilgiris, October (*Rao*); 1 ex. Distinct from *rhodochrysa* by dark fuscous hindwings. . . ."

Type: The specimen referred to above, erroneously recorded as a female, dated ".10.17". Slide No. 7665.

Figure 2, right wings (image reversed); 2a, lateral aspect of aedeagus; 2b, enlarged view of cucullus to show short, stout setae; 2c, ventral view of male genitalia with aedeagus removed; 2d, modification of seventh and eighth segments.

Aeolanthes deltogramma Meyrick

Plate 200, Figures 3–3b

Aeolanthes deltogramma Meyrick, 1923, Exotic Microlepidoptera, 2: 611.

"♂. 25 mm. . . . Assam, Shillong, 5,000 feet, July (*Fletcher*); 1 ex."

Type: The male denoted above, "Shillong, Assam. TBF. .22." Slide No. 7664.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Aeolanthes diacritica Meyrick

Plate 200, Figures 4–4b

Aeolanthes diacritica Meyrick, 1918, Exotic Microlepidoptera, 2: 196.

"♂. 19 mm. . . . Assam, Shillong, 5,000 feet, September (*Fletcher*); one specimen."

Type: The above indicated male dated ".9.17". Slide No. 7663.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a

1b


2

2a

2b

2c

2d


3

3a


3b


4

4a

4b

AEOLANTHES













AEOLANTHES

Aeolanthes dicraea Meyrick

Plate 201, Figures 1–1c

Aeolanthes dicraea Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 637.

"♂. 20–22 mm. . . . Khasis, in November and April; two specimens."

Lectotype: The specimen labelled "Khasi Hills, Assam. .4.1906." Both specimens are females, the opposite sex having been recorded erroneously. Slide No. 7668.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Aeolanthes erebomicta Meyrick

Plate 201, Figures 2–2b

Aeolanthes erebomicta Meyrick, 1931, *in* Caradja, Bull. Sec. sci. Acad. roum., 14: 73.

"♂♀. 18–27 mm. . . . 4 ex. July, August."

Lectotype: ♂, 18 mm., "Kwanhsien, China. F. .7.30." Slide No. 7670.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Aeolanthes erythrantis Meyrick

Plate 201, Figures 3–3b

Aeolanthes erythrantis Meyrick, 1935, *in* Caradja and Meyrick, Materialien zu einer Microlepidopteren Fauna der Chinesischen Provinzen Kiangsu, Chekiang und Hunan, 83.

". . . ♂♀. 21 mm. . . . Nearest *rhodochrysa*. 12 ex., Lungtan V, VI."

Lectotype: ♂, "Nanking, China, H. 31.5.33." Slide No. 7677.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Aeolanthes euryatma Meyrick

Plate 201, Figures 4–4b

Aeolanthes euryatma Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 638.

"♂. 17 mm. . . . Khasis, in July; one specimen."

Type: The male indicated above, labelled "Khasi Hills, Assam. .7.1906." Slide No. 7661.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Aeolanthes haematopa Meyrick

Plate 202, Figures 1–1b

Aeolanthes haematopa Meyrick, 1931, *in* Caradja, Bull. Sect. sci. Acad. roum., 14: 73.

"♂♀. 15 mm. . . . 3 ex., July. Nearest *rhodochrysa*."

Lectotype: ♂, "Kwanhsien, China. F. .7.30." Slide No. 7671. The three specimens bear identical data.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed (left cucullus broken off).

Aeolanthes lychnidias Meyrick

Plate 202, Figures 2–2b

Aeolanthes lychnidias Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 638.

"♂. 22 mm. . . . Palni Hills, in November (*Campbell*); one specimen."

Type: The male denoted above, labelled "Palni Hills, S. India. Campbell. .3.06." The discrepancy in date seems to be a printing error since all other details agree with the record. Slide No. 7667.

Figure 2, right wings (image reversed); 2a, lateral aspect of aedeagus; 2b, lateral view of male genitalia with aedeagus removed.

Aeolanthes megalophthalma Meyrick

Plate 202, Figures 3–3b

Aeolanthes megalophthalma Meyrick, 1930, Exotic Microlepidoptera, 4: 11.

"♂♀. 24–30 mm. . . . China, Kwanhsien, July; 5 ex. Allied to *callidora*."

Lectotype: ♂, 24 mm. All specimens bear identical data and are dated ".7.28". Slide No. 7674.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Aeolanthes meniscias Meyrick

Plate 202, Figures 4–4d

Aeolanthes meniscias Meyrick, 1918, Exotic Microlepidoptera, 2: 196.

"♀. 25 mm. . . . Burma, Koni, October (*Manders*); one specimen."

Type: The female denoted above labelled "Koni, Burma. NM. /10/88." Slide No. 7662.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signa; 4c, ostium; 4d, detail of genital plate.



AEOLANTHES



AEOLANTHES

Aeolanthes rhodochrysa Meyrick

Plate 203, Figures 1–1b

Aeolanthes rhodochrysa Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 739.

"♂. 18–19 mm. . . . Khasi Hills; two specimens."

Lectotype: ♂, 18 mm., "Khasi Hills, Assam. D. .1906." Slide No. 7672. There are ten specimens in the Meyrick collection, all dated 1906, from the same locality. Two or three might be excluded from consideration as possible types on the basis of size, but the selection of a lectotype from the remainder is purely arbitrary.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Aeolanthes sagulata Meyrick

Plate 203, Figures 2–2a

Aeolanthes sagulata Meyrick, 1917, Exotic Microlepidoptera, 2: 53.

"♀. 19 mm. . . . Bengal, Chapra (*Mackenzie*); one specimen."

Type: The female indicated above, labelled "Chapra, Bengal. M. .16." Slide No. 7675.

Figure 2, left wings; 2a, ventral view of female genitalia.

Aeolanthes semiostrina Meyrick

Plate 203, Figure 3

Aeolanthes semiostrina Meyrick, 1935, *in* Caradja and Meyrick, Materialien zu einer Microlepidopteren Fauna der Chinesischen Provinzen Kiangsu, Chekiang und Hunan, 82.

". . . ♀. 19 mm. . . . 4 ex., Tienmushan, VI, VII . . . allied to *rhodochrysa*."

Type: The above indicated specimen in the Meyrick collection. The abdomen of the type is missing.

Figure 3, left wings.

Aeolanthes siphonias Meyrick

Plate 203, Figures 4–4b

Aeolanthes siphonias Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 637.

"♂♀. 22–29 mm. . . . Simla, in July, nine specimens."

Lectotype: ♂, 23 mm. All specimens bear identical data and are labelled "Simla, India. R. .7.89." Slide No. 7669. One specimen is missing.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

AGRIOPHARA Rosenstock

Agriophara halarcta Meyrick

Plate 204, Figures 1–1b

Agriophara halarcta Meyrick, 1917, Exotic Microlepidoptera, 2: 59.

"♂. 26 mm. . . . Madagascar, Antananarivo; one specimen."

Type: The above indicated male, "Antananarivo, Madagascar. LeM. .15." Slide No. 8525.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Agriophara parilis Meyrick

Plate 204, Figures 2–2b

Agriophara parilis Meyrick, 1918, Exotic Microlepidoptera, 2: 197.

"♂. 18 mm. . . . New Guinea, Setekwa River, 2–3,000 feet; one specimen. Very like *neoxanta*. . . ."

Type: The above indicated male, "Setekwa R. New Guinea. 2–3,000′. .17." Slide No. 9214.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.







AGRIOPHARA







AMORBAEA

AMORBAEA Meyrick

Amorbaea Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 627. (*Typus generis: Amorbaea hepatica* Meyrick, ibid. [by original designation].)

Amorbaea hepatica Meyrick

Plate 205, Figures 1–1f

Amorbaea hepatica Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 627.

"♂♀. 22–26 mm. . . . Bombay (*Young*); two specimens."

Lectotype: ♀, 22 mm., "Konkan, Bombay. LCY. .05." Slide No. 7700. Both specimens are females with identical data.

Amorbaea, suppressed as a synonym of *Ptochoryctis* by Meyrick, is distinguished from the latter genus by the stalking of veins 3 and 4 and the strong preservation of 1c of the forewing, and the presence of a cross vein between the cell and vein 8 of the hindwing. I hereby remove *Amorbaea* from synonymy.

Figure 1, left wings; 1a, venation of right wings; 1b, ventral view of female genitalia; 1c, dorsal aspect of part of denuded abdomen to show spines; 1d, lateral aspect of head to show palpus; 1e, signum; 1f, detail of genital plate and ostium.

AMPHITRIAS Meyrick

Amphitrias Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 631. (*Typus generis: Amphitrias cynica* Meyrick, ibid. [monotypy].)

Amphitrias cynica Meyrick

Plate 206, Figures 1–1d

Amphitrias cynica Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 631.

"♂♀. 17–21 mm. . . . Patipola and Maskeliya, Ceylon, from February to May (*Alston, Pole*); six specimens."

Lectotype: ♂, "Patipola, Ceylon. GCA. .4.06." Slide No. 7636.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.





AMPHITRIAS







ANTITHYRA

ANTITHYRA Meyrick

Antithyra Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 404. (*Typus generis: Antithyra vineata* Meyrick, ibid. [monotypy].)

Antithyra vineata Meyrick

Plate 207, Figures 1–1d

Antithyra vineata Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 404.

"♀. 11–12 mm. . . . Five specimens, Peradeniya, Ceylon, in February and March (*Green*)."

Lectotype: ♀, 11 mm., dated ".3.05". Slide No. 7637.

Figure 1, left wings; 1a, venation of right wings; 1b, frontal aspect of head to show palpus; 1c, ventral view of female genitalia; 1d, detail of genital plate and ostium.

ANTOLAEA Meyrick

Antolaea Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 779. (*Typus generis: Antolaea xanthopa* Meyrick, l.c., 780 [monotypy].)

Antolaea xanthopa Meyrick

Plate 208, Figures 1–1d

Antolaea xanthopa Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 780.

"♂♀. 13–14 mm. . . . Khasis, in April; two specimens."

Lectotype: ♂, "Khasi Hills, Assam. .4.1907." Slide No. 7639.

Figure 1, left wings; 1a, venation of right wings; 1b, anterior view of head showing palpi; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.





ANTOLAEA









DELORYCTIS

DELORYCTIS Meyrick

Deloryctis Meyrick, 1934, Exotic Microlepidoptera, 4: 464. (*Typus generis: Deloryctis corticivora* Meyrick, ibid. [monotypy].)

Deloryctis corticivora Meyrick

Plate 209, Figures 1–1e

Deloryctis corticivora Meyrick, 1934, Exotic Microlepidoptera, 4: 464.

"♂♀. 12–15 mm. . . . Java, Telawa, bred August, September from larvae feeding on bark of *Schleichera trijuga* (*Sapindaceae*) (*Dr. L. G. E. Kalshoven*); 5 ex."

Lectotype: ♂, 13 mm., labelled "Seneng, Java, K. bred .8.32." Slide No. 7691. Meyrick recorded this species from "Telawa" an apparent lapsus, since all three of the five original specimens in the Meyrick collection are labelled "Seneng". Kalshoven bred much material from Telawa so the specimens may be mislabelled.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of part of denuded abdomen to show spined and strongly sclerotised tergites; 1d, ventral view of male genitalia with aedeagus removed; 1e, lateral aspect of aedeagus.

DURRANTIA Busck

Durrantia flaccescens Meyrick

Plate 210, Figures 1–1b

Durrantia flaccescens Meyrick, 1925, Exotic Microlepidoptera, 3: 161.

"♂. 17 mm. . . . Peru, Jurimaguas, March (*Parish*); 1 ex."

Type: The above indicated male dated "3.20". Slide No. 4791.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Durrantia montivola Meyrick

Plate 210, Figures 2–2b

Durrantia montivola Meyrick, 1927, Exotic Microlepidoptera, 3: 364.

"♂. 18–20 mm. . . . Texas, Alpine, 6,000–7,000 feet, April, May; 4 ex."

Lectotype: ♂, "Brenster Co., Texas. 7,000 ft., .4.26." Slide No. 4792.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

ASAPHARCHA Meyrick

Asapharcha nephelomicta Meyrick

Plate 210, Figures 3–3c

Asapharcha nephelomicta Meyrick, 1930, Exotic Microlepidoptera, 3: 555.

"♀. 24 mm. . . . New Mexico, Bent, 7,000 feet, October; 1 ex. This interesting insect (of non-American type) appears truly related to the two African species of the genus."

Type: The above indicated specimen dated ".10.27". Slide No. 4794.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.









DURRANTIA and ASAPHARCHA





EPICHOSTIS

EPICHOSTIS Meyrick

Epichostis Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 404. (*Typus generis: Epichostis elephantias* Meyrick, l.c., 405 [monotypy].)

Epichostis elephantias Meyrick

Plate 211, Figures 1–1d

Epichostis elephantias Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 405.

"♂. 15–16 mm. . . . Three specimens, Maskeliya, Ceylon, in April, October and November (*Pole, Green, de Mowbray*)."

Lectotype: ♂, "Maskeliya, Ceylon. Green. .4.05." Slide No. 7655.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

Epichostis antigama (Meyrick)

Plate 212, Figures 1–1b

Hermogenes antigama Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 630.

"♂♀. 18–22 mm. . . . Palni Hills, 6,000 feet (*Campbell*); three specimens."

Lectotype: ♂, 18 mm., "Palni Hills, S. India. Campbell. 6,000'. .06." Slide No. 7652.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Epichostis barathrias (Meyrick)

Plate 212, Figures 2–2b

Hermogenes barathrias Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 629.

"♂♀. 13–17 mm. . . . Patipola, Newera Eliya and Maskeliya, Ceylon, in November and from March to May (*Alston, Pole*); eight specimens."

Lectotype: ♂, 15 mm., "Patipola, Ceylon. A. .5.06." Slide No. 7658.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Epichostis cryphaea (Meyrick)

Plate 212, Figures 3–3b

Hermogenes cryphaea Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 630.

"♀. 18–19 mm. . . . Khasis; three specimens."

Lectotype: ♀, 19 mm., "Khasi Hills, Assam. D. .1906." Slide No. 7651.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

Epichostis dicremna (Meyrick)

Plate 212, Figures 4–4b

Hermogenes dicremna Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 629.

"♂♀. 16–19 mm. . . . Maskeliya, Ceylon, in July and from October to March (*Pole*); sixteen specimens."

Lectotype: ♂, 16 mm., dated ".10.06". Slide No. 7653. Two specimens are missing.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

1

1a



1b

2

2a

2b



3

3a

3b

4

4a



4b

EPICHOSTIS



EPICHOSTIS

Epichostis leptorthra Meyrick

Plate 213, Figures 1–1b

Epichostis leptorthra Meyrick, 1931, *in* Caradja, Bull. Sect. sci. Acad. roum., 14: 74.

"♂. 16–18 mm. . . . 5 ex., July."

Lectotype: ♂, 17 mm., "Kwanhsien, China. F. .7.30." Slide No. 7649. A hair-pencil from hindwing fits into a dorsal groove on abdomen.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Epichostis leucorma (Meyrick)

Plate 213, Figures 2–2b; 3c

Hermogenes leucorma Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 627.

"♂♀. 20–22 mm. . . . Khasis, in July; four specimens."

Lectotype: ♂, 21 mm., "Khasi Hills, Assam. .7.1906." Slide No. 7650.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral edge of harpe showing clasper; 3c, ventral view of male genitalia with aedeagus removed.

Epichostis melanocona (Meyrick)

Plate 213, Figures 3–3b

Hermogenes melanocona Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 629.

"♂. 13–17 mm. . . . Maskeliya, Ceylon, from August to January (*Pole, de Mowbray*); eight specimens."

Lectotype: ♂, 14 mm., "Maskeliya, Ceylon. Pole. .10.06." Slide No. 7654.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Epichostis metrodelta (Meyrick)

Plate 213, Figures 4–4b

Chaliniastis metrodelta Meyrick, 1905, Journ. Bombay Nat. Hist. Soc. 16: 591.

"♂. 15 mm. . . . One specimen, Maskeliya, Ceylon, in March (*Pole*)."

Type: The male denoted above dated ".3.03". Slide No. 7659.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Epichostis microdelta Meyrick

Plate 214, Figures 1–1b

Epichostis microdelta Meyrick, 1927, *in* Caradja, Mem. Sect. Sti. Acad. Română, 4: 422.

". . . Kwan. 28.VII. 1 ♂ [*Talbot*] . . . ♂ 15 mm."

Type: The above indicated male, "Kwanhsien, China. C. 28.7.27." Slide No. 7660.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed (harpes damaged before dissection).

Epichostis stelota (Meyrick)

Plate 214, Figures 2–2b

Hermogenes stelota Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 628.

"♂. 17 mm. . . . Khasis; one specimen."

Type: The male referred to above, labelled "Khasi Hills, Assam. D. .1906." Slide No. 7657. The genitalia of this species are typically oecophorid but I retain the species in this family for the present.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Epichostis tympanias (Meyrick)

Plate 214, Figures 3–3c

Hermogenes tympanias Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 628.

"♂♀. 18–23 mm. . . . Khasis, from June to September; fifteen specimens. Allied to *elephantias*, . . ."

Lectotype: ♂, 18 mm., "Khasi Hills, Assam. .9.1906." Three specimens are missing. Slide No. 7656.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed; 3c, ventral view of denuded abdomen to show modifications of segments.



EPICHOSTIS







EPIMACTIS

EPIMACTIS Meyrick

Epimactis Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 741. (*Typus generis: Epimactis monodoxa* Meyrick, ibid. [monotypy].)

Epimactis monodoxa Meyrick

Plate 215, Figures 1–1e

Epimactis monodoxa Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 741.

"♂♀. 19–23 mm. . . . Bhotan, in June; four specimens (*Dudgeon*)."

Type: Lost. There is one specimen, a female, "Sikkim, 4,500′. GCD. .7.94", 23 mm., which fits both the generic and specific descriptions. I figure this specimen which is labelled "*monodoxa*" in Meyrick's hand. Slide No. 9081.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, ventral view of female genitalia; 1d, detail of genital plate and ostium; 1e, signum.

Epimactis crambalea Meyrick

Plate 216, Figures 1–1b

Epimactis crambalea Meyrick, 1910, Trans. Ent. Soc. London, 1910: 458.

"♂. 16 mm. . . . Java, Bandong; one specimen."

Type: The above indicated specimen with "R. .07" at the bottom of the pin-label. Slide No. 7628.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Epimactis infulata Meyrick

Plate 216, Figures 2–2a

Epimactis infulata Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 779.

"♂. 14 mm. . . . Kegalle, Ceylon (*Alston*); one specimen."

Type: The above indicated male, "Kegalle, Ceylon. CCA. .08." Slide No. 7629.

Figure 2, left wings; 2a, ventral view of male genitalia (aedeagus missing).

Epimactis melithorax Meyrick

Plate 216, Figures 3–3b

Epimactis melithorax Meyrick, 1923, Exotic Microlepidoptera, 2: 613.

"♂. 22 mm. . . . Kumaon, Bhim Tal, 4,500 feet, August (*Fletcher*); 1 ex."

Type: The male denoted above dated "8.18". Slide No. 7632.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.



EPIMACTIS


1

2

3


4

2a

2b


3a


3b

4a


4b


4c

EPIMACTIS

Epimactis spasmodes Meyrick

Plate 217, Figure 1

Epimactis spasmodes Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 779.

"♀. 21 mm. . . . Palnis (*Campbell*); one specimen."

Type: The female, without abdomen, indicated above, "Palni Hills, S. India. Campbell."

Figure 1, left wings.

Epimactis strombodes Meyrick

Plate 217, Figures 2–2b

Epimactis strombodes Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 779.

"♂ 17 mm., ♀ 24 mm. . . . Kandy, Ceylon, in March and May (*Mackwood, Green*); two specimens."

Lectotype: ♂, "Kandy, Ceylon, FMM. .3.09." Slide No. 7630.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Epimactis talantias Meyrick

Plate 217, Figures 3–3b

Epimactis talantias Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 631.

"♂♀. 14–19 mm. . . . Matale and Kandy, Ceylon, in August, November, December, and March (*Pole*); six specimens."

Lectotype: ♂, "Matale, Ceylon. JP. .11.04." Slide No. 7633.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Epimactis turbida Meyrick

Plate 217, Figures 4–4c

Epimactis turbida Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 779.

"♀. 14–18 mm. . . . Khasis, in September; three specimens."

Lectotype: ♀, 16 mm., "Khasi Hills, Assam. .9.1906." Slide No. 7631. All specimens bear identical data.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, detail of bursa copulatrix; 4c, detail of genital plate and ostium.

GEMORODES Meyrick

Gemorodes delphinopa Meyrick

Plate 218, Figures 2–2b

Gemorodes delphinopa Meyrick, 1930, Exotic Microlepidoptera, 4: 13.

"♀. 18 mm. . . . Burma, Pyinmana, March (*Fletcher*); 1 ex. This species indicates the affinity of the genus with *Prothamnodes*, . . ."

Type: The specimen denoted above, labelled "Pyinmana, Burma. AGR. 10.3.18." The initials should read "TBF" or Meyrick recorded Fletcher's name in error. Slide No. 7635.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium.

LASIODICTIS Meyrick

Lasiodictis Meyrick, 1912, Exotic Microlepidoptera, 1: 41. (*Typus generis: Lasiodictis melistoma* Meyrick, ibid. [monotypy].)

Lasiodictis melistoma Meyrick

Plate 218, Figures 1–1d

Lasiodictis melistoma Meyrick, 1912, Exotic Microlepidoptera, 1: 41.

"♂♀. 16–18 mm. . . . Assam, Khasi Hills, in May, June and September; seven specimens."

Lectotype: ♂, 18 mm., "Khasi Hills, Assam. .5.1906." Slide No. 9236.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.



1

1a



1b

1c

1d

2

2a



2b

GEMORODES and LASIODICTIS







LETOGENES and LINOCLOSTIS

LETOGENES Meyrick

Letogenes festalis Meyrick

Plate 219, Figures 3–3b

Letogenes festalis Meyrick, 1930, Exotic Microlepidoptera, 4: 16.

"♀. 19 mm. . . . S. Ussuri, Narva, July (*Kordakoff*); 1 ex."

Type: The above indicated female, "Narva, S. Ussuri. K. .7.21." Slide No. 7638.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

LINOCLOSTIS Meyrick

Linoclostis Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 626. (*Typus generis: Linoclostis gonatias* Meyrick, ibid. [monotypy].) *See* p. 442.

Linoclostis brachyloga Meyrick

Plate 219, Figures 1–1b

Linoclostis brachyloga Meyrick, 1917, Exotic Microlepidoptera, 2: 54.

"♂. 17 mm. . . . Java, Bandong; one specimen."

Type: The male denoted above, "Bandong, Java. R. .07." Slide No. 7676.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Linoclostis musicodes Meyrick

Plate 219, Figures 2–2b

Linoclostis musicodes Meyrick, 1910, Trans. Ent. Soc. London, 1910: 458.

"♂♀. 11–13 mm. . . . Borneo, Kuching (*Hewitt*), in April and August; two specimens."

Lectotype: ♂, "Kuching, Borneo. JH. .4.06." Slide No. 7679.

Figure 2, left wings; 2a, lateral aspect of two denuded tergites to show spines; 2b, ventral view of male genitalia.

Linoclostis gonatias Meyrick

Plate 220, Figures 1–1f

Linoclostis gonatias Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 626.

"♀. 23 mm. . . . Khasis, in June; one specimen."

Type: The female denoted above, "Khasi Hills, Assam. .6.1906." Slide No. 7678.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, ventral view of female genitalia; 1d, dorsal view of denuded abdomen to show spined tergites; 1e, detail of genital plate and ostium; 1f, signum.





LINOCLOSTIS



METATHRINCA

METATHRINCA Meyrick

Metathrinca Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 625. (*Typus generis: Metathrinca ancistrias* Meyrick, l.c., p. 625 [by original designation].)

Metathrinca ophiura Meyrick

Plate 221, Figures 1–1d

Metathrinca ophiura Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 626.

"♂. 13–15 mm. . . . Maskeliya, Puttalam, Eppanela, Wellawaya, and Galle, Ceylon, in April, May, September and December (*Alston, Pole, Green, Fletcher*); six specimens."

Lectotype: ♂, "Galle, Ceylon. BF. 2.5.07." Slide No. 7687.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

Metathrinca ancistrias (Meyrick)

Plate 222, Figures 1–1b

Ptochoryctis ancistrias Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 403.

"♂♀. 13–23 mm. . . . Seven specimens, Maskeliya and Puttalam, Ceylon, from January to May (*Pole, de Mowbray, Alston*); the single Puttalam specimen is much the smallest. . . ."

Lectotype: ♂, 19 mm., "Maskeliya, Ceylon. Pole. .3.06." Slide No. 7688. The Puttalam specimen, referred to above, now forms part of Meyrick's series of *ophiura*.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Metathrinca ceromorpha (Meyrick), new combination

Plate 222, Figures 2–2b

Ptochoryctis ceromorpha Meyrick, 1923, Exotic Microlepidoptera, 2: 612.

"♂. 25 mm. . . . Assam, Shillong, June (*Fletcher*); 2 ex."

Lectotype: ♂, "Shillong, Assam. Fletcher. .6.20." Slide No. 7701. The second specimen is missing.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Metathrinca illuvialis (Meyrick,) new combination

Plate 222, Figures 3–3c

Ptochoryctis illuvialis Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 778.

"♀. 26 mm. . . . Khasis, in April; one specimen."

Type: The female denoted above, "Khasi Hills, Assam. .4.1906." Slide No. 7698. The abdomen is spined.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Metathrinca intacta (Meyrick), new combination

Plate 222, Figures 4–4b

Ptochoryctis intacta Meyrick, 1938, *in* Caradja and Meyrick, Deuts. Ent. Zeit. Iris, 52: 14.

". . . ♂. 25 mm., . . . Yül., 2,800 m., VIII, 1 ex. Nearest the Indian *inviolata* Meyr., . . ."

Type: The above indicated male, "Likiang, China. H. .8.34." Slide No. 7686.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a

1b

2

2a


2b

3

3a


3b 3c

4

4a

4b

METATHRINCA



1

1a

1b



2

2a



2b

3

3a

3b

3c

4

4a

4b

METATHRINCA

Metathrinca loranthivora (Meyrick), new combination

Plate 223, Figures 1–1b

Ptochoryctis loranthivora Meyrick, 1937, Exotic Microlepidoptera, 5: 150.

"♂♀. 15–18 mm. . . . Java, Telawa, bred July, August, from larvae feeding in shoots of *Loranthus* (*Dr. L. G. Kalshoven*); 6 ex."

Lectotype: ♂, 16 mm., dated ".7.35". Slide No. 7692. Three specimens are missing. Abdomen spined.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Metathrinca memnon Meyrick

Plate 223, Figures 2–2b

Metathrinca memnon Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 778.

"♂. 24 mm. . . . Hakgala, Ceylon, in April (*Green*); one specimen."

Type: The male denoted above, "Hakgala, Ceylon. EEG. .4.07." Slide No. 7684.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Metathrinca parabola (Meyrick), new combination

Plate 223, Figures 3–3c

Ptochoryctis parabola Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 778.

"♀. 25 mm. . . . N. Coorg, 3,500 feet, in May (*Newcome*); one specimen. Similar to *rosaria*, . . ."

Type: The above indicated female, "Dibidi, N. Coorg. Newcome. 19.5.07." Slide No. 7685.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signum; 3c, detail of genital plate and ostium.

Metathrinca rosaria (Meyrick), new combination

Plate 223, Figures 4–4b

Ptochoryctis rosaria Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 740.

"♂♀. 23–24 mm. . . . Bhotan, in April and May; two specimens (*Dudgeon*)."

Lectotype: ♂, "Bhotan. GCD. 16.5.95." Slide No. 7696.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

MYRIOPLEURA Meyrick

Myriopleura Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 405. (*Typus generis: Xyloryeta psilotis* Meyrick, 1905, Journ. Bombay Nat. Hist. Soc., 16: 602 [by original designation].)

Myriopleura psilotis (Meyrick)

Plate 224, Figures 1–1d

Xyloryctis psilotis Meyrick, 1905, Journ. Bombay Nat. Hist. Soc., 16: 602.

"♂. 16–17 mm. . . . Four specimens, Peradeniya and Yatiyantola, Ceylon, in March, September and November (*Green*)."

Lectotype: ♂, "Peradeniya, Ceylon. Green. .11.01." Slide No. 7622. This genus is a synonym of *Odites*.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.



MYRIOPLEURA


1


1a


1b

1c

1d


2

2a


2b


3

3a

3b

NEOSPASTIS

NEOSPASTIS Meyrick

Neospastis Meyrick, 1917, Exotic Microlepidoptera, 2: 59. (*Typus generis: Agriophara encrpyhias* Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 743 [by original designation].)

Neospastis encryphias (Meyrick)

Plate 225, Figures 1–1d

Agriophara encryphias Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 743.

"♂♀. 23–24 mm. . . . Khasi Hills; two specimens (*Swinhoe*)."

Lectotype: ♂, "Khasi Hills, Assam. CS. .1906." Slide No. 7582.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

Neospastis calpidias Meyrick

Plate 225, Figures 2–2b

Neospastis calpidias Meyrick, 1917, Exotic Microlepidoptera, 2: 59.

"♂. 23 mm. . . . S. India, Nilgiris (Pykara), 7,000 feet, in March (*Andrewes*); one specimen. . . ."

Type: The above indicated specimen dated "III.13". Slide No. 7583.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Neospastis ichnaea (Meyrick)

Plate 225, Figures 3–3b

Stenoma ichnaea Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 23: 118.

"♂♀. 22–25 mm. . . . Anshi, Kanara, bred in January (*Maxwell*); six specimens."

Lectotype: ♂, 22 mm., dated "17.1.12". Slide No. 7584.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

NEPHANTIS Meyrick

Nephantis Meyrick, 1905, Journ. Bombay Nat. Hist. Soc., 16: 603. (*Typus generis: Nephantis serinopa* Meyrick, ibid. [monotypy].)

Nephantis serinopa Meyrick

Plate 226, Figures 1–1d

Nephantis serinopa Meyrick, 1905, Journ. Bombay Nat. Hist. Soc., 16: 603.

"♂♀. 20–28 mm. . . . Five specimens, Batticaloa, Ceylon, in April (*Green*)."

Lectotype: ♂. The five specimens bear identical data and are dated ".4.00". Slide No. 7646.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.



NEPHANTIS


1

1a

1b

2


1c

2a

2b


3

3a


3b

4


4a

4b

ODITES

ODITES Walsingham

Odites actuosa Meyrick

Plate 227, Figures 1–1c

Odites actuosa Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 780.

"♀. 18 mm. . . . N. Coorg, 3,500 feet, in February (*Newcome*). Resembles *sphendonistis*. . . ."

Type: The female indicated above labelled "Dibidi, N. Coorg. Newcome. 23.2.07." Slide No. 7601.

Figure 1, left wings; 1a ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signum.

Odites agraula Meyrick

Plate 227, Figures 2–2b

Odites agraula Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 632.

"♂. 19–23 mm. . . . Palni Hills, 6,000 feet (*Campbell*); five specimens."

Lectotype: ♂, 22 mm., "Palni Hills, S. India. Campbell." Slide No. 7604.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Odites analogica Meyrick

Plate 227, Figures 3–3b

Odites analogica Meyrick, 1917, Exotic Microlepidoptera, 2: 57.

"♀. 12 mm. . . . Comoro Is., Anjouan, in July (*Leigh*); one specimen."

Type: The above indicated specimen. Slide No. 4471.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

Odites anisocarpa Meyrick

Plate 227, Figures 4–4b

Odites anisocarpa Meyrick, 1930, Exotic Microlepidoptera, 4: 17.

"♂. 20 mm. . . . Madagascar, Forest of Antsianaka (*Humblot*); 1 ex. (Coll. Paravicini)."

Type: The above indicated specimen, "Forest of Antsianaka, Madagascar. L. Humblot. 1888." Slide No. 4506.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Odites aspasta Meyrick

Plate 228, Figures 1–1b

Odites aspasta Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 632.

"♂♀. 23–24 mm. . . . Khasis, in May and September; three specimens."

Lectotype: ♂, "Khasi Hills, Assam. .5.1906." Slide No. 7603.

Figure 1, left wings; 1a, ventral view of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Odites atmopa Meyrick

Plate 228, Figures 2–2c

Odites atmopa Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 780.

"♀. 26–27 mm. . . . Kandy, Ceylon, in December (*Green*, *Mackwood*); two specimens. Allied to *paracyrta*."

Lectotype: ♀, "Kandy, Ceylon. FM. .11." Slide No. 7616.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signa.

Odites atonopa Meyrick

Plate 228, Figures 3–3b

Odites atonopa Meyrick, 1918, Exotic Microlepidoptera, 2: 197.

"♂. 20 mm. . . . N.W. India, Abbottabad, June (*Fletcher*); one specimen."

Type: The male indicated above dated ".6.16". Slide No. 7625.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Odites balanospila Meyrick

Plate 228, Figures 4–4a

Odites balanospila Meyrick, 1930, Exotic Microlepidoptera, 3: 555.

[balanospila]

"♂. 17 mm. . . . Sierra Leone, Njala, September (*E. Hargreaves*); 1 ex. (Brit. Mus.)."

Type: The above indicated specimen so marked in the British Museum. Slide No. 4488.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*. The apex of the aedeagus is curved ventrad but in the illustration this is foreshortened.



ODITES

1


1a


2

2a


2b

3

3a

3b


4

4a

4b

ODITES

Odites balanospila Meyrick

Plate 229, Figures 1–1a

Odites haplogramma Meyrick, 1930, Exotic Microlepidoptera, 4: 17 (new synonymy).

[haplogramma]

"♂. 16–17 mm. . . . Cameroons, Lododorf, January, February (*Conradt*); 2 ex. (Coll. Paravicini)."

Type: The specimen so marked in the British Museum, dated "30–I". The locality should read "Lolodorf". Slide No. 4503.

Figure 1, left wings; 1a, ventral view of male genitalia with aedeagus *in situ*.

Odites brachyclista Meyrick

Plate 229, Figures 2–2b

Odites brachyclista Meyrick, 1928, Exotic Microlepidoptera, 3: 434.

"♂. 22 mm. . . . Philippines, Mt. Makiling, Luzon (*Baker*); 1 ex."

Type: The above indicated male with "B. .27" at the bottom of the pin-label. Slide No. 7618.

Figure 2, left wings; 2a, lateral view of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Odites cataxantha Meyrick

Plate 229, Figures 3–3b

Odites cataxantha Meyrick, 1915, Exotic Microlepidoptera, 1: 379.

"♀. 14 mm. . . . Madagascar, Antananarivo; one specimen."

Type: The above indicated specimen, "Antananarivo, Madagascar. R. .11." Slide No. 4460.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

Odites choricopa Meyrick

Plate 229, Figures 4–4b

Odites choricopa Meyrick, 1931, *in* Caradja, Bull. Sect. sci. Acad. roum., 14: 74.

"♂♀. 17–18 mm. . . . 4 ex., July. Nearest to *pragmatias*."

Lectotype: ♂, "Kwanhsien, China. F. .7.28." Slide No. 7613.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Odites circiformis Meyrick

Plate 230, Figures 1–1b

Odites circiformis Meyrick, 1930, Exotic Microlepidoptera, 4: 18.

"♂ 24 mm., ♀ 34 mm. . . . Cameroons, Bitje, Ja River, October (*G. L. Bates*); 2 ex."

Lectotype: The male. Slide No. 4491.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Odites citrantha Meyrick

Plate 230, Figures 2–2a

Odites citrantha Meyrick, 1908, Proc. Zool. Soc. London, 1908: 729.

"♂. 21 mm. . . . Natal, Durban, in August (*Leigh*); one specimen."

Type: The above indicated specimen. Slide No. 4496.

Figure 2, left wings; 2a, ventral view of male genitalia with aedeagus *in situ*.

Odites consecrata Meyrick

Plate 230, Figures 3–3b

Odites consecrata Meyrick, 1917, Exotic Microlepidoptera, 2: 57.

"♂. 15 mm. . . . Madagascar, Antananarivo; one specimen."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 4458.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Odites continua Meyrick

Plate 230, Figures 4–4b

Odites continua Meyrick, 1935, *in* Caradja and Meyrick, Materialien zu einer Microlepidopteren Fauna der Chinesischen Provinzen Kiangsu, Chekiang und Hunan, 84.

". . . ♂, 13 mm. . . . 1 ex., Tienmushan, VIII. Nearest the Indian *spoliatrix* Meyr."

Type: The male denoted above, labelled "Tien-Mu-Shan, China. C. 5,000′. .8.32." Slide No. 7602.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a

1b

2

2a

3

3a

3b


4


4a


4b

ODITES

1

1a 1b


2c

2


2a


2b


3

3a


3b


4

4a

4b

ODITES

Odites crossophanta Meyrick

Plate 231, Figures 1–1b

Odites crossophanta Meyrick, 1930, Exotic Microlepidoptera, 4: 17.

"♀. 16 mm. . . . Cameroons, Lolodorf, January (*Conradt*); 1 ex. (Coll. Paravicini)."

Type: The above indicated specimen so marked in the British Museum. Slide No. 4487.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium.

Odites diopta Meyrick

Plate 231, Figures 2–2c

Odites diopta Meyrick, 1917, Exotic Microlepidoptera, 2: 57.

"♀. 12 mm. . . . French Congo, Fort Crampel; one specimen. Generally similar to the Indian *Swinhoei*, . . ."

Type: The above indicated specimen with "LeM. .15" at the bottom of the pin-label. Slide No. 4459.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, bursa copulatrix showing multiple signa; 2c, detail of genital plate and ostium.

Odites encarsia Meyrick

Plate 231, Figures 3–3b

Odites encarsia Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 634.

"♂. 16–17 mm. . . . N. Coorg, 3,500 feet, in May (*Newcome*); two specimens."

Lectotype: ♂, 16 mm., "Dibidi, N. Coorg. Newcome. 14.5.07." Slide No. 7607.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Odites eriopa Meyrick

Plate 231, Figures 4–4b

Odites eriopa Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 633.

"♂♀. 16–18 mm. . . . Khasis, in July and August; six specimens."

Lectotype: ♂, "Khasi Hills, Assam. .7.1906." Slide No. 7605.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Odites euphema Meyrick

Plate 232, Figures 1–1c

Odites euphema Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 780.

"♀. 15–17 mm. . . . Mundgod and Pola, Kanara, in October (*Maxwell*); N. Coorg, 3,500 feet, in August (*Newcome*); three specimens."

Lectotype: ♀, "Pola, Kanara. RM. 20.10.11." Slide No. 7587.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium; 1c, signa.

Odites fruticosa Meyrick

Plate 232, Figures 2–2b

Odites fruticosa Meyrick, 1915, Exotic Microlepidoptera, 1: 380.

"♂. 17 mm. . . . Coorg, Dibidi, 3,500 feet, in February (*Newcome*); one specimen."

Type: The male denoted above, "Dibidi, N. Coorg. LN. .2.12." Slide No. 7611.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Odites furfurosa (Meyrick)

Plate 232, Figures 3–3b

Myriopleura furfurosa Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 405.

"♂♀. 14–19 mm. . . . Six specimens, Puttalam, Ceylon, in January, February, September and October (*Pole*)."

Lectotype: ♂, 14 mm., dated ".10.04". Slide No. 7623.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Odites glaphyra Meyrick

Plate 232, Figures 4–4b

Odites glaphyra Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 632.

"♂. 26 mm. . . . Sikkim, in March; one specimen."

Type: The male indicated above, "Sikkim. P. 26.3.89." Slide No. 7600.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1b


1c


2


2b


3

3b


4


4b

ODITES


1

1a


1b

2

2a


2b


3


3a

4

4a

ODITES

Odites gomphias Meyrick

Plate 233, Figures 1–1b

Odites gomphias Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 633.

"♂♀. 15–23 mm. . . . Maskeliya, Maturatta, Puttalam, and Eppawela, Ceylon; in March, September and November (*Pole, Green, de Mowbray*); five specimens."

Lectotype: ♂, "Eppawela, Ceylon. EEG. .9.05." Slide No. 7606.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Odites haplogramma Meyrick

See Odites balanospila Meyrick, p. 461.

Odites haplonoma Meyrick

Plate 233, Figures 2–2b

Odites haplonoma Meyrick, 1915, Exotic Microlepidoptera, 1: 379.

"♀. 20 mm. . . . Madagascar, Antanosy; one specimen."

Type: The above indicated specimen with "L. .93" at the bottom of the pin-label. Slide No. 4457.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium.

Odites hemipercna Meyrick

Plate 233, Figures 3–3a

Odites hemipercna Meyrick, 1914, Exotic Microlepidoptera, 1: 282.

"♂♀. 10 mm. . . . Nyassaland, Mt. Mlanje, in March (*Neave*); three specimens. . . . Type in British Museum."

Type: The male so marked in the British Museum, labelled "Mt. Mlanje, Nyassaland. 17–III–1913. S. A. Neave." Slide No. 4489.

Figure 3, left wings; 3a, ventral view of male genitalia with aedeagus *in situ*.

Odites heptasticta Meyrick

Plate 233, Figures 4–4a

Odites heptasticta Meyrick, 1914, Exotic Microlepidoptera, 1: 281.

"♂♀. 18 mm. . . . Nyassaland, Mt. Mlanje, in January and April (*Neave*); two specimens. Type in British Museum."

Type: The male so marked in the British Museum. Slide No. 4482.

Figure 4, left wings; 4a, ventral view of male genitalia with aedeagus *in situ*.

Odites hermatica Meyrick

Plate 234, Figures 1–1b

Odites hermatica Meyrick, 1915, Exotic Microlepidoptera, 1: 378.

"♂♀. 15–16 mm. . . . Comoro Is., Mayotte, in June (*Leigh*); four specimens."

Lectotype: ♂, with "L. .6.11" at the bottom of the pin-label. Slide No. 4468.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Odites holocitra Meyrick

Plate 234, Figures 2–2c

Odites holocitra Meyrick, 1925, Exotic Microlepidoptera, 3: 156.

"♀. 31 mm. . . . Cameroons, Bitje, Ja R., 2,000 feet (*Bates*); 1 ex."

Type: The above indicated specimen, "Bitje, Ja R., Cameroons, 2,000 ft. Dry season. G. L. Bates.' Slide No. 4493.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signa.

Odites holotorna Meyrick

Plate 234, Figures 3–3c

Odites holotorna Meyrick, 1925, Exotic Microlepidoptera, 3: 157.

"♀. 26 mm. . . . Cameroons, Bitje, Ja R., 2,000 feet (*Bates*); 1 ex."

Type: The above indicated specimen with "Dry season, G. L. Bates" on the label in addition to the data recorded above. Slide No. 4492.

Figure 3, left wings; 3a, signum; 3b, detail of genital plate and ostium; 3c, ventral view of female genitalia.

Odites incallida Meyrick

Plate 234, Figures 4–4c

Odites incallida Meyrick, 1915, Exotic Microlepidoptera, 1: 381.

"♀. 17 mm. . . . S. India, Cuddapah, 4,000 feet (*Campbell*); one specimen."

Type: The female denoted above, "Cuddapah, S. India. WHC. 4,000′. .08." Slide No. 7626.

Figure 4, left wings; 4a, ventral view of female genitalia; 4b, signa; 4c, detail of genital plate and ostium.

1

1a

1b


2

2a

2b


3


3a

3b


2c

3c


4

4a

4b

4c

ODITES



1

1a

1b

2

2a

2b



3

3a



3b



4

5

ODITES

Odites isocentra (Meyrick)

Plate 235, Figures 1–1b

Myriopleura isocentra Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 406.

"♂. 11–12 mm. . . . Two specimens, Puttalam, Ceylon, in October and December (*Pole*)."

Lectotype: ♂, dated ".12.04". Slide No. 7624.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Odites leucostola (Meyrick), new combination

Plate 235, Figures 2–2b

Protobathra leucostola Meyrick, 1921, Exotic Microlepidoptera, 2: 436.

"♂. 17 mm. . . . Japan, Tokio, July; 1 ex."

Type: The male denoted above, "Tokio, Japan. N. .7.19." Slide No. 9038.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Odites lividula Meyrick

Plate 235, Figures 3–3b

Odites lividula Meyrick, 1932, Exotic Microlepidoptera, 4: 286.

"♂. 18 mm. . . . Japan, Tokio, Inokasira, July (*S. Issiki*); 1 ex. Nearest the Chinese *xenophaea*."

Type: The specimen denoted above, "Tokio, Japan. SI. 16.7.20." Slide No. 7614.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Odites malivora Meyrick

Plate 235, Figure 4

Odites malivora Meyrick, 1930, Exotic Microlepidoptera, 3: 555.

"♀. 22 mm. . . . Manchuria, bred August from larva rolling leaf of *Pyrus malus*; 1 ex. Near *assidua*."

Type: The female denoted above, without abdomen, dated ".8.28".

Figure 4, left wings.

Odites matura Meyrick

Plate 235, Figure 5

Odites matura Meyrick, 1914, Exotic Microlepidoptera, 1: 282.

"♂. 19 mm. . . . Nyassaland, Mt. Mlanje, in April (*Neave*); one specimen. Type in British Museum."

Type: The above indicated specimen so marked in the British Museum. The type consists of the two forewings, part of the thorax and head.

Figure 5, left forewing.

Odites metaclista Meyrick

Plate 236, Figures 1–1b

Odites metaclista Meyrick, 1915, Exotic Microlepidoptera, 1: 379.

"♀. 14–15 mm. . . . Madagascar, Antananarivo; two specimens."

Lectotype: One of the above indicated specimens with "R. .11" at the bottom of the pin-label. Slide No. 4461.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium.

Odites microbolista Meyrick

See Odites pelochrosta Meyrick, p. 477.

Odites navigatrix (Meyrick)

Plate 236, Figures 2–2b

Xylorycta navigatrix Meyrick, 1912, Trans. Ent. Soc. London, 1911: 705.

"♂. 25 mm. . . . Colombia, San Antonio, 5,800 feet, in November; one specimen."

Type: The specimen denoted above dated ".11.07". Slide No. 4793. This species is obviously misplaced generically, appears to be nearest to *Acria*, and probably will require a new genus.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Odites notocapna Meyrick

Plate 236, Figures 3–3b

Odites notocapna Meyrick, 1925, *in* Caradja, Mem. Sect. sti. Acad. Română, 3: 382.

"♂. 16 mm. . . . Canton; one example."

Type: The male denoted above, "Canton, China. C. .24." Slide No. 7694.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Odites oligectis Meyrick

Plate 236, Figure 4

Odites oligectis Meyrick, 1917, Exotic Microlepidoptera, 2: 56.

"♂. 17 mm. . . . S. India, Shevaroys, 4,500 feet, in December (*Fletcher*); one specimen."

Type: The above indicated specimen, "Shevaroy Hills, S. India. TBF. 4,500′. .12.13." The abdomen of the type is missing.

Figure 4, left wings.

Odites orthometra Meyrick

Plate 236, Figure 5

Odites orthometra Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 634.

"♂♀. 16–18 mm. . . . Maskeliya, Ceylon, in April (*Pole*), one ♂ (type); Khasis in August, one ♀."

Type: The specimen designated by Meyrick dated ".4.06". The abdomen of the type is missing.

Figure 5, left wings.


1

1a

1b


2

2a

2b


3

3a

3b


4


5

ODITES

1

1a

1b


2

2a

2b

3

3a

3b


4

4a

4b

ODITES

Odites pancyclia Meyrick

Plate 237, Figures 1–1b

Odites pancyclia Meyrick, 1928, Exotic Microlepidoptera, 3: 434.

"♂. 18 mm. . . . Philippines, Mt. Makiling, Luzon (*Baker*); 1 ex."

Type: The above indicated male with "B. .27" at the bottom of the pin-label. Slide No. 7617.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Odites paracyrta (Meyrick), new combination

Plate 237, Figures 2–2b

Xylorycta paracyrta Meyrick, 1905, Journ. Bombay Nat. Hist. Soc., 16: 602.

"♂. 23 mm. . . . One specimen, Haputale, Ceylon, in February (*Green*)."

Type: The above indicated specimen, "Haputale, Ceylon. EEG. .2.00." Slide No. 7627.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Odites pelochrosta Meyrick

Plate 237, Figures 3–3b

Odites pelochrosta Meyrick, 1933, Exotic Microlepidoptera, 4: 434.
Odites microbolista Meyrick, 1937, Exotic Microlepidoptera, 5: 152 (new synonymy).

[pelochrosta]

"♀. 20 mm. . . . Sierra Leone, Njala, July (*E. Hargreaves*); 1 ex. (British Museum)."

Type: The specimen so marked in the British Museum, without abdomen. The specimen was erroneously recorded as a female.

No figure.

[microbolista]

"♀. 22 mm. . . . Uganda, Kampala, bred April from larvae rolling leaves of *Coffea* (*H. Hargreaves*); 1 ex. (Brit. Mus.)."

Type: The above indicated specimen so marked in the British Museum. Slide No. 4485.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

Odites perissopis Meyrick

Plate 237, Figures 4–4b

Odites perissopis Meyrick, 1936, Exotic Microlepidoptera, 5: 27.

"♂. 19 mm. . . . Japan, Tokyo, July (*Prof. S. Issiki*); 1 ex."

Type: The above indicated specimen dated "12.7.32". Slide No. 7609.

Figure 4, left wings; 4a, ventral view of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Odites plocamopa Meyrick

Plate 238, Figures 1–1b

Odites plocamopa Meyrick, 1935, *in* Caradja and Meyrick, Materialien zu einer Microlepidopteren Fauna der Chinesischen Provinzen Kiangsu, Chekiang und Hunan, 84.

". . . ♂, 18 mm. . . . 1 ex., Tienmushan, IX. Allied to *atmopa* Meyr. . . ."

Type: The male indicated above, "Tien-Mu-Shan, China. C. 5,000'. .9.32." Slide No. 7621.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Odites practoria Meyrick

Plate 238, Figures 2–2b

Odites practoria Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 631.

"♂♀. 15–18 mm. . . . Khasis, in June and September; seven specimens."

Lectotype: ♂, 17 mm., "Khasi Hills, Assam. .9.1906." Slide No. 7599.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Odites pragmatias Meyrick

Plate 238, Figures 3–3c

Odites pragmatias Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 780.

"♀. 17–18 mm. . . . Pykara, Nilgiris, 7,000 feet, in April (*Andrewes*); two specimens."

Lectotype: ♀, 18 mm., "Nilgiri Hills, Pykara, 7,000 ft., H. L. Andrewes. IV.13." Slide No. 7586.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, signa; 3c, detail of posterior section of ductus bursae and ostium.

Odites procellosa Meyrick

Plate 238, Figures 4–4b

Odites procellosa Meyrick, 1908, Proc. Zool. Soc. London, 1908: 729.

"♂♀. 17–18 mm. . . . S. Nigeria, Wari, in May and August; three specimens."

Lectotype: ♂, "Wari, Benin. R. .8.97." Slide No. 4484. All specimens are males.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1

1a

1b


2

2a

2b


3

3a

3b


3c

4


4a


4b



ODITES

Odites prosedra Meyrick

Plate 239, Figures 1–1c

Odites prosedra Meyrick, 1915, Exotic Microlepidoptera, 1: 378.

"♀. 20 mm. . . . Nigeria, Ogrugu; one specimen."

Type: The above indicated specimen with "D. .06" at the bottom of the pin-label. Slide No. 4481.

Figure 1, left wings; 1a, bursa copulatrix and ductus bursae; 1b, posterior part of female genitalia; 1c, detail of genital plate and ostium.

Odites repetita Meyrick

(No figure.)

Odites repetita Meyrick, 1932, Exotic Microlepidoptera, 4: 286.

"♀. 16 mm. . . . Uganda, Bukaloba, October (*T. W. Chorley*); 1 ex. (Brit. Mus.)."

Type: The specimen denoted above, so marked in the British Museum. The type is badly worn and the tips of the wings and abdomen are gone. The species appears to be nearest *heptasticta* if not identical with it.

Odites semisepta Meyrick

Plate 239, Figures 2–2c

Odites semisepta Meyrick, 1930, Exotic Microlepidoptera, 4: 18.

"♀. 14 mm. . . . Cameroons, Johann-Albrecht's Höhe Station (*Conradt*); 1 ex. (Coll. Paravicini). Allied to *metaclista*."

Type: The specimen so marked in the British Museum. Slide No. 4504.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium; 2c, signa.

Odites siccinervis Meyrick

Plate 239, Figures 3–3b

Odites siccinervis Meyrick, 1930, Exotic Microlepidoptera, 4: 17.

"♂. 18 mm. . . . Uganda, Kampala, bred October from larva on leaf of *Coffea* (*Hargreaves*); 1 ex. (Brit. Mus.)."

Type: The specimen so marked in the British Museum. Slide No. 4486.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed (tegumen and gnathos broken off).

Odites sphenidias Meyrick

Plate 239, Figure 4

Odites sphenidias Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 781.

"♂. 16 mm. . . . Khasis, in October; one specimen."

Type: The above indicated specimen, "Khasi Hills, Assam. .10.1906", without abdomen.

Figure 4, left wings.

Odites periscias Meyrick

Plate 239, Figure 5

Odites periscias Meyrick, 1928, Exotic Microlepidoptera, 3: 434.

"♂. 19 mm. . . . Philippines, Mt. Makiling, Luzon (*Baker*); 1 ex."

Type: The above indicated male, without abdomen, with "B. .27" at the bottom of the pin-label.

Figure 5, left wings.

Odites sucinea Meyrick

Plate 240, Figures 1–1b

Odites sucinea Meyrick, 1915, Exotic Microlepidoptera, 1: 378.

"♂. 24 mm. . . . Natal, Pinetown, in January (*Leigh*); one specimen."

Type: The above indicated specimen. Slide No. 4490.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Odites typota Meyrick

Plate 240, Figures 2–2c

Odites typota Meyrick, 1915, Exotic Microlepidoptera, 1: 379.

"♀. 17–18 mm. . . . Madagascar, Antananarivo; two specimens."

Type: ♀, "Antananarivo, Madagascar. M. .11." Slide No. 4473.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of ostium and posterior part of ductus bursae; 2c, bursa copulatrix and signum.

Odites velipotens Meyrick

Plate 240, Figures 3–3b

Odites velipotens Meyrick, 1935, Exotic Microlepidoptera, 4: 572.

"♂. 18 mm. . . . W. China, Mt. Omei, 4,000 feet, July (*Franck*); 1 ex."

Type: The above indicated male, "Mt. Omei, W. China. 4,000′. .7.32." Slide No. 7615.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Odites xenophaea Meyrick

Plate 240, Figures 4, 4b, 4c

Odites xenophaea Meyrick, 1931, *in* Caradja, Bull. Sect. sci. Acad. roum., 14: 74.

"♂ 19 mm., ♀ 22–24 mm., . . . 3 ex., July."

Lectotype: ♂, "Kwanhsien, China. F. .7.28." Slide No. 7612.

Figure 4, left wings; 4b, lateral aspect of aedeagus; 4c, ventral view of male genitalia with aedeagus removed.

1

1a

1b


2

2a

2b

2c


3

3a

3b


4

4b

4c

ODITES



PHTHONERODES

PHTHONERODES Meyrick

Phthonerodes Meyrick, 1890, Trans. R. Soc. South Australia, 14: 44. (*Typus generis: Phthonerodes scotarcha* Meyrick, l.c., 45 [monotypy].)

Phthonerodes scotarcha Meyrick

Plate 241, Figures 1–1d

"Both sexes 13–21 mm., . . . Quorn and Wirrabara, South Australia, in October; taken plentifully on a fence."

Lectotype: ♂, 14 mm., "Quorn, S. Australia. 23/10/82." There are ten specimens in the Meyrick collection which probably represent all the original material. Slide No. 8803.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

PROCOMETIS Meyrick

Procometis mistharna (Meyrick), new combination

Plate 242, Figures 1–1b

Odites mistharna Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 634.

"♂♀. 11–14 mm. . . . Puttalam and Trincomali, Ceylon, from June to September (*Pole*, *Fletcher*); five specimens."

Lectotype: ♂, 11 mm., "Puttalam, Ceylon. Pole. .8.04." Slide No. 7608.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Procometis sphendonistis (Meyrick), new combination

Plate 242, Figures 2–2b

Odites sphendonistis Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 634.

"♂. 14 mm. . . . Puttalam, Ceylon, in September (*Pole*); one specimen."

Type: The above indicated male dated ".9.04". Slide No. 7598.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Procometis spoliatrix (Meyrick), new combination

Plate 242, Figures 3–3b

Odites spoliatrix Meyrick, 1916, Exotic Microlepidoptera, 1: 509.

"♂ 16 mm., ♀ 18 mm., S. India, Coimbatore, 1 ♂ bred in August from larva forming gallery in nest of social spider (*Fletcher*); Bombay, Kokan, 1 ♀ (*Young*). Perhaps nearest *furfurosa*. . . ."

Lectotype: ♂, "Coimbatore, India. TBF. Bred 13.8.15." Slide No. 7597.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Procometis trochala Meyrick

Plate 242, Figures 4–4b

Procometis trochala Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 635.

"♂ 21 mm., ♀ 35 mm., Pusa, Bengal, bred in June from larvae feeding on dried leaves of sugar-cane (*Maxwell-Lefroy*); two specimens."

Lectotype: ♂, "Pusa, Bengal. HML. Bred .6.06." Slide No. 7647. A note on an elongate white label, beneath the date label, reads "larva on leaves of sugar-cane."

Figure 4, left wings; 4a, lateral aspect of abdomen to show spined tergites; 4b, ventral view of male genitalia with aedeagus (unusually small) *in situ*.

1

1a


1b


2

2a


2b

3

3a

3b

4


4a

4b

PROCOMETIS





PROTHAMNODES

PROTHAMNODES Meyrick

Prothamnodes Meyrick, 1923, Exotic Microlepidoptera, 2: 613. (*Typus generis: Prothamnodes platycycla* Meyrick, ibid. [monotypy].)

Prothamnodes platycycla Meyrick

Plate 243, Figures 1–1d

Prothamnodes platycycla Meyrick, 1923, Exotic Microlepidoptera, 2: 613.

"♀. 23 mm. . . . Burma, Thaton, March (*Fletcher*); 1 ex."

Type: The above indicated female dated ".3.18". Slide No. 7634.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, ventral view of female genitalia; 1d, detail of genital plate and ostium.

PTOCHORYCTIS Meyrick

Ptochoryctis Meyrick, 1894, Trans. Ent. Soc. London, 1894: 19. (*Typus generis: Ptochoryctis eremopa* Meyrick, ibid. [monotypy].)

Ptochoryctis eremopa Meyrick

Plate 244, Figures 1–1d

Ptochoryctis eremopa Meyrick, 1894, Trans. Ent. Soc. London, 1894: 19.

"♂. 17 mm. . . . Koni; one specimen."

Type: The male indicated above, "Koni, Burma. NM. .88." Slide No. 7683.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of denuded abdomen to show spined tergites; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.







PTOCHORYCTIS


1

1a

1b


2

2a

2b


2c


3

3a


3b

4

4a 4b

4c

PTOCHORYCTIS

Ptochoryctis acrosticta Meyrick

Plate 245, Figures 1–1b

Ptochoryctis acrosticta Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 403.

"♂♀. 11–15 mm. . . . Seven specimens, Puttalam and Hambantota, Ceylon, in January, February and October (*Pole*)."

Lectotype: ♂, "Hambantota, Ceylon. JP. .04." Slide No. 7690. Two specimens are missing.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Ptochoryctis alma (Meyrick)

Plate 245, Figures 2–2c

Amorbaea alma Meyrick, 1908, Journ. Bombay Nat. Hist. Soc., 18: 627.

"♀. 24 mm. . . . Coorg, 5,000 feet, in June (Newcome); one specimen."

Type: The female denoted above, "Dibidi, N. Coorg. *Newcome.* 15.6.06." Slide No. 7699.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, signum; 2c, detail of genital plate and ostium.

Ptochoryctis anguillaris Meyrick

Plate 245, Figures 3–3b

Ptochoryctis anguillaris Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 778.

"♂♀. 12–15 mm. . . . Hambantota and Maskeliya, Ceylon, in October and December (*Fletcher, Alston*); two specimens."

Lectotype: ♂, "Maskeliya, Ceylon. Alston. .12.06." Slide No. 7697.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Ptochoryctis galbanea (Meyrick)

Plate 245, Figures 4–4c

Amorbaea galbanea Meyrick, 1914, Journ. Bombay Nat. Hist. Soc., 22: 778.

"♀. 30–32 mm. . . . Maskeliya, Ceylon, from February to April (*Green, Alston*); three specimens. . . ."

Lectotype: ♀, 31 mm., "Maskeliya, Ceylon. Green. .4.05." Slide No. 7703.

Figure 4, left wings; 4a, signum; 4b, ventral view of female genitalia; 4c, detail of genital plate and ostium.

Ptochoryctis ochrograpta Meyrick

Plate 246, Figures 1–1c

Ptochoryctis ochrograpta Meyrick, 1923, Exotic Microlepidoptera, 2: 612.

"♀. 22 mm. . . . Burma, Kalaw, 4,300 feet, May (*Fletcher*); 1 ex."

Type: The above indicated female dated ".5.18". Slide No. 7702.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, signum; 1c, detail of genital plate and ostium.

Ptochoryctis scionota Meyrick

Plate 246, Figures 2–2b

Ptochoryctis scionota Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 403.

"♂. 16–17 mm. . . . Three specimens, Puttalam, Ceylon, in April and August (*Pole*)."

Lectotype: ♂, 17 mm., dated ".8.04". Slide No. 7689.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Ptochoryctis simbleuta Meyrick

Plate 246, Figures 3–3b

Ptochoryctis simbleuta Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 18: 150.

"♂♀. 11–17 mm. . . . Gazepore, Assam, in March and April; eight specimens bred (*Antram*)."

Lectotype: ♂, 12 mm., "Gazepore, Assam. A. bred. .3.07." Slide No. 7695.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, lateral aspect of male genitalia with aedeagus removed.



PTOCHORYCTIS



RHIZOSTHENES

RHIZOSTHENES Meyrick

Rhizosthenes Meyrick, 1935, *in* Caradja and Meyrick, Materialien zu einer Microlepidopteren Fauna der Chinesischen Provinzen Kiangsu, Chekiang und Hunan, 83. (*Typus generis: Rhizosthenes falciformis* Meyrick, ibid. [monotypy].)

Rhizosthenes falciformis Meyrick

Plate 247, Figures 1–1d

Rhizosthenes falciformis Meyrick, 1935, *in* Caradja and Meyrick, Materialien zu einer Microlepidopteren Fauna der Chinesischen Provinzen Kiangsu, Chekiang und Hunan, 83.

". . . ♂, 23 mm. . . . 4 ex. Lungtan 6–14 VI."

Type: The male in the Meyrick collection, "Nanking, China. H. 6.6.33." Slide No. 7641.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

SYNCHALARA Meyrick

Synchalara Meyrick, 1917, Exotic Microlepidoptera, 2: 60. (*Typus generis: Agriophara rhombota* Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 981 [by original designation].)

Synchalara rhombota (Meyrick)

Plate 248, Figures 1–1d

Agriophara rhombota Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 981.

"♂♀. 28–38 mm. . . . Fourteen specimens, Khasi Hills and Silchar, Assam, from June to August (*Antram*). . . ."

Lectotype: ♂, 32 mm., "Khasi Hills, Assam. .7.1906." Slide No. 7580. Four specimens are missing.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.



SYNCHALARA



SYNCHALARA

Synchalara argoplaca (Meyrick)

Plate 249, Figures 1–1b

Agriophara argoplaca Meyrick, 1907, Journ. Bombay Nat. Hist. Soc. 17: 743.

"♂. 25–28 mm. . . . Maskeliya, Ceylon, in March, June and October; three specimens (*de Mowbray, Pole*)."

Lectotype: ♂, dated ".6.05". Slide No. 7581.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Synchalara byrsina (Meyrick)

Plate 249, Figures 2–2b

Agriophara byrsina Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 18: 151.

"♂♀. 22–25 mm. . . . Khasi Hills, from July to September; five specimens."

Lectotype: ♂, 24 mm., "Khasi Hills, Assam. .7.1906." Slide No. 7579.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Synchalara minax (Meyrick)

Plate 249, Figures 3–3b

Agriophara minax Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 18: 152.

"♂. 22–25 mm. . . . Khasi Hills, in July; two specimens."

Lectotype: ♂, 22 mm., "Khasi Hills, Assam. .7.1906." Slide No. 7578.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

THYMIATRIS Meyrick

Thymiatris Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 738. (*Typus generis: Thymiatris melitacma* Meyrick, ibid. [monotypy].)

Thymiatris melitacma Meyrick

Plate 250, Figures 1–1d

Thymiatris melitacma Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 738.

"♂. 42–44 mm. . . . Khasi Hills; two specimens."

Lectotype: ♂, 44 mm., "Khasi Hills, Assam. .5.1906." Slide No. 7648.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.



THYMIATRIS









TRICHERNIS

TRICHERNIS Meyrick

Trichernis Meyrick, 1894, Trans. Ent. Soc. London, 1894: 20. (*Typus generis: Trichernis centrias* Meyrick, ibid. [monotypy].)

Trichernis centrias Meyrick

Plate 251, Figures 1–1d

Trichernis centrias Meyrick, 1894, Trans. Ent. Soc. London, 1894: 20.

"♂. 15 mm. . . . Koni; one specimen."

Type: The male indicated above, "Koni, Burma. NM. .9.88." Slide No. 7610. This genus is synonymous with *Odites* Walsingham.

Figure 1, left wings; 1a, venation of right wings; 1b, anterior aspect of head showing palpi; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia (damaged before dissection) with aedeagus removed.

TRYPHERANTIS Meyrick

Trypherantis Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 740. (*Typus generis: Trypherantis atelogramma* Meyrick, ibid. [monotypy].)

Trypherantis atelogramma Meyrick

Plate 252, Figures 1–1d

Trypherantis atelogramma Meyrick, 1907, Journ. Bombay Nat. Hist. Soc., 17: 740.

"♂. 24–26 mm. . . . Bhotan, Sikkim, in May and June; three specimens (*Dudgeon*)."

Lectotype: ♂, "Sikkim. GCD. 18.6.90." Slide No. 7704. Two specimens are missing.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.







TRYPHERANTIS

COPROMORPHIDAE

1



1a

1b

1c

1d

AEGIDOMORPHA

AEGIDOMORPHA Meyrick

Aegidomorpha Meyrick, 1932, Exotic Microlepidoptera, 4: 228. (*Typus generis: Aegidomorpha psammodina* Meyrick, l.c., 229 [monotypy].)

Aegidomorpha psammodina Meyrick

Plate 253, Figures 1–1d

Aegidomorpha psammodina Meyrick, 1932, Exotic Microlepidoptera, 4: 229.

"♂. 30 mm. . . . China, Szechwan, Mt. Omei, 4,500 feet, August (*Franck*); 1 ex."

Type: The male denoted above, "Omei, China. C. .8.28." Slide No. 9293.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

CATHELOTIS Meyrick

Cathelotis Meyrick, 1926, Exotic Microlepidoptera, 3: 241. (*Typus generis: Cathelotis sanidopa* Meyrick, l.c., 242 [monotypy].)

Cathelotis sanidopa Meyrick

Plate 254, Figures 1–1d

Cathelotis sanidopa Meyrick, 1926, Exotic Microlepidoptera, 3: 242.

"♂. 23 mm. . . . Colombia, Mt. Tolima, 12,500 feet, October; 1 ex. In poor condition, but an interesting new generic type".

Type: The specimen indicated above dated ".10.20". Slide No. 5622.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.



CATHELOTIS

1

1a


1b

1c


1d


2

2a

2b


2c

CHERSOMORPHA

CHERSOMORPHA Meyrick

Chersomorpha Meyrick, 1926, Exotic Microlepidoptera, 3: 243. (*Typus generis: Chersomorpha taospila* Meyrick, ibid. [monotypy].)

Chersomorpha taospila Meyrick

Plate 255, Figures 1–1d

Chersomorpha taospila Meyrick, 1926, Exotic Microlepidoptera, 3: 243.

"♂. 23 mm. . . . New Ireland, January (*Eichhorn*); 1 ex."

Type: The above indicated male, "New Ireland, XII.23–I.24. A. F. Eichhorn." Slide No. 9517.

Figure 1, left wings,; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

Chersomorpha cataxeranta Meyrick

Plate 255, Figures 2–2c

Chersomorpha cataxeranta Meyrick, 1938, Trans. R. Ent. Soc. London, 87: 519.

"♂♀. 16–18 mm. . . . Kokoda, 1,200 feet, May, July. 3 ex. Allied to . . . *neurobapta* Low."

Type: The male so marked in the British Museum, labelled (by Stringer?) "Type ♀". Pin-label data are "Papua: Kokoda. 1,200 ft. 16.V.1933. L. E. Cheesman." A second label bears the inscription "Ch.90". Slide No. 9295. There is a second specimen, undoubtedly another species, labelled "type ♂", which is in poor condition and does not agree with Meyrick's description.

Figure 2, left wings; 2a, male abdomen showing modifications of eighth segment; 2b, lateral view of aedeagus; 2c, ventral view of male genitalia with aedeagus removed.

COPROMORPHA Meyrick

Copromorpha Meyrick, 1886, Trans. Ent. Soc. London, 1886: 281. (*Typus generis: Copromorpha gypsota* Meyrick, l.c., 282 [monotypy].)

Copromorpha gypsota Meyrick

Plate 256, Figures 1–1d

Copromorpha gypsota Meyrick, 1886, Trans. Ent. Soc. London, 1886: 282.

"♂. 26–30 mm. . . . Fiji (*Mathew, Lucas*); two specimens."

Type: Unknown. The species is figured from a specimen from "Fiji, Polynesia, Mathew 1887, 4321." Slide No. 9530.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

Copromorpha bryanthes Meyrick

Plate 256, Figures 2–2b

Copromorpha bryanthes Meyrick, 1926, Exotic Microlepidoptera, 3: 245.

"♀. 27 mm. . . . Borneo, Sarawak, Bidi (*Brooks*); 1 ex."

Type: The above indicated specimen, "Bidi, Sarawak. 1907–1908. C. J. Brooks", erroneously recorded as a female. Slide No. 9292.

Figure 2, left wings; 2a, lateral view of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

1

1a

1b

1c

1d

2

2a

2b

COPROMORPHA

1

1a

1b

2

2a

2b


3


3a

3b


4

4a


4b

COPROMORPHA

Copromorpha cryptochlora Meyrick

Plate 257, Figures 1–1b

Copromorpha cryptochlora Meyrick, 1930, Exotic Microlepidoptera, 3: 590.

"♂. 14 mm. . . . Comoro Is., Grand Comoro (*Humblot*); 1 ex. (Coll. Paravicini). Allied to *aeruginea*."

Type: The above indicated male, so marked in the British Museum, "Grande Comore. L. Humblot. 1884." A small white label bears the inscription "M985". Slide No. 9287.

Figure 1, left wings; 1a, lateral aspect of aedeagus; 1b, ventral view of male genitalia with aedeagus removed.

Copromorpha efflorescens Meyrick

Plate 257, Figures 2–2b

Copromorpha efflorescens Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 412.

"♀. 17–19 mm. . . . Four specimens, Matale and Madulsima, Ceylon, in April and from October to December (*Pole, Vaughan*)."

Lectotype: ♀, 19 mm., "Madulsima, Ceylon. Vaughn. .4.05." Slide No. 9285.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium.

Copromorpha mesobactris Meyrick

Plate 257, Figures 3–3b

Copromorpha mesobactris Meyrick, 1930, Exotic Microlepidoptera, 3: 590.

"♂ 18 mm., ♀ 20 mm. . . . Comoro Is., Mayote, June; 1 ♂. Also 1 ♀ in Coll. Paravicini from Réunion (*Roussel*)."

Type: The female so marked in the British Museum, "Bourbon, Dr. Roussel". A small white label reads "M1025". Slide No. 9288.

Figure 3, left wings; 3a, ventral view of female genitalia; 3b, detail of genital plate and ostium.

Copromorpha metallitis Meyrick

Plate 257, Figures 4–4b

Copromorpha metallitis Meyrick, 1906, Journ. Bombay Nat. Hist. Soc., 17: 411.

"♂♀. 15–19 mm. . . . Five specimens, Puttalam, Ceylon, in April and October (*Pole*)."

Lectotype: ♂, 15 mm., dated ".4.04". Slide No. 9284.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.

Copromorpha myrmecias Meyrick

Plate 258, Figures 1–1b

Copromorpha myrmecias Meyrick, 1930, Exotic Microlepidoptera, 3: 590.

"♀. 16–18 mm. . . . Bihar, Pusa, bred July, September, October from larvae boring top-shoots of *Ficus bengalensis* (*Fletcher*); 4 ex."

Lectotype: ♀, 16 mm., "Pusa, India. TBF. Bred. .7.29." Slide No. 9286.

Figure 1, left wings; 1a, ventral view of female genitalia; 1b, detail of genital plate and ostium.

Copromorpha orthidias Meyrick

Plate 258, Figures 2–2b

Copromorpha orthidias Meyrick, 1927, Insects of Samoa, 3, Lepidoptera, fasc., 2: 96.

"♂. 16 mm. . . . Upolu, Malololelei, 2,000 feet, December; 1 ex. Nearest to the much larger Fijian *C. gypsota*."

Type: The male so marked in the British Museum, "Malololelei, Upolu, Samoa, 2,000 feet. Buxton & Hopkins. Dec. 1925." Slide No. 9290. There is a paired hair-pencil from the intersegmental membrane between segment 8 and the genitalia; eighth segment modified.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Copromorpha pleurophanes Meyrick

Plate 258, Figures 3–3b

Copromorpha pleurophanes Meyrick, 1905, Journ. Bombay Nat. Hist. Soc., 16: 606.

"♂♀. 15–20 mm. . . . Three specimens, Maskeliya, Ceylon, in March (*Pole*); Margherita, Assam."

Lectotype: ♂, 15 mm., "Margherita, Assam. WD. /90." Slide No. 9291. There is a paired hair-pencil from the intersegmental membrane between segment 8 and the genitalia; eighth segment modified.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.

Copromorpha pyrrhoscia Meyrick

Plate 258, Figures 4–4b

Copromorpha pyrrhoscia Meyrick, 1935, Exotic Microlepidoptera, 4: 574.

"♂. 17 mm. . . . Fiji, Vunidawa, August (*H. Phillips*); 1 ex. (Brit. Mus.)."

Type: The male so marked in the British Museum, "Vunidawa, Fiji. 26.8.1932. H. Phillips. 701." A long lavender-coloured label, in Meyrick's hand reads, "*Copromorpha pyrrhoscia* n. sp." Slide No. 9289.

Figure 4, left wings; 4a, lateral aspect of aedeagus; 4b, ventral view of male genitalia with aedeagus removed.


1


1a


1b

2

2a


2b

3

3a

3b


4

4a

4b

COPROMORPHA







NEOPHYLARCHA

NEOPHYLARCHA Meyrick

Neophylarcha Meyrick, 1926, Exotic Microlepidoptera, 3: 240. (*Typus generis: Neophylarcha helicosema* Meyrick, ibid. [monotypy].)

Neophylarcha helicosema Meyrick

Plate 259, Figures 1–1d

Neophylarcha helicosema Meyrick, 1926, Exotic Microlepidoptera, 3: 240.

"♂♀. 15–17 mm. . . . British Guiana, Bartica, January (*Parish*); French Guiana, R. Maroni, July; 2 ex."

Lectotype: The male labelled "St. Jean, R. Maroni, Fr. Guiana. LeM. .7.15." Slide No. 5620.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, lateral aspect of aedeagus; 1d, ventral view of male genitalia with aedeagus removed.

ORDRUPIA Busck

Ordrupia friserella Busck

Plate 260, Figures 1–1c; 2–2b

Ordrupia friserella Busck, 1911, Proc. U.S. Nat. Mus., 40: 228.
Ordrupia fabricata Meyrick, 1915, Trans. Ent. Soc. London, 1915: 215.
Ordrupia dasyleuca Meyrick, 1926, Exotic Microlepidoptera, 3: 242 (new synonymy).

[dasyleuca]

"♂. 34 mm. . . . Peru, Rio Napo, May (*Parish*); 1 ex."

Type: The specimen indicated above dated "5.20". Slide No. 5623. The abdominal tufts in this species are indeciduous.

Figure 1, left wings; 1a, dorsal view of aedeagus; 1b, ventral view of male genitalia with aedeagus removed; 1c, posterior abdominal tufts.

[fabricata]

"♂. 31 mm. . . . British Guiana, R. Demerara; Dutch Guiana, R. Saramacca, in May; two specimens."

Lectotype: ♂, "Saramacca R., Dutch Guiana. R. .5.93." Slide No. 5626.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

Ordrupia macroctenis Meyrick

Plate 260, Figures 3–3b

Ordrupia macroctenis Meyrick, 1926, Exotic Microlepidoptera, 3: 243.

"♂♀. 34–37 mm. . . . Peru, Contamana, R. Ucuyali; 2 ex. This species differs from all others in the much longer antennal pectinations of ♂."

Lectotype: ♂, with ".15" in the lower right of the pin-label. Slide No. 5624. Abdominal tufts readily deciduous.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.



ORDRUPIA

1



1a



1b

1c

1d

2



2a



2b

OSIDRYAS

OSIDRYAS Meyrick

Osidryas Meyrick, 1916, Exotic Microlepidoptera, 2: 7. (*Typus generis: Heterocrita chersodes* Turner, 1913, Proc. Linnean Soc. New South Wales, 38: 223. [*Osidryas* Meyrick is a substitute name for *Heterocrita* Turner, which is preoccupied, and takes the same type.])

Osidryas chersodes (Turner)

Plate 261, Figures 1–1d

Heterocrita chersodes Turner, 1913, Proc. Linnean Soc. New South Wales, 38: 223.

The illustrations are made from a female in the British Museum, "Cairns, Queensland. PPD. .8.04." Slide No. 9581.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, ventral view of female genitalia; 1d, detail of genital plate and ostium.

Osidryas chlorotribes Meyrick

Plate 261, Figures 2–2b

Osidryas chlorotribes Meyrick, 1939, Trans. R. Ent. Soc. London, 89: 59.

"Male, female, 13–14 mm. . . . Java, Telawa, 3 ex., bred March from larvae on leaves of unidentified plant (*Dr. Kalshoven*)."

Lectotype: ♂, 14 mm., "Telawa, Java. K. Bred. .3.35." Slide No. 9296. One specimen is missing.

Figure 2, left wings; 2a, lateral aspect of aedeagus; 2b, ventral view of male genitalia with aedeagus removed.

PHYCOMORPHA Meyrick

Phycomorpha Meyrick, 1914, Trans. Proc. New Zealand Inst., 46: 106. *Typus generis: Phycomorpha metachrysa* Meyrick, ibid. [monotypy].)

Phycomorpha metachrysa Meyrick

Plate 262, Figures 1–1a

Phycomorpha metachrysa Meyrick, 1914, Trans. Proc. New Zealand Inst., 46: 106.

There are three specimens of this species in the British Museum, all without abdomens, one of which I have used to illustrate the wings and wing venation. Slide No. 9587, wings only.

Figure 1, left wings; 1a, wing venation.

Phycomorpha bryophylla Meyrick

Plate 262, Figures 2–2b

Phycomorpha bryophylla Meyrick, 1927, Insects of Samoa, 3, Lepidoptera, fasc., 2: 96.

"♂♀. 19–23 mm. . . . Upolu, Malololelei, 2,000 feet, Apia, 1,000 feet; June, October, December; 4 ex. Extremely similar to *Ph. prasinochroa*, . . ."

Type: The female so marked in the British Museum, "Samoan Is. Upolu Is. Malololelei, 2,000 ft., XII.1925. P. A. Buxton and G. H. Hopkins." Slide No. 9294.

Figure 2, left wings; 2a, ventral view of female genitalia; 2b, detail of genital plate and ostium.

Phycomorpha escharitis Meyrick

Plate 262, Figures 3–3b

Phycomorpha escharitis Meyrick, 1916, Exotic Microlepidoptera, 1: 555.

"♂. 23–29 mm. . . . Colombia, San Antonio, 5,800 feet, in November; four specimens."

Lectotype: One of the above indicated specimens all of which bear identical data and are dated "11.07". Slide No. 5625.

Figure 3, left wings; 3a, lateral aspect of aedeagus; 3b, ventral view of male genitalia with aedeagus removed.



PHYCOMORPHA







RHOPALOSETIA

RHOPALOSETIA Meyrick

Rhopalosetia Meyrick, 1926, Exotic Microlepidoptera, 3: 241. (*Typus generis: Rhopalosetia phlyctaenopa* Meyrick, ibid. [monotypy].)

Rhopalosetia phlyctaenopa Meyrick

Plate 263, Figures 1–1e

Rhopalosetia phlyctaenopa Meyrick, 1926, Exotic Microlepidoptera, 3: 241.

"♂. 24 mm. . . . French Guiana, R. Maroni; 1 ex."

Type: The above indicated specimen, erroneously recorded as a male, labelled "R. Maroni, Fr. Guiana. LeM. .15." Slide No. 5621.

Figure 1, left wings; 1a, venation of right wings; 1b, lateral aspect of head to show palpus; 1c, ventral view of female genitalia; 1d, signum; 1e, detail of genital plate and ostium.

PRESENTED
25 MAY 1955